


# Real Functions of Several Variables - Line Int...

Leif Mejlbro



Download free books at

**bookboon**.com

Leif Mejlbro

---

# Real Functions of Several Variables

## Examples of Line Integrales

Calculus 2c-7

---

---

Real Functions of Several Variables – Examples of Line Integrals – Calculus 2c-7  
© 2007 Leif Mejlbro & Ventus Publishing Aps  
ISBN 978-87-7681-258-4

# Contents

	<b>Preface</b>	<b>5</b>
<b>1</b>	<b>Line integrals, rectangular coordinates</b>	<b>6</b>
<b>2</b>	<b>Line integral, polar coordinates</b>	<b>25</b>
<b>3</b>	<b>Arc lengths and parametric descriptions by the arc length</b>	<b>32</b>
<b>4</b>	<b>Tangential line integrals</b>	<b>57</b>

Please click the advert



## Masters in Management



Designed for high-achieving graduates across all disciplines, London Business School's Masters in Management provides specific and tangible foundations for a successful career in business.

This 12-month, full-time programme is a business qualification with impact. In 2010, our MiM employment rate was 95% within 3 months of graduation\*; the majority of graduates choosing to work in consulting or financial services.

As well as a renowned qualification from a world-class business school, you also gain access to the School's network of more than 34,000 global alumni – a community that offers support and opportunities throughout your career.

For more information visit [www.london.edu/mm](http://www.london.edu/mm), email [mim@london.edu](mailto:mim@london.edu) or give us a call on **+44 (0)20 7000 7573**.

\* Figures taken from London Business School's Masters in Management 2010 employment report

## Preface

In this volume I present some examples of *line integrals*, cf. also *Calculus 2b, Functions of Several Variables*. Since my aim also has been to demonstrate some solution strategy I have as far as possible structured the examples according to the following form

**A** *Awareness*, i.e. a short description of what is the problem.

**D** *Decision*, i.e. a reflection over what should be done with the problem.

**I** *Implementation*, i.e. where all the calculations are made.

**C** *Control*, i.e. a test of the result.

This is an ideal form of a general procedure of solution. It can be used in any situation and it is not linked to Mathematics alone. I learned it many years ago in the Theory of Telecommunication in a situation which did not contain Mathematics at all. The student is recommended to use it also in other disciplines.

One is used to from high school immediately to proceed to **I. Implementation**. However, examples and problems at university level are often so complicated that it in general will be a good investment also to spend some time on the first two points above in order to be absolutely certain of what to do in a particular case. Note that the first three points, **ADI**, can always be performed.

This is unfortunately not the case with **C Control**, because it from now on may be difficult, if possible, to check one's solution. It is only an extra securing whenever it is possible, but we cannot include it always in our solution form above.

I shall on purpose not use the logical signs. These should in general be avoided in Calculus as a shorthand, because they are often (too often, I would say) misused. Instead of  $\wedge$  I shall either write "and", or a comma, and instead of  $\vee$  I shall write "or". The arrows  $\Rightarrow$  and  $\Leftrightarrow$  are in particular misunderstood by the students, so they should be totally avoided. Instead, write in a plain language what you mean or want to do.

It is my hope that these examples, of which many are treated in more ways to show that the solutions procedures are not unique, may be of some inspiration for the students who have just started their studies at the universities.

Finally, even if I have tried to write as careful as possible, I doubt that all errors have been removed. I hope that the reader will forgive me the unavoidable errors.

Leif Mejlbro  
14th October 2007

# 1 Line integrals, rectangular coordinates

**Example 1.1** Calculate in each of the following cases the given line integral, where the curve  $\mathcal{K}$  is given by the parametric description

$$\mathcal{K} = \{\mathbf{x} \in \mathbb{R}^k \mid \mathbf{x} = \mathbf{r}(t), \quad t \in I\}, \quad k = 2 \text{ or } k = 3.$$

1) The line integral  $\int_{\mathcal{K}} ds$ , where

$$\mathbf{r}(t) = (a(1 - \cos t), a(t - \sin t)), \quad t \in [0, 4\pi].$$

2) The line integral  $\int_{\mathcal{K}} \sqrt{x} ds$ , where

$$\mathbf{r}(t) = (a(1 - \cos t), a(t - \sin t)), \quad t \in [0, 4\pi].$$

3) The line integral  $\int_{\mathcal{K}} z ds$ , where

$$\mathbf{r}(t) = (t, 3t^2, 6t^3), \quad t \in [0, 2].$$

4) The line integral  $\int_{\mathcal{K}} \frac{1}{1+6y} ds$ , where

$$\mathbf{r}(t) = (t, 3t^2, 6t^3), \quad t \in [0, 2].$$

5) The line integral  $\int_{\mathcal{K}} (x + e^z)$ , where

$$\mathbf{r}(t) = (\cos t, \sin t, \ln \cos t), \quad t \in \left[0, \frac{\pi}{4}\right].$$

[Cf. **Example 3.3.6.**]

6) The line integral  $\int_{\mathcal{K}} (x^2 + y^2 + z^2) ds$ , where

$$\mathbf{r}(t) = (e^t \cos t, e^t \sin t, e^t), \quad t \in [0, 2].$$

7) The line integral  $\int_{\mathcal{K}} \frac{x+y}{z^2} ds$ , where

$$\mathbf{r}(t) = \frac{1}{\sqrt{3}} (e^t, e^t \sin t, e^t), \quad t \in [0, u].$$

[Cf. **Example 3.3.7.**]

8) The line integral  $\int_{\mathcal{K}} (x^2 + y^2) ds$ , where

$$\mathbf{r}(t) = \left( \frac{1-t^2}{1+t^2}, \frac{2t}{1+t^2} \right), \quad t \in \mathbb{R}.$$

9) The line integral  $\int_{\mathcal{K}} ds$ , where

$$\mathbf{r}(t) = \left( 2 \operatorname{Arcsin} t, \ln(1-t^2), \ln \frac{1+t}{1-t} \right), \quad t \in \left[0, \frac{1}{\sqrt{2}}\right].$$

10) The line integral  $\int_K x e^y ds$ , where

$$\mathbf{r}(t) = \left( 2 \operatorname{Arccsin} t, \ln(1 - t^2), \ln \frac{1+t}{1-t} \right), \quad t \in \left[ 0, \frac{1}{\sqrt{2}} \right].$$

11) The line integral  $\int_K \frac{1}{\sqrt{1+3x^2+z^2}} ds$ , where

$$\mathbf{r}(t) = (\cos t, 2 \sin t, e^t), \quad t \in [-1, 1].$$

**A** Line integrals.

**D** First find  $\|\mathbf{r}'(t)\|$  in each case. Then compute the line integral.

Please click the advert



You're full of *energy*  
and *ideas*. And that's  
just what we are looking for.

Looking for a career where your ideas could really make a difference? UBS's Graduate Programme and internships are a chance for you to experience for yourself what it's like to be part of a global team that rewards your input and believes in succeeding together.

Wherever you are in your academic career, make your future a part of ours by visiting [www.ubs.com/graduates](http://www.ubs.com/graduates).

[www.ubs.com/graduates](http://www.ubs.com/graduates)



© UBS 2010. All rights reserved.




Figure 1: The plane curve  $\mathcal{K}$  of **Example 1.1.1** and **Example 1.1.2** for  $a = 1$ .

I 1) Here,

$$\mathbf{r}'(t) = a(\sin t, 1 - \cos t),$$

so

$$\|\mathbf{r}'(t)\| = a\sqrt{\sin^2 t + (1 - \cos t)^2} = a\sqrt{2 - 2\cos t} = a\sqrt{4\sin^2 \frac{t}{2}} = 2a \left| \sin \frac{t}{2} \right|.$$

Then accordingly,

$$\int_{\mathcal{K}} ds = \int_0^{4\pi} 2a \left| \sin \frac{t}{2} \right| dt = 4a \int_0^{2\pi} |\sin u| du = 8a \int_0^{\pi} \sin u du = 16a.$$

2) It follows from 1) that

$$\|\mathbf{r}'(t)\| = a\sqrt{2(1 - \cos t)},$$

thus

$$\begin{aligned} \int_{\mathcal{K}} \sqrt{x} ds &= \int_0^{4\pi} \sqrt{a(1 - \cos t)} \cdot a\sqrt{2(1 - \cos t)} dt \\ &= a\sqrt{2a} \int_0^{4\pi} |1 - \cos t| dt = a\sqrt{2a} \int_0^{4\pi} (1 - \cos t) dt = 4\sqrt{2}\pi a\sqrt{a}. \end{aligned}$$

3) It follows from  $\mathbf{r}'(t) = (1, 6t, 18t^2)$  that

$$\|\mathbf{r}'(t)\| = \sqrt{1 + 36t^2 + 324t^4} = \sqrt{(1 + 18t^2)^2} = 1 + 18t^2,$$

hence

$$\int_{\mathcal{K}} z ds = \int_0^2 6t^3(1 + 18t^2) dt = \left[ \frac{6}{4}t^4 + \frac{6 \cdot 18}{6}t^6 \right]_0^2 = \frac{3}{2} \cdot 16 + 18 \cdot 64 = 24 + 1152 = 1176.$$






Figure 2: The curve  $\mathcal{K}$  for  $t \in [0, \frac{7}{10}]$ . It is used in **Example 1.1.3** and **Example 1.1.4** for  $t \in [0, 2]$ .

4) It follows from 3 above that  $\|\mathbf{r}'(t)\| = 1 + 18t^2$ , so

$$\int_{\mathcal{K}} \frac{1}{1+6y} ds = \int_0^2 \frac{1+18t^2}{1+18t^2} dt = 2.$$




Figure 3: The curve  $\mathcal{K}$  of **Example 1.1.5**.

5) It follows from

$$\mathbf{r}'(t) = \left( -\sin t, \cos t, -\frac{\sin t}{\cos t} \right),$$

that

$$\|\mathbf{r}'(t)\| = \sqrt{\sin^2 t + \cos^2 t + \frac{\sin^2 t}{\cos^2 t}} = \sqrt{1 + \frac{\sin^2 t}{\cos^2 t}} = \frac{1}{|\cos t|},$$

hence

$$\int_{\mathcal{K}} (x + e^z) ds = \int_0^{\frac{\pi}{4}} (\cos t + e^{\ln \cos t}) \frac{1}{\cos t} dt = \int_0^{\frac{\pi}{4}} 2 dt = \frac{\pi}{2}.$$




Figure 4: The curve  $\mathcal{K}$  of **Example 1.1.6** and – apart from e factor  $1/\sqrt{3}$  – of **Example 1.1.7**.

6) It follows from

$$\mathbf{r}'(t) = (e^t(\cos t - \sin t), e^t(\sin t + \cos t), e^t),$$

that

$$\|\mathbf{r}'(t)\| = e^t \sqrt{(\cos t - \sin t)^2 + (\sin t + \cos t)^2 + 1} = \sqrt{3} e^t,$$

thus

$$\begin{aligned} \int_{\mathcal{K}} (x^2 + y^2 + z^2) ds &= \int_0^2 e^{2t} (\cos^2 t + \sin^2 t + 1) \cdot \sqrt{3} e^t dt \\ &= 2\sqrt{3} \int_0^2 e^{3t} dt = \frac{2\sqrt{3}}{3} (e^6 - 1). \end{aligned}$$

7) If we first divide by  $\sqrt{3}$ , we get by **Example 1.1.6** the more nice expression  $\|\mathbf{r}'(t)\| = e^t$ . Then the line integral becomes

$$\int_{\mathcal{K}} ds = \int_0^u e^t dt = e^u - 1.$$

8) We get by just computing

$$\begin{aligned} \mathbf{r}'(t) &= \left( \frac{-2t(1+t^2) - 2t(1-t^2)}{(1+t^2)^2}, \frac{2(1+t^2) - 2t \cdot 2t}{(1+t^2)^2} \right) \\ &= \left( -\frac{4t}{(1+t^2)^2}, \frac{2(1-t^2)}{(1+t^2)^2} \right) = \frac{1}{(1+t^2)^2} (-2t, 1-t^2), \end{aligned}$$

hence

$$\|\mathbf{r}'(t)\| = \frac{1}{(1+t^2)^2} \sqrt{4t^2 + (1-t^2)^2} = \frac{1}{(1+t^2)^2} \sqrt{(1+t^2)^2} = \frac{1}{1+t^2}.$$

Then finally,

$$\begin{aligned} \int_{\mathcal{K}} (x^2 + y^2) ds &= \int_{-\infty}^{+\infty} \frac{1}{(1+t^2)^2} \{ (1-t^2)^2 + 4t^2 \} \cdot \frac{2}{1+t^2} dt \\ &= \int_{-\infty}^{+\infty} \frac{(1+t^2)^2}{(1+t^2)^2} \cdot \frac{2}{1+t^2} dt = [2 \operatorname{Arctan} t]_{-\infty}^{+\infty} = 2\pi. \end{aligned}$$




Figure 5: The curve  $\mathcal{K}$  of **Example 1.1.8**, i.e. a circle except for the point  $(-1, 0)$ .

ALTERNATIVE. The computation above was a little elaborated. However, the line integral is independent of the chosen parametric description, and  $\mathcal{K}$  is a circle with the exception of the point  $(-1, 0)$ , which is of no importance for the integration. Therefore, we can apply the simpler parametric description

$$\mathbf{r}(t) = (\cos t, \sin t), \quad t \in ]-\pi, \pi[ ,$$

where

$$\mathbf{r}'(t) = (-\sin t, \cos t) \quad \text{og} \quad \|\mathbf{r}'(t)\| = \sqrt{\sin^2 t + \cos^2 t} = 1.$$

Then the line integral becomes almost trivial,

$$\int_{\mathcal{K}} (x^2 + y^2) ds = \int_{-\pi}^{\pi} 1^2 dt = 2\pi.$$




Figure 6: The curve  $\mathcal{K}$  of **Example 1.1.9** and **Example 1.1.10**.

9) Here

$$\mathbf{r}'(t) = \left( \frac{2}{\sqrt{1-t^2}}, -\frac{2t}{1-t^2}, \frac{1}{1+t} + \frac{1}{1-t} \right) = \frac{2}{1-t^2} \left( \sqrt{1-t^2}, -t, 1 \right),$$

hence

$$\|\mathbf{r}'(t)\| = \frac{2}{1-t^2} \sqrt{1-t^2+t^2+1} = \frac{2\sqrt{2}}{1-t^2} = \sqrt{2} \left( \frac{1}{1+t} - \frac{1}{t-1} \right).$$

The line integral is

$$\begin{aligned} \int_{\mathcal{K}} ds &= \int_0^{\frac{1}{\sqrt{2}}} \sqrt{2} \left( \frac{1}{1+t} - \frac{1}{t-1} \right) dt = \sqrt{2} \left[ \ln \frac{1-t}{1+t} \right]_0^{\frac{1}{\sqrt{2}}} \\ &= \sqrt{2} \ln \left( \frac{1+\frac{1}{\sqrt{2}}}{1-\frac{1}{\sqrt{2}}} \right) = \sqrt{2} \ln \left( \frac{\sqrt{2}+1}{\sqrt{2}-1} \right) = 2\sqrt{2} \ln(\sqrt{2}+1). \end{aligned}$$

10) We consider the same curve as in **Example 1.1.9**, so we can reuse that

$$\|\mathbf{r}'(t)\| = \frac{2\sqrt{2}}{1-t^2} = \sqrt{2} \left( \frac{1}{1+t} - \frac{1}{t-1} \right), \quad t \in \left[ 0, \frac{1}{\sqrt{2}} \right],$$

Please click the advert



Discover the truth at [www.deloitte.ca/careers](http://www.deloitte.ca/careers)

**Deloitte.**

© Deloitte & Touche LLP and affiliated entities.

Download free ebooks at [bookboon.com](http://bookboon.com)

and the line integral becomes

$$\begin{aligned}
 \int_{\mathcal{K}} x e^y ds &= \int_0^{\frac{1}{\sqrt{2}}} 2 \operatorname{Arctan} t \cdot (1-t^2) \cdot \frac{2\sqrt{2}}{1-t^2} dt = 4\sqrt{2} \int_0^{\frac{1}{\sqrt{2}}} \operatorname{Arctan} t dt \\
 &= 4\sqrt{2} \int_0^{\frac{\pi}{4}} u \cos u du = 4\sqrt{2} [u \sin u + \cos u]_0^{\frac{\pi}{4}} = 4\sqrt{2} \left\{ \frac{\pi}{4} \cdot \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} - 1 \right\} \\
 &= 4 + \pi - 4\sqrt{2} = \pi - 4(\sqrt{2} - 1).
 \end{aligned}$$




Figure 7: The curve  $\mathcal{K}$  of **Example 1.1.11**.

11) Here

$$\mathbf{r}'(t) = (-\sin t, 2 \cos t, e^t),$$

so

$$\|\mathbf{r}'(t)\| = \sqrt{\sin^2 t + 4 \cos^2 t + e^{2t}} = \sqrt{1 + 3 \cos^2 t + e^{2t}}.$$

The parametric description of the integrand restricted to the curve is

$$\sqrt{1 + 3x^2 + z^2} = \sqrt{1 + 3 \cos^2 t + e^{2t}},$$

so the line integral becomes easy to compute

$$\int_{\mathcal{K}} \frac{1}{\sqrt{1 + 3x^2 + z^2}} ds = \int_{-1}^1 \frac{1}{\sqrt{1 + 3 \cos^2 t + e^{2t}}} \sqrt{1 + 3 \cos^2 t + e^{2t}} dt = \int_{-1}^1 1 dt = 2.$$

**Example 1.2** Calculate in each of the following cases the given line integral along the given plane curve  $\mathcal{K}$  of the equation  $y = Y(x)$ ,  $x \in I$ .

1) The line integral  $\int_{\mathcal{K}} x^2 ds$  along the curve

$$y = Y(x) = \ln x, \quad x \in [1, 2\sqrt{2}].$$

2) The line integral  $\int_{\mathcal{K}} \frac{1}{1+4y} ds$  along the curve

$$y = Y(x) = x^2, \quad x \in [0, 1].$$

3) The line integral  $\int_{\mathcal{K}} y^2 ds$  along the curve

$$y = Y(x) = x, \quad x \in [1, 2].$$

4) The line integral  $\int_{\mathcal{K}} \frac{1}{\sqrt{2-y^2}} ds$  along the curve

$$y = Y(x) = \sin x, \quad x \in [0, \pi].$$

5) The line integral  $\int_{\mathcal{K}} \frac{1}{\sqrt{2+y^2}} ds$  along the curve

$$y = Y(x) = \sinh x, \quad x \in [0, 2].$$

6) The line integral  $\int_{\mathcal{K}} y e^x ds$  along the curve

$$y = Y(x) = e^x, \quad x \in [0, 1].$$

**A** Line integrals along plane curves.

**D** Sketch if possible the den plane curve. Compute the weight function  $\sqrt{1+Y'(x)^2}$  and finally reduce the line integral to an ordinary integral.




Figure 8: The curve  $\mathcal{K}$  of **Example 1.2.1**.

**I** 1) It follows from  $Y'(x) = \frac{1}{x}$  that

$$\sqrt{1+Y'(x)^2} = \frac{1}{x} \sqrt{1+x^2}, \quad x \in [1, 2\sqrt{2}].$$

Thus we get the line integral


$$\begin{aligned}\int_{\mathcal{K}} x^2 ds &= \int_1^{2\sqrt{2}} x^2 \cdot \frac{1}{x^2} \sqrt{1+x^2} dx = \int_0^{2\sqrt{2}} \sqrt{1+x^2} \cdot x dx \\ &= \left[ \frac{1}{2} \cdot \frac{2}{3} (1+x^2)^{\frac{3}{2}} \right]_1^{2\sqrt{2}} = \frac{1}{3} \left\{ 9^{\frac{3}{2}} - 2^{\frac{3}{2}} \right\} = \frac{1}{3} (27 - 2\sqrt{2}) = 9 - \frac{2}{3} \sqrt{2}.\end{aligned}$$

Please click the advert

It's only an opportunity if you act on it

[IKEA.SE/STUDENT](http://IKEA.SE/STUDENT)

© Inter IKEA Systems B.V. 2009


Figure 9: The curve  $\mathcal{K}$  of **Example 1.2.2**.

2) From  $Y'(x) = 2x$  follows that

$$\sqrt{1 + Y'(x)^2} = \sqrt{1 + 4x^2},$$

and thus

$$\begin{aligned} \int_{\mathcal{K}} \frac{1}{1+4y} ds &= \int_0^1 \frac{\sqrt{1+4x^2}}{1+4x^2} dx = \int_0^1 \frac{1}{\sqrt{1+(2x)^2}} dx = \frac{1}{2} \int_0^2 \frac{1}{\sqrt{1+t^2}} dt \\ &= \frac{1}{2} \left[ \ln(t + \sqrt{1+t^2}) \right]_0^2 = \frac{1}{2} \ln\left(\frac{2+\sqrt{5}}{1}\right) = \frac{1}{2} \ln(2+\sqrt{5}). \end{aligned}$$

Figure 10: The curve  $\mathcal{K}$  of **Example 1.2.3**.


3) Here clearly  $\sqrt{1 + Y'(x)^2} = \sqrt{1 + 1^2} = \sqrt{2}$ , so

$$\int_{\mathcal{K}} y^2 ds = \int_1^2 x^2 \sqrt{2} ds = \frac{\sqrt{2}}{3} [x^3]_1^2 = \frac{7}{3} \sqrt{2}.$$

4) We get by differentiation of  $Y(x) = \sin x$  that  $Y'(x) = \cos x$ , hence the weight function is


$$\sqrt{1 + Y'(x)^2} = \sqrt{1 + \cos^2 x} = \sqrt{2 - \sin^2 x}.$$



Figure 11: The curve  $\mathcal{K}$  of **Example 1.2.4**.

We finally get the line integral by insertion

$$\int_{\mathcal{K}} \frac{1}{\sqrt{2-y^2}} ds = \int_0^{\pi} \frac{1}{\sqrt{2-\sin^2 x}} \sqrt{2-\sin^2 x} dx = \int_0^{\pi} dx = \pi.$$

Figure 12: The curve  $\mathcal{K}$  of **Example 1.2.5**.

- 5) When  $Y(x) = \sinh x$ , then  $Y'(x) = \cosh x$ , and the weight becomes


$$\sqrt{1+Y'(x)^2} = \sqrt{1+\cosh^2 x} = \sqrt{2+\sinh^2 x}.$$

We get finally the line integral by insertion

$$\int_{\mathcal{K}} \frac{1}{\sqrt{2+y^2}} ds = \int_0^2 \frac{1}{\sqrt{2+\sinh^2 x}} \sqrt{2+\sinh^2 x} dx = \int_0^2 dx = 2.$$

- 6) When  $Y(x) = e^x$  then also  $Y'(x) = e^x$ , so the weight function becomes

$$\sqrt{1+Y'(x)^2} = \sqrt{1+e^{2x}}.$$

Figure 13: The curve  $\mathcal{K}$  of **Example 1.2.6**.

We get the line integral by insertion

$$\begin{aligned} \int_{\mathcal{K}} y e^x ds &= \int_0^1 e^x \cdot e^x \cdot \sqrt{1+e^{2x}} dx = \int_0^1 \sqrt{1+e^{2x}} \cdot e^{2x} dx \\ &= \frac{1}{2} \int_{x=0}^1 \sqrt{1+e^{2x}} d(1+e^{2x}) = \frac{1}{2} \cdot \frac{2}{3} \left[ (1+e^{2x})^{\frac{3}{2}} \right]_0^1 = \frac{1}{3} \left\{ (1+e^2)^{\frac{3}{2}} - 1 \right\}. \end{aligned}$$

Please click the advert

## YOUR CHANCE TO CHANGE THE WORLD

Here at Ericsson we have a deep rooted belief that the innovations we make on a daily basis can have a profound effect on making the world a better place for people, business and society. Join us.

In Germany we are especially looking for graduates as Integration Engineers for

- Radio Access and IP Networks
- IMS and IPTV

We are looking forward to getting your application!  
To apply and for all current job openings please visit our web page: [www.ericsson.com/careers](http://www.ericsson.com/careers)

ericsson.  
com



**Example 1.3** A space curve  $\mathcal{K}$  is given by the parametric description

$$\mathbf{r}(t) = (e^t, t\sqrt{2}, e^{-t}), \quad t \in [0, \ln 3].$$

Prove that the curve element  $ds$  is given by  $(e^t + e^{-t}) dt$ , and then find the line integral

$$\int_{\mathcal{K}} x^3 z ds.$$

**A** Curve element and line integral.

**D** Follow the guidelines.




Figure 14: The curve  $\mathcal{K}$ .

**I** The curve is clearly of class  $C^\infty$ . Furthermore,

$$\|\mathbf{r}'(t)\|^2 = (e^t)^2 + (\sqrt{2})^2 + (e^{-t})^2 = e^{2t} + 2 + e^{-2t} = (e^t + e^{-t})^2,$$

and we get the curve element

$$ds = \|\mathbf{r}'(t)\| dt = (e^t + e^{-t}) dt$$

with respect to the given parametric description.

Then compute the line integral,

$$\begin{aligned} \int_{\mathcal{K}} x^3 z ds &= \int_0^{\ln 3} e^{3t} e^{-t} (e^t + e^{-t}) dt = \int_0^{\ln 3} (e^{3t} + e^t) dt \\ &= \left[ \frac{1}{3} e^{3t} + e^t \right]_0^{\ln 3} = \frac{1}{3} \cdot 3^3 + 3 - \frac{1}{3} - 1 = \frac{32}{3}, \end{aligned}$$

where we alternatively first can apply the change of variables  $u = e^t$ , from which

$$\int_{\mathcal{K}} x^3 z ds = \int_1^3 (u^2 + 1) du = \left[ \frac{1}{3} u^3 + u \right]_1^3 = 9 + 3 - \frac{1}{3} - 1 = \frac{32}{3}.$$

**Example 1.4** A space curve  $\mathcal{K}$  is given by the parametric description

$$\mathbf{r}(t) = \left( t + 4 \cos t, \frac{4}{3}t - 3 \cos t, 5 \sin t \right), \quad t \in \left[ 0, \frac{\pi}{2} \right].$$

Find the value of the line integral

$$\int_{\mathcal{K}} x \, ds.$$

**A** Line integral.

**D** First find the curve element  $ds = \|\mathbf{r}'(t)\| \, dt$ .




Figure 15: The space curve  $\mathcal{K}$ .

**I** From

$$\mathbf{r}'(t) = \left( 1 - 4 \sin t, \frac{4}{3} + 3 \sin t, 5 \cos t \right),$$

follows that

$$\begin{aligned} \|\mathbf{r}'(t)\|^2 &= (1 - 4 \sin t)^2 + \left( \frac{4}{3} + 3 \sin t \right)^2 + 25 \cos^2 t \\ &= 1 - 8 \sin t + 16 \sin^2 t + \frac{16}{9} + 8 \sin t + 9 \sin^2 t + 25 \cos^2 t = \frac{25}{9} + 25 = \frac{25}{9} \cdot 10, \end{aligned}$$

thus

$$ds = \|\mathbf{r}'(t)\| \, dt = \frac{5}{3} \sqrt{10} \, dt.$$

The line integral is

$$\begin{aligned} \int_{\mathcal{K}} x \, ds &= \int_0^{\frac{\pi}{2}} (t + 4 \cos t) \cdot \frac{5}{3} \sqrt{10} \, dt = \frac{5}{3} \sqrt{10} \left[ \frac{t^2}{2} + 4 \sin t \right]_0^{\frac{\pi}{2}} = \frac{5}{3} \sqrt{10} \left( \frac{\pi^2}{8} + 4 \right) \\ &= \frac{5\pi^2 \sqrt{10}}{24} + \frac{20\sqrt{10}}{3}. \end{aligned}$$

**Example 1.5** A space curve  $\mathcal{K}$  is given by the parametric description

$$\mathbf{r}(t) = \left( t^2, e^{2t}, \frac{1}{2} t^4 \right), \quad t \in [1, 2].$$

Find the value of the line integral

$$\int_{\mathcal{K}} \frac{1}{\sqrt{x + 2xz + y^2}} ds.$$

**A** Line integral.

**D** First calculate the weight function  $\|\mathbf{r}'(t)\|$ .




Figure 16: The curve  $\mathcal{K}$ . Notice the different scales on the axes.

**I** We get from

$$\mathbf{r}'(t) = (2t, 2e^{2t}, 2t^3) = 2(t, e^{2t}, t^3)$$

that

$$\|\mathbf{r}'(t)\| = 2\sqrt{e^{4t} + t^2 + t^6}.$$

Then by insertion and reduction,

$$\int_{\mathcal{K}} \frac{1}{\sqrt{x + 2xz + y^2}} ds = \int_1^2 \frac{1}{\sqrt{t^2 + 2t^2 \cdot \frac{1}{2} t^4 + e^{4t}}} \cdot 2\sqrt{e^{4t} + t^2 + t^6} dt = 2.$$

**Example 1.6** A space curve  $\mathcal{K}$  is given by the parametric description

$$\mathbf{r}(t) = \left( \ln t, t^2, \frac{1}{2} t^4 \right), \quad t \in [1, 2].$$

Prove that the curve element  $ds$  is given by  $\left( \frac{1}{t} + 2t^3 \right) dt$ , and then compute the line integral

$$\int_{\mathcal{K}} \frac{y e^x}{z} ds.$$

**A** Line integral.

**D** Follow the guidelines.




Figure 17: The curve  $\mathcal{K}$ .

**I** Clearly,  $\mathbf{r}(t)$  is of class  $C^\infty$  for  $t \in ]1, 2[$ . Then

$$\mathbf{r}'(t) = \left( \frac{1}{t}, 2t, 2t^3 \right), \quad t \in ]1, 2[,$$

implies that

$$\|\mathbf{r}'(t)\|^2 = \frac{1}{t^2} + 4t^2 + 4t^6 = \left( \frac{1}{t} + 2t^3 \right)^2,$$

so

$$ds = \|\mathbf{r}'(t)\| dt = \left| \frac{1}{t} + 2t^3 \right| dt = \left( \frac{1}{t} + 2t^3 \right) dt \quad \text{for } t \in ]1, 2[.$$

We get by insertion of the parametric description,

$$\begin{aligned} \int_{\mathcal{K}} \frac{y e^x}{z} ds &= \int_1^2 \frac{t^2 \cdot t}{\frac{1}{2} t^4} \cdot \left( \frac{1}{t} + 2t^3 \right) dt = 2 \int_1^2 \left( \frac{1}{t^2} + 2t^2 \right) dt \\ &= 2 \left[ -\frac{1}{t} + \frac{2}{3} t^3 \right]_1^2 = 2 \left( -\frac{1}{2} + \frac{16}{3} + 1 - \frac{2}{3} \right) = \frac{31}{3}. \end{aligned}$$

**Example 1.7** A space curve  $\mathcal{K}$  is given by the parametric description

$$\mathbf{r}(t) = (\ln t, t^2, 2t), \quad t \in \left[\frac{1}{2}, \frac{3}{2}\right].$$

1) Find a parametric description of the tangent to  $\mathcal{K}$  at the point  $\mathbf{r}(1)$ .

2) Prove that the curve element  $ds$  is given by  $\left(\frac{1}{t} + 2t\right) dt$ .

3) Compute the value of the line integral

$$\int_{\mathcal{K}} (e^x + \sqrt{y} + 2z) ds.$$

**A** Space curve; tangent; curve element; line integral.

**D** Find  $\mathbf{r}'(t)$ , and apply that  $ds = \|\mathbf{r}'(t)\| dt$ .




Figure 18: The curve  $\mathcal{K}$  and its tangent at  $(0, 1, 2)$ .

**I** 1) Since  $\mathbf{r}(1) = (\ln 1, 1^2, 2 \cdot 1) = (0, 1, 2)$ , and

$$\mathbf{r}'(t) = \left(\frac{1}{t}, 2t, 2\right), \quad \mathbf{r}'(1) = (1, 2, 2),$$

a parametric description of the tangent is given by

$$(x(u), y(u), z(u)) = (0, 1, 2) + u(1, 2, 2) = (u, 2u + 1, 2u + 2), \quad u \in \mathbb{R}.$$

2) Since

$$\|\mathbf{r}'(t)\|^2 = \frac{1}{t^2} + 4t^2 + 4 = \left(\frac{1}{t} + 2t\right)^2,$$

we get for  $t \in \left[\frac{1}{2}, \frac{3}{2}\right]$  that

$$ds = \|\mathbf{r}'(t)\| dt = \left|\frac{1}{t} + 2t\right| dt = \left(\frac{1}{t} + 2t\right) dt.$$

3) Then by insertion and computation we get the line integral

$$\begin{aligned}
 \int_{\mathcal{K}} (e^x + \sqrt{y} + 2z) \, ds &= \int_{\frac{1}{2}}^{\frac{3}{2}} \left( e^{\ln t} + \sqrt{t^2} + 2 \cdot 2t \right) \cdot \left( \frac{1}{t} + 2t \right) dt \\
 &= \int_{\frac{1}{2}}^{\frac{3}{2}} (t + t + 4t) \cdot \frac{1}{t} (1 + 2t^2) dt = 6 \int_{\frac{1}{2}}^{\frac{3}{2}} (1 + 2t^2) dt \\
 &= [6t + 4t^3]_{\frac{1}{2}}^{\frac{3}{2}} = 6 + 4 \left\{ \left( \frac{3}{2} \right)^3 - \left( \frac{1}{2} \right)^3 \right\} = 6 + \frac{1}{2} (27 - 1) = 19.
 \end{aligned}$$

Please click the advert

SIMPLY CLEVER

ŠKODA



We will turn your CV into  
an opportunity of a lifetime



Do you like cars? Would you like to be a part of a successful brand?  
We will appreciate and reward both your enthusiasm and talent.  
Send us your CV. You will be surprised where it can take you.

Send us your CV on  
[www.employerforlife.com](http://www.employerforlife.com)





## 2 Line integral, polar coordinates

**Example 2.1** Compute in each of the following cases the given line integral along the plane curve  $\mathcal{K}$  which is given by an equation in polar coordinates.

- 1) The line integral  $\int_{\mathcal{K}} (x^2 + y^2) ds$  along the curve given by

$$\varrho = e^\varphi, \quad \varphi \in [0, 4].$$

- 2) The line integral  $\int_{\mathcal{K}} y ds$  along the curve given by

$$\varrho = a(1 - \cos \varphi), \quad \varphi \in [0, \pi].$$

- 3) The line integral  $\int_{\mathcal{K}} \sqrt{y} ds$  along the curve given by

$$\varphi = \operatorname{Arcsin} \varrho, \quad \varrho \in [0, 1].$$

- 4) The line integral  $\int_{\mathcal{K}} \frac{y}{\sqrt{4a-3\varrho}} ds$  along the curve given by

$$\varrho = a \cos^2 \varphi, \quad \varphi \in \left[0, \frac{\pi}{2}\right].$$

- 5) The line integral  $\int_{\mathcal{K}} \frac{1}{x^2 + y^2} ds$  along the curve given by

$$\varrho = \frac{a}{\cos \varphi}, \quad \varphi \in \left[0, \frac{\pi}{4}\right].$$

- 6) The line integral  $\int_{\mathcal{K}} \left(\sqrt{x^2 + y^2} - 1\right) ds$  along the curve given by

$$\varphi = \varrho - \ln \varrho, \quad \varrho \in [1, 2].$$

- 7) The line integral  $\int_{\mathcal{K}} (x^2 + y^2) ds$  along the curve given by

$$\varphi = \varrho, \quad \varrho \in [1, 2].$$

**A** Line integral in polar coordinates.

**D** First compute the weight function  $\sqrt{\varrho^2 + \left(\frac{d\varrho}{d\varphi}\right)^2}$  (possibly  $\sqrt{1 + \left(\varrho \frac{d\varphi}{d\varrho}\right)^2}$ ), and then the line integral.

**I** 1) From  $\frac{d\varrho}{d\varphi} = e^\varphi = \varrho$  follows that

$$\sqrt{\varrho^2 + \left(\frac{d\varrho}{d\varphi}\right)^2} = \sqrt{2} \cdot \varrho = \sqrt{2} \cdot e^\varphi,$$

and thus

$$\int_{\mathcal{K}} (x^2 + y^2) ds = \int_0^4 \varrho^2 \sqrt{2} \cdot e^\varphi d\varphi = \sqrt{2} \int_0^4 e^{3\varphi} d\varphi = \frac{\sqrt{2}}{3} (e^{12} - 1).$$




Figure 19: The curve  $\mathcal{K}$  of **Example 2.1.1**.





Figure 20: The curve  $\mathcal{K}$  of **Example 2.1.2**.

Figure 21: The curve  $\mathcal{K}$  of **Example 2.1.3**.

2) From  $\frac{d\rho}{d\varphi} = a \sin \varphi$ , follows that

$$\rho^2 + \left( \frac{d\rho}{d\varphi} \right)^2 = a^2 \{ (1 - \cos \varphi)^2 + \sin^2 \varphi \} = a^2 \cdot 2(1 - \cos \varphi),$$


Please click the advert

With us you can  
shape the future.  
Every single day.

For more information go to:  
[www.eon-career.com](http://www.eon-career.com)

Your energy shapes the future.

**e-on**

Figure 22: The curve  $\mathcal{K}$  of **Example 2.1.4**.

hence

$$\begin{aligned} \int_{\mathcal{K}} y \, ds &= \int_0^{\pi} \varrho(\varphi) \sin \varphi \cdot a\sqrt{2} \cdot \sqrt{1 - \cos \varphi} \, d\varphi = a^2\sqrt{2} \int_0^{\pi} (1 - \cos \varphi)^{\frac{3}{2}} \sin \varphi \, d\varphi \\ &= a^2\sqrt{2} \left[ \frac{2}{5}(1 - \cos \varphi)^{\frac{5}{2}} \right]_0^{\pi} = a^2\sqrt{2} \cdot \frac{2}{5} \cdot 2^{\frac{5}{2}} = \frac{16a^2}{5}. \end{aligned}$$

3) It follows from

$$\sqrt{1 + \left\{ \varrho \frac{d\varphi}{d\varrho} \right\}^2} = \sqrt{1 + \left\{ \varrho \cdot \frac{1}{\sqrt{1 - \varrho^2}} \right\}^2} = \frac{1}{\sqrt{1 - \varrho^2}},$$

that

$$\begin{aligned} \int_{\mathcal{K}} \sqrt{y} \, ds &= \int_0^1 \sqrt{\varrho \cdot \sin \varphi(\varrho)} \cdot \frac{1}{\sqrt{1 - \varrho^2}} \, d\varrho = \int_0^1 \sqrt{\varrho^2} \cdot \frac{1}{\sqrt{1 - \varrho^2}} \, d\varrho \\ &= \int_0^1 \frac{\varrho}{\sqrt{1 - \varrho^2}} \, d\varrho = \left[ -\sqrt{1 - \varrho^2} \right]_0^1 = 1. \end{aligned}$$

ALTERNATIVELY,  $\varrho = \sin \varphi$ ,  $\varphi \in \left[0, \frac{\pi}{2}\right]$  and

$$\sqrt{\varrho^2 + \left( \frac{d\varrho}{d\varphi} \right)^2} = \sqrt{\sin^2 \varphi + \cos^2 \varphi} = 1,$$

thus

$$\int_{\mathcal{K}} \sqrt{y} \, ds = \int_0^{\frac{\pi}{2}} \sqrt{\sin^2 \varphi} \cdot 1 \, d\varphi = \int_0^{\frac{\pi}{2}} \sin \varphi \, d\varphi = 1.$$

4) It follows from

$$\frac{d\varrho}{d\varphi} = -2a \sin \varphi \cdot \cos \varphi,$$




Figure 23: The curve  $\mathcal{K}$  of **Example 2.1.5**. (In fact, a segment of the line  $x = a$ .)

that

$$\begin{aligned}\varrho^2 + \left(\frac{d\varrho}{d\varphi}\right)^2 &= a^2 \cos^4 \varphi + 4a^2 \sin^2 \varphi \cdot \cos^2 \varphi = a^2 \cos^2 \varphi \{\cos^2 \varphi + 4 \sin^2 \varphi\} \\ &= a^2 \cos^2 \varphi \{4 - 3 \cos^2 \varphi\}.\end{aligned}$$

Now,  $\cos \varphi \geq 0$  for  $\varphi \in \left[0, \frac{\pi}{2}\right]$ , so

$$\sqrt{\varrho^2 + \left(\frac{d\varrho}{d\varphi}\right)^2} = a \cos \varphi \sqrt{4 - 3 \cos^2 \varphi},$$

and the line integral becomes

$$\begin{aligned}\int_{\mathcal{K}} \frac{y}{\sqrt{4a - 3\varrho}} ds &= \int_{\mathcal{K}} \frac{\varrho \sin \varphi}{\sqrt{4a - 3\varrho}} ds = \int_0^{\frac{\pi}{2}} \frac{a \cos^2 \varphi \cdot \sin \varphi}{\sqrt{4a - 3a \cos^2 \varphi}} \cdot a \cos \varphi \sqrt{4 - 3 \cos^2 \varphi} d\varphi \\ &= a\sqrt{a} \int_0^{\frac{\pi}{2}} \cos^3 \varphi \cdot \sin \varphi d\varphi = a\sqrt{a} \left[ -\frac{\cos^4 \varphi}{4} \right]_0^{\frac{\pi}{2}} = \frac{a\sqrt{a}}{4}.\end{aligned}$$

5) If  $\varrho = \frac{a}{\cos \varphi}$ , then

$$\frac{d\varrho}{d\varphi} = \frac{a \sin \varphi}{\cos^2 \varphi},$$

hence



$$\varrho^2 + \left(\frac{d\varrho}{d\varphi}\right)^2 = a^2 \left\{ \frac{1}{\cos^2 \varphi} + \frac{\sin^2 \varphi}{\cos^4 \varphi} \right\} = \frac{a^2}{\cos^4 \varphi},$$

where

$$\sqrt{\varrho^2 + \left(\frac{d\varrho}{d\varphi}\right)^2} = \frac{a}{\cos^2 \varphi}.$$

The line integral is obtained by insertion,

$$\int_{\mathcal{K}} \frac{1}{x^2 + y^2} ds = \int_{\mathcal{K}} \frac{1}{\varrho^2} ds = \int_0^{\pi} \frac{\cos^2 \varphi}{a^2} \cdot \frac{a}{\cos^2 \varphi} d\varphi = \frac{\pi}{4a}.$$

Figure 24: The curve  $\mathcal{K}$  of **Example 2.1.6**.Figure 25: The curve  $\mathcal{K}$  of **Example 2.1.7**.

6) If  $\varphi = \varrho - \ln \varrho$ , then

$$\frac{d\varphi}{d\varrho} = 1 - \frac{1}{\varrho} = \frac{\varrho - 1}{\varrho},$$

hence

$$\sqrt{1 + \left(\varrho \frac{d\varphi}{d\varrho}\right)^2} = \sqrt{1 + (\varrho - 1)^2}.$$

Finally, we get the line integral by insertion,

$$\begin{aligned} \int_{\mathcal{K}} \left( \sqrt{x^2 + y^2} - 1 \right) ds &= \int_{\mathcal{K}} (\varrho - 1) ds = \int_1^2 \sqrt{1 + (\varrho - 1)^2} \cdot (\varrho - 1) d\varrho \\ &= \frac{1}{3} \left[ \{1 + (\varrho - 1)^2\}^{\frac{3}{2}} \right]_1^2 = \frac{1}{3} (2\sqrt{2} - 1). \end{aligned}$$

7) If  $\varphi = \frac{1}{\varrho}$ , then  $\frac{d\varphi}{d\varrho} = -\frac{1}{\varrho^2}$ , så

$$\sqrt{1 + \left(\varrho \frac{d\varphi}{d\varrho}\right)^2} = \sqrt{1 + \frac{1}{\varrho^2}} = \frac{\sqrt{1 + \varrho^2}}{\varrho}.$$

We get the line integral by insertion

$$\int_{\mathcal{K}} (x^2 + y^2) ds = \int_1^2 \varrho^2 \cdot \frac{\sqrt{1 + \varrho^2}}{\varrho} d\varrho = \frac{1}{3} \left[ \{1 + \varrho^2\}^{\frac{3}{2}} \right]_1^2 = \frac{1}{3} \{5\sqrt{5} - 2\sqrt{2}\}.$$

Please click the advert



**Nido**

**Luxurious accommodation**

**Central zone 1 & 2 locations**

**Meet hundreds of international students**

**BOOK NOW and get a £100 voucher from voucherexpress**

**Nido Student Living - London**

**Visit [www.NidoStudentLiving.com/Bookboon](http://www.NidoStudentLiving.com/Bookboon) for more info.**

**+44 (0)20 3102 1060**

Download free ebooks at [bookboon.com](http://bookboon.com)

### 3 Arc lengths and parametric descriptions by the arc length

**Example 3.1** Compute in each of the following cases the arc length of the plane curve  $\mathcal{K}$  given by an equation of the form  $y = Y(x)$ ,  $x \in I$ .

1) The arc length  $\int_{\mathcal{K}} ds$  of the curve

$$y = Y(x) = \frac{x^4 + 48}{24x}, \quad x \in [2, 4].$$

2) The arc length  $\int_{\mathcal{K}} ds$  of the curve

$$y = Y(x) = a \cosh \frac{x}{a}, \quad x \in [-a, a].$$

[Cf. **Example 3.4.1** and **Example 4.1.8**.]

3) The arc length  $\int_{\mathcal{K}} ds$  of the curve

$$y = Y(x) = \ln \frac{e^x - 1}{e^x + 1}, \quad x \in [2, 4].$$

4) The arc length  $\int_{\mathcal{K}} ds$  of the curve

$$y = Y(x) = x^{\frac{3}{2}}, \quad x \in [0, 1].$$

5) The arc length  $\int_{\mathcal{K}} ds$  of the curve

$$y = Y(x) = x^{\frac{2}{3}}, \quad x \in [0, 1].$$

**A** Arc lengths of plane curves.

**D** Sketch the plane curve. Calculate the weight function  $\sqrt{1 + Y'(x)^2}$  and reduce the line integral of integrand 1 to an ordinary integral.




Figure 26: The curve  $\mathcal{K}$  of **Example 3.1.1**.



I 1) It follows from

$$Y(x) = \frac{x^3}{24} + \frac{2}{x}, \quad \text{thus} \quad Y'(x) = \frac{x^2}{8} - \frac{2}{x^2} = \frac{x^4 - 16}{8x^2},$$

that

$$\sqrt{1 + Y'(x)^2} = \frac{1}{8x^2} \sqrt{64x^4 + (x^4 - 16)^2} = \frac{1}{8x^2} (x^4 + 16) = \frac{x^2}{8} + \frac{2}{x^2}.$$

We get the arc length by insertion,

$$\begin{aligned} \int_{\mathcal{K}} ds &= \int_2^4 \sqrt{1 + Y'(x)^2} dx = \int_2^4 \left\{ \frac{x^2}{8} + \frac{2}{x^2} \right\} dx = \left[ \frac{x^3}{24} - \frac{2}{x} \right]_2^4 \\ &= \frac{64 - 8}{24} - \frac{2}{4} + \frac{2}{2} = \frac{56}{24} - \frac{1}{2} + 1 = \frac{7}{3} + \frac{1}{2} = \frac{17}{6}. \end{aligned}$$




Figure 27: The curve  $\mathcal{K}$  of **Example 3.1.2** for  $a = 1$ .

2) From  $Y'(x) = \sinh \frac{x}{a}$  follows that

$$\sqrt{1 + Y'(x)^2} = \sqrt{1 + \sinh^2 \left( \frac{x}{a} \right)} = \cosh \frac{x}{a}.$$

The arc length is


$$\int_{\mathcal{K}} ds = \int_{-a}^a \cosh \frac{x}{a} dx = a \left[ \sinh \frac{x}{a} \right]_{-a}^a = 2a \sinh 1 = \frac{a}{e} (e^2 - 1).$$

3) It follows from

$$Y'(x) = \frac{e^x}{e^x - 1} - \frac{e^x}{e^x + 1} = \frac{2e^x}{e^{2x} - 1},$$

that

$$\sqrt{1 + Y'(x)^2} = \frac{1}{e^{2x} - 1} \sqrt{(e^{2x} - 1)^2 + 4e^{2x}} = \frac{e^{2x} + 1}{e^{2x} - 1} = \frac{\cosh x}{\sinh x},$$

Figure 28: The curve  $\mathcal{K}$  of **Example 3.1.3**.

so the arc length becomes


$$\begin{aligned} \int_{\mathcal{K}} ds &= \int_2^4 \frac{\cosh x}{\sinh x} dx = [\ln \sinh x]_2^4 = \ln \left( \frac{\sinh 4}{\sinh 2} \right) = \ln \left( \frac{2 \sinh 2 \cosh 2}{\sinh 2} \right) \\ &= \ln(2 \cosh 2) = \ln(e^2 + e^{-2}) = \ln(e^4 + 1) - 2. \end{aligned}$$

4) Here,  $Y'(x) = \frac{3}{2} \sqrt{x}$ , so

$$\sqrt{1 + Y'(x)^2} = \sqrt{1 + \frac{9}{4}x}.$$

The arc length is

$$\int_{\mathcal{K}} ds = \int_0^1 \sqrt{1 + \frac{9}{4}x} dx = \frac{4}{9} \cdot \frac{2}{3} \left[ \left(1 + \frac{9}{4}x\right)^{\frac{3}{2}} \right]_0^1 = \frac{8}{27} \left\{ \left(\frac{13}{4}\right)^{\frac{3}{2}} - 1 \right\} = \frac{1}{27} \{13\sqrt{13} - 8\}.$$

Figure 29: The curve  $\mathcal{K}$  of **Example 3.1.5**.

- 5) Since the arc length of  $y = x^{\frac{2}{3}}$ ,  $x \in [0, 1]$ , is equal to the arc length of  $x = y^{\frac{3}{2}}$ , it follows from **Example 3.1.4** that

$$\int_{\mathcal{K}} ds = \frac{1}{27} \{13\sqrt{13} - 9\}.$$

ALTERNATIVELY,  $Y'(x) = \frac{2}{3}x^{-\frac{1}{3}}$ , thus

$$\begin{aligned} \int_{\mathcal{K}} ds &= \int_0^1 \sqrt{1 + \frac{4}{9}x^{-\frac{2}{3}}} dx = \int_0^1 x^{-\frac{1}{3}} \sqrt{x^{\frac{2}{3}} + \frac{4}{9}} dx = \frac{3}{2} \int_0^1 \sqrt{t + \frac{4}{9}} dt \\ &= \left[ \left( t + \frac{4}{9} \right)^{\frac{3}{2}} \right]_0^1 = \left( \frac{13}{9} \right)^{\frac{3}{2}} - \left( \frac{4}{9} \right)^{\frac{3}{2}} = \frac{13\sqrt{13}}{27} - \frac{8}{27} = \frac{1}{27} \{13\sqrt{13} - 8\}. \end{aligned}$$

Please click the advert

I joined MITAS because  
I wanted **real responsibility**

The Graduate Programme  
for Engineers and Geoscientists  
**Maersk.com/Mitas**



**Month 16**


I was a construction  
supervisor in  
the North Sea  
advising and  
helping foremen  
solve problems

Real work  
International opportunities  
Three work placements



 **MAERSK**

Download free ebooks at [bookboon.com](http://bookboon.com)

Figure 30: The curve  $\mathcal{K}$  of **Example 3.2.1**.

**Example 3.2** Compute in each of the following cases the arc length of the given plane curve  $\mathcal{K}$  by an equation in polar coordinates.

1) The arc length  $\int_{\mathcal{K}} ds$  of the curve given by

$$\varrho = a \cos^4 \frac{\varphi}{4}, \quad \varphi \in [0, 4\pi].$$

2) The arc length  $\int_{\mathcal{K}} ds$  of the curve given by

$$\varrho = a(1 + \cos \varphi), \quad \varphi \in [0, 2\pi].$$

3) The arc length  $\int_{\mathcal{K}} ds$  of the curve given by

$$\varphi = \ln \varrho, \quad \varrho \in [1, e].$$

4) The arc length  $\int_{\mathcal{K}} ds$  of the curve given by

$$\varrho = a \sin^3 \frac{\varphi}{3}, \quad \varphi \in [0, 3\pi].$$

**A** Arc lengths in polar coordinates.



**D** First calculate the weight function  $\sqrt{\varrho^2 + \left(\frac{d\varrho}{d\varphi}\right)^2}$  (possibly  $\sqrt{1 + \left(\varrho \frac{d\varphi}{d\varrho}\right)^2}$ ), and then the line integral.

1) Since

$$\frac{d\varrho}{d\varphi} = -a \cdot \cos^3 \frac{\varphi}{4} \cdot \sin \frac{\varphi}{4},$$

the weight is given by

$$\varrho^2 + \left(\frac{d\varrho}{d\varphi}\right)^2 = a^2 \cos^8 \frac{\varphi}{4} + a^2 \cos^6 \frac{\varphi}{4} \cdot \sin^2 \frac{\varphi}{4} = a^2 \cos^6 \frac{\varphi}{4},$$

Figure 31: The curve  $\mathcal{K}$  of **Example 3.2.2**.Figure 32: The curve  $\mathcal{K}$  of **Example 3.2.3**. (Part of the curve of **Example 2.1.2**).

hence


$$\begin{aligned} \int_{\mathcal{K}} ds &= \int_0^{4\pi} \sqrt{\varrho^2 + \left(\frac{d\varrho}{d\varphi}\right)^2} d\varphi = \int_0^{4\pi} a \left| \cos^3 \frac{\varphi}{4} \right| d\varphi = 4a \int_0^{\pi} |\cos^3 t| dt \\ &= 8a \int_0^{\frac{\pi}{2}} \cos^3 t dt = 8a \int_0^{\frac{\pi}{2}} (1 - \sin^2 t) \cos t dt = 8a \left[ \sin t - \frac{1}{3} \sin^3 t \right]_0^{\frac{\pi}{2}} = \frac{16a}{3}. \end{aligned}$$

2) In this case,

$$\sqrt{\varrho^2 + \left(\frac{d\varrho}{d\varphi}\right)^2} = a \sqrt{(1 + \cos \varphi)^2 + \sin^2 \varphi} = a \sqrt{2(1 + \cos \varphi)} = a \sqrt{4 \cos^2 \frac{\varphi}{2}} = 2a \left| \cos \frac{\varphi}{2} \right|,$$

so

$$\int_{\mathcal{K}} ds = \int_0^{2\pi} 2a \left| \cos \frac{\varphi}{2} \right| d\varphi = 4a \int_0^{\pi} |\cos t| dt = 8a \int_0^{\frac{\pi}{2}} \cos t dt = 8a.$$

Figure 33: The curve  $\mathcal{K}$  of **Example 3.2.4**.

3) From

$$\sqrt{1 + \left\{ \varrho \frac{d\varphi}{d\varrho} \right\}^2} = \sqrt{1 + \left\{ \varrho \cdot \frac{1}{\varrho} \right\}^2} = \sqrt{2},$$

follows that

$$\int_{\mathcal{K}} ds = \int_1^e \sqrt{2} d\varrho = \sqrt{2}(e-1).$$

ALTERNATIVELY,  $\varrho = e^\varphi$ ,  $\varphi \in [0, 1]$ , so (cf. **Example 2.1.1**)

$$\sqrt{\varrho^2 + \left( \frac{d\varrho}{d\varphi} \right)^2} = \sqrt{2} e^\varphi,$$

hence


$$\int_{\mathcal{K}} ds = \int_0^1 \sqrt{2} e^\varphi d\varphi = \sqrt{2}(e-1).$$

4) Here  $\frac{d\varrho}{d\varphi} = a \cdot \sin^2 \frac{\varphi}{3} \cdot \cos \frac{\varphi}{3}$ , so

$$\sqrt{\varrho^2 + \left( \frac{d\varrho}{d\varphi} \right)^2} = a \sqrt{\sin^6 \frac{\varphi}{3} + \sin^4 \frac{\varphi}{3} \cdot \cos^2 \frac{\varphi}{3}} = a \cdot \sin^2 \frac{\varphi}{3},$$

thus

$$\int_{\mathcal{K}} ds = \int_0^{3\pi} a \sin^2 \frac{\varphi}{3} d\varphi = 3a \int_0^\pi \sin^2 t dt = \frac{3a}{2} \int_0^\pi (1 - \cos 2t) dt = \frac{3a\pi}{2}.$$

Figure 34: The space curve of **Example 3.3.1**.

**Example 3.3** Below are given some space curves by their parametric descriptions  $\mathbf{x} = \mathbf{r}(t)$ ,  $t \in I$ . Express for each of the curves their parametric description with respect to arc length from the point of the parametric value  $t_0$ .

1) The curve  $\mathbf{r}(t) = (\cos t, \sin t, \ln \cos t)$ , from  $t_0 = 0$  in the interval  $I = \left[0, \frac{\pi}{2}\right]$ .

2) The curve  $\mathbf{r}(t) = \frac{1}{\sqrt{3}}(e^t \cos t, e^t \sin t, e^t)$  from  $t_0 = 0$  in the interval  $I = \mathbb{R}$ .

[Cf. **Example 1.1.7**.]

3) The curve  $\mathbf{r}(t) = (\ln \cos t, \ln \sin t, \sqrt{2}t)$  from  $t_0 = \frac{\pi}{4}$  in the interval  $I = \left]0, \frac{\pi}{2}\right[$ .

4) The curve  $\mathbf{r}(t) = (7t + \cos t, 7t - \cos t, \sqrt{2} \sin t)$  from  $t_0 = \frac{\pi}{2}$  in the interval  $I = \mathbb{R}$ .

5) The curve  $\mathbf{r}(t) = (\cos(2t), \sin(2t), 2 \cosh t)$  from  $t_0 = 0$  in the interval  $I = \mathbb{R}$ .

6) The curve  $\mathbf{r}(t) = (\cos t, \sin t, \ln \cos t)$  from  $t_0 = 0$  in the interval  $I = \left]-\frac{\pi}{2}, \frac{\pi}{2}\right[$ .

[Cf. **Example 1.1.5**.]

**A** Parametric description by the arc length.

**D** Find  $s'(t) = \|\mathbf{r}'(t)\|$  and then  $s = s(t)$  and  $t = \tau(s)$ , where we integrate from  $t_0$ . Finally, insert in  $\mathbf{x} = \mathbf{r}(t) = \mathbf{r}(\tau(s))$ .

**I** 1) From

$$\mathbf{r}'(t) = \left(-\sin t, \cos t, -\frac{\sin t}{\cos t}\right), \quad t \in \left[0, \frac{\pi}{2}\right],$$

follows that

$$s'(t) = \|\mathbf{r}'(t)\| = \sqrt{\sin^2 t + \cos^2 t + \frac{\sin^2 t}{\cos^2 t}} = \frac{1}{\cos t},$$

hence

$$\begin{aligned} s(t) &= \int_0^t \frac{1}{\cos u} du = \int_0^t \frac{\cos u}{1 - \sin^2 u} du = \int_0^t \frac{1}{2} \left( \frac{1}{1 + \sin u} + \frac{1}{1 - \sin u} \right) \cos u du \\ &= \frac{1}{2} \left[ \ln \left( \frac{1 + \sin u}{1 - \sin u} \right) \right]_0^t = \frac{1}{2} \ln \left( \frac{1 + \sin t}{1 - \sin t} \right). \end{aligned}$$

Please click the advert



**Brain power**

By 2020, wind could provide one-tenth of our planet's electricity needs. Already today, SKF's innovative know-how is crucial to running a large proportion of the world's wind turbines.

Up to 25 % of the generating costs relate to maintenance. These can be reduced dramatically thanks to our systems for on-line condition monitoring and automatic lubrication. We help make it more economical to create cleaner, cheaper energy out of thin air.


By sharing our experience, expertise, and creativity, industries can boost performance beyond expectations. Therefore we need the best employees who can meet this challenge!

**The Power of Knowledge Engineering**

Plug into The Power of Knowledge Engineering.  
Visit us at [www.skf.com/knowledge](http://www.skf.com/knowledge)

**SKF**



Figure 35: The space curve of **Example 3.3.2**.

Then

$$\frac{1 + \sin t}{1 - \sin t} = e^{2s}, \quad \text{dvs.} \quad \sin t = \frac{e^{2s} - 1}{e^{2s} + 1} = \tanh s, \quad s \geq 0.$$

Notice that it follows from  $t \in \left[0, \frac{\pi}{2}\right[$  that

$$\cos t = \frac{2e^s}{e^{2s} + 1} = \frac{1}{\cosh s}.$$

Thus

$$t = \operatorname{Arcsin}\left(\frac{e^{2s} - 1}{e^{2s} + 1}\right) = \operatorname{Arcsin}(\tanh s), \quad s \geq 0,$$

and the parametric description by the arc length is

$$\begin{aligned} \mathbf{r}(s) &= (\cos t, \sin t, \ln \cos t) = \left( \frac{2e^s}{e^{2s} + 1}, \frac{e^{2s} - 1}{e^{2s} + 1}, \ln \left( \frac{2e^s}{e^{2s} + 1} \right) \right) \\ &= \left( \frac{1}{\cosh s}, \tanh s, -\ln \cosh s \right), \quad s \geq 0. \end{aligned}$$

2) Here


$$\mathbf{r}'(t) = \frac{1}{\sqrt{3}} e^t (\cos t - \sin t, \cos t + \sin t, 1),$$

so

$$s'(t) = \|\mathbf{r}'(t)\| = \frac{1}{\sqrt{3}} e^t \sqrt{(\cos t - \sin t)^2 + (\cos t + \sin t)^2 + 1} = e^t,$$

hence

$$s(t) = \int_0^t e^u du = e^t - 1, \quad \text{and} \quad t = \ln(s + 1), \quad s > -1.$$

Figure 36: The space curve of **Example 3.3.3**.

Finally, we get the parametric description by the arc length,

$$\begin{aligned}\mathbf{r}(s) &= \frac{1}{\sqrt{3}} ((s+1) \cos(\ln(s+1)), (s+1) \sin(\ln(s+1)), s+1) \\ &= \frac{s+1}{\sqrt{3}} (\cos(\ln(s+1)), \sin(\ln(s+1)), 1), \quad s > -1.\end{aligned}$$

3) From

$$\mathbf{r}'(t) = \left( -\frac{\sin t}{\cos t}, \frac{\cos t}{\sin t}, \sqrt{2} \right), \quad t \in \left] 0, \frac{\pi}{2} \right[ ,$$

follows that

$$s'(t) = \|\mathbf{r}'(t)\| = \sqrt{\frac{\sin^2 t}{\cos^2 t} + 2 + \frac{\cos^2 t}{\sin^2 t}} = \frac{\sin t}{\cos t} + \frac{\cos t}{\sin t} = \frac{1}{\cos t \sin t},$$

as  $t \in \left] 0, \frac{\pi}{2} \right[$ . Then

$$s(t) = \int_{\frac{\pi}{4}}^t \frac{1}{\cos u \sin u} du = \int_{\frac{\pi}{4}}^t \left( \frac{\sin u}{\cos u} + \frac{\cos u}{\sin u} \right) du = \ln \frac{\sin t}{\cos t} = \ln \tan t,$$

and thus  $s \in \mathbb{R}$  and  $\tan t = e^s$ , and



$$\cos t = \frac{+1}{\sqrt{1 + \tan^2 t}} = \frac{1}{\sqrt{1 + e^{2s}}}, \quad \text{and} \quad \sin t = \frac{e^s}{\sqrt{1 + e^{2s}}}.$$

The parametric description by the arc length is

$$\mathbf{r}(s) = \left( -\frac{1}{2} \ln(1 + e^{2s}), s - \frac{1}{2} \ln(1 + e^{2s}), \sqrt{2} \operatorname{Arctan}(e^s) \right), \quad s \in \mathbb{R}.$$

4) Here,

$$\mathbf{r}'(t) = (7 - \sin t, 7 + \sin t, \sqrt{2} \cos t),$$

Figure 37: The space curve of **Example 3.3.4**.Figure 38: The space curve of **Example 3.3.5**.

so

$$\begin{aligned} s'(t) &= \|\mathbf{r}'(t)\| = \sqrt{(7 - \sin t)^2 + (7 + \sin t)^2 + 2 \cos^2 t} \\ &= \sqrt{2 \cdot 49 + 2 \sin^2 t + 2 \cos^2 t} = \sqrt{98 + 2} = 10, \end{aligned}$$

and thus

$$s(t) = \int_{\frac{\pi}{2}}^t 10 \, du = 10 \left( t - \frac{\pi}{2} \right), \quad \text{so} \quad t = \frac{s}{10} + \frac{\pi}{2}, \quad s \in \mathbb{R},$$

and the parametric description with the arc length as parameter from the point  $\left( \frac{7\pi}{2}, \frac{7\pi}{2}, \sqrt{2} \right)$  is


$$\mathbf{r}(s) = \left( \frac{7s + 35\pi}{10} - \sin\left(\frac{s}{10}\right), \frac{7s + 35\pi}{10} + \sin\left(\frac{s}{10}\right), \sqrt{2} \cos\left(\frac{s}{10}\right) \right),$$

for  $s \in \mathbb{R}$ .

5) It follows from

$$\mathbf{r}'(t) = (-2 \sin 2t, 2 \cos 2t, 2 \sinh t),$$

Please click the advert



## Are you considering a European business degree?

**LEARN BUSINESS** at university level. We mix cases with cutting edge research working individually or in teams and everyone speaks English. Bring back valuable knowledge and experience to boost your career.

**MEET** a culture of new foods, music and traditions and a new way of studying business in a safe, clean environment – in the middle of Copenhagen, Denmark.


**ENGAGE** in extra-curricular activities such as case competitions, sports, etc. – make new friends among CBS' 18,000 students from more than 80 countries.

**Copenhagen Business School**  
HANDELSHØJSKOLEN

EFMD **EQUIS** AACSB ACCREDITED BY ASSOCIATION OF MBAs CEMS EQUIS

See what we look like and how we work on [cbs.dk](http://cbs.dk)

Download free ebooks at [bookboon.com](http://bookboon.com)

Figure 39: The space curve of **Example 3.3.6**; cf. **Example 3.3.1**.

that

$$s'(t) = \|\mathbf{r}'(t)\| = 2\sqrt{\sin^2(2t) + \cos^2(2t) + \sinh^2 t} = 2 \cosh t,$$

hence

$$s(t) \int_0^t 2 \cosh u \, du = 2 \sinh t,$$

so  $s \in \mathbb{R}$  and

$$t = \operatorname{Arsinh} \left( \frac{s}{2} \right) = \ln \left( \frac{1}{2} \left( s + \sqrt{s^2 + 4} \right) \right), \quad s \in \mathbb{R}.$$

The parametric description with the arc length as parameter is

$$\begin{aligned} \mathbf{r}(s) &= \left( \cos \left( 2 \operatorname{Arsinh} \left( \frac{s}{2} \right) \right), \sin \left( 2 \operatorname{Arsinh} \left( \frac{s}{2} \right) \right), 2\sqrt{1 + \frac{s^2}{4}} \right) \\ &= \left( \cos \left( 2 \operatorname{Arsinh} \left( \frac{s}{2} \right) \right), \sin \left( 2 \operatorname{Arsinh} \left( \frac{s}{2} \right) \right), \sqrt{4 + s^2} \right), \end{aligned}$$

for  $s \in \mathbb{R}$ .

- 6) This is an extension of the curve of **Example 3.3.1**, with the same parametric description evaluated from the same point  $t_0 = 0$ . We can therefore reuse this example, since the only change is that  $s \in \mathbb{R}$ ,

$$\mathbf{r}(s) = \left( \frac{1}{\cosh s}, \tanh s, -\ln \cosh s \right), \quad \text{for } s \in \mathbb{R}.$$




Figure 40: The chain curve for  $a = 1$ , cf. **Example 3.4.1**.

**Example 3.4** Find for every one of the given plane curves below an equation of the form

(1)  $\psi = \Psi(s)$ ,

where the signed arc length  $s$  is computed from a fixed point  $P_0$  on the curve, while  $\psi$  is the angle between the oriented tangents at  $P_0$  and at the point  $P$  on the curve given by  $s$ .

- 1) The chain curve given by  $y = a \cosh \frac{x}{a}$ , from  $P_0$  given by  $x = 0$ .

[Cf. **Example 1.2.2**.]

- 2) The asteroid given by

$$\mathbf{r}(t) = a(-\cos^3 t, \sin^3 t), \quad t \in \left[0, \frac{\pi}{2}\right],$$

from  $P_0$  given by  $t = 0$ .

- 3) The winding of a circle given by

$$\mathbf{r}(t) = a(\cos t + t \sin t, \sin t - t \cos t), \quad t \in \mathbb{R}_+,$$

from  $P_0$  given by  $t = 0$ .

- 4) The cycloid given by


$$\mathbf{r}(t) = a(t - \sin t, 1 - \cos t), \quad t \in [0, 2\pi],$$

from  $P_0$  given by  $t = \pi$ .

It can be proved that (1) determines the curve uniquely with exception of its placement in the plane. Therefore, (1) is also called the natural equation of the curve.

**A** Natural equation.

**D** Find the arc length  $s$ , and then  $\psi$  by a geometrical analysis.

Figure 41: The astroid of **Example 3.4.2**.

- 1) The point  $P_0$  has the coordinates  $(0, a)$ . A parametric description of the chain curve is e.g.

$$\mathbf{r}(t) = \left( t, a \cosh \frac{t}{a} \right),$$

hence

$$\mathbf{r}'(t) = \left( 1, \sinh \frac{t}{a} \right), \quad \text{where } \mathbf{r}'(0) = (1, 0),$$

and thus  $\psi = \text{Arctan} \left( \sinh \frac{t}{a} \right)$ .

From

$$s'(t) = \|\mathbf{r}'(t)\| = \sqrt{1 + \sinh^2 \frac{t}{a}} = \cosh \frac{t}{a},$$

follows that

$$s(t) = \int_0^t \cosh \frac{u}{a} dy = a \left[ \sinh \frac{u}{a} \right]_0^t = a \sinh \left( \frac{t}{a} \right).$$

The natural equation is

$$\psi = \Psi(s) = \text{Arctan} \left( \frac{s}{a} \right).$$


- 2) The point  $P_0$  has the coordinates  $(-a, 0)$ , and

$$\mathbf{r}'(t) = a (3 \cos^2 t \cdot \sin t, 3 \sin^2 t \cdot \cos t) = 3a \cos t \cdot \sin t (\cos t, \sin t).$$

For  $t \rightarrow 0+$  we get  $\mathbf{r}'(0) = \mathbf{0}$ , and by considering a figure we may conclude that we have a horizontal half tangent. Then it follows that  $\psi = t$ .

It follows from

$$s'(t) = \|\mathbf{r}'(t)\| = 3a \cos t \cdot \sin t,$$

Figure 42: The winding of the circle of **Example 3.4.3**.

that

$$s(t) = \int_0^t 3a \cos u \cdot \sin u \, du = \frac{3a}{2} \int_0^t \sin 2u \, du = \frac{3a}{4} \{1 - \cos 2t\},$$

hence

$$\cos 2t = 1 - \frac{4s}{3a},$$

and

$$\psi = \Psi(s) = t = \frac{1}{2} \operatorname{Arccos} \left( 1 - \frac{4s}{3a} \right).$$

- 3) The point  $P_0$  has the coordinates  $(a, 0)$ , and

$$\mathbf{r}'(t) = a(-\sin t + \sin t + t \cos t, \cos t - \cos t + t \sin t) = at(\cos t, \sin t).$$

It follows that  $\psi = t$ .

As  $t > 0$  we have

$$s'(t) = \|\mathbf{r}'(t)\| = at,$$

thus

$$s(t) = a \int_0^t u \, du = \frac{a}{2} t^2, \quad \text{so } t = \sqrt{\frac{2s}{a}},$$


and hence

$$\psi = \Psi(s) = \sqrt{\frac{2s}{a}}.$$

- 4) The point  $P_0$  is described by  $\mathbf{r}(\pi) = a(\pi, 0)$ . The curve has a vertical half tangent at  $P_0$ . From

$$\mathbf{r}'(t) = a(1 - \cos t, \sin t) = a \left( 2 \sin^2 \frac{t}{2}, 2 \sin \frac{t}{2} \cos \frac{t}{2} \right) = 2a \sin \frac{t}{2} \left( \sin \frac{t}{2}, \cos \frac{t}{2} \right),$$



Figure 43: The cycloid of **Example 3.4.4**.

follows that

$$s'(t) = \|\mathbf{r}'(t)\| = 2a \sin \frac{t}{2},$$

Please click the advert



## The financial industry needs a strong software platform

That's why we need you

SimCorp is a leading provider of software solutions for the financial industry. We work together to reach a common goal: to help our clients succeed by providing a strong, scalable IT platform that enables growth, while mitigating risk and reducing cost. At SimCorp, we value commitment and enable you to make the most of your ambitions and potential.

Are you among the best qualified in finance, economics, IT or mathematics?

Find your next challenge at  
[www.simcorp.com/careers](http://www.simcorp.com/careers)



[www.simcorp.com](http://www.simcorp.com)

MITIGATE RISK | REDUCE COST | ENABLE GROWTH

so

$$s(t) = \int_{\pi}^t 2a \sin \frac{u}{2} du = 4a \left[ -\cos \frac{u}{2} \right]_{\pi}^t = 4a \left\{ -\cos \frac{t}{2} \right\},$$

and hence

$$\cos \frac{t}{2} = -\frac{s}{4a}, \quad \text{dvs.} \quad t = 2 \operatorname{Arccos} \left( -\frac{s}{4a} \right).$$

Since  $\psi$  must have the form  $at + b$ , it is easy to derive that

$$\psi = \pi - t = \pi - 2 \operatorname{Arccos} \left( -\frac{s}{4a} \right).$$

**Example 3.5** A plane curve  $\mathcal{K}$  is given by the parametric description

$$\mathbf{r}(t) = \left( a \int_0^t \sin(u^2) du, a \int_0^t \cos(u^2) du \right), \quad t \in \mathbb{R}.$$

The signed arc length from the point  $(0,0)$  is called  $s$ .

1. Find  $s$ , and find the parametric description of the curve given by the arc length.

It is proved in Differential Geometry that any plane curve has a curvature

$$\kappa(t) = \frac{\{\mathbf{e}_z \times \mathbf{r}'(t)\} \cdot \mathbf{r}''(t)}{\|\mathbf{r}'(t)\|^3},$$

where we let the plane of the curve be the  $(X, Y)$ -plane in the space.

2. Prove that  $\kappa$  is proportional to  $s$  for  $\mathcal{K}$ .

The curve under consideration has many names: the clothoid, Euler's spiral, Cornu's spiral.

REMARK. "Clothoid" means in koiné, i.e. Ancient Greek:  $\kappa\lambda\omega\theta\omega = \text{I spin}$ .  $\diamond$

**A** Parametric description with respect to arc length, curvature.

**D** Find  $ds$  and then compute.

**I** 1) As  $s'(t) = \|\mathbf{r}'(t)\|$  and  $\mathbf{r}'(t) = a(\sin(t^2), \cos(t^2))$ , fås


$$s'(t) = a\sqrt{\sin^2(t^2) + \cos^2(t^2)} = a,$$

we get

$$s(t) = at \quad \text{and} \quad t(s) = \frac{1}{a}s.$$

The parametric description with the arc length is

$$\mathbf{x} = \mathbf{r}(t) = a \left( \int_0^t \sin(u^2) du, \int_0^t \cos(u^2) du \right) = a \left( \int_0^{\frac{s}{a}} \sin(u^2) du, \int_0^{\frac{s}{a}} \cos(u^2) du \right).$$

Figure 44: The clothoid for  $a = 1$  and  $s \in [-4, 4]$ .

2) From

$$\mathbf{r}'(t) = a (\sin(t^2), \cos(t^2)) \sim a (\sin(t^2), \cos(t^2), 0),$$

and

$$\mathbf{r}''(t) = 2ta (\cos(t^2), -\sin(t^2)) \sim 2ta (\cos(t^2), -\sin(t^2), 0),$$

follows that

$$\begin{aligned} \{\mathbf{e}_z \times \mathbf{r}'(t)\} \cdot \mathbf{r}''(t) &= \begin{vmatrix} 2ta \cos(t^2) & -2ta \sin(t^2) & 0 \\ 0 & 0 & 1 \\ a \sin(t^2) & a \cos(t^2) & 0 \end{vmatrix} = - \begin{vmatrix} 2ta \cos(t^2) & -2ta \sin(t^2) \\ a \sin(t^2) & a \cos(t^2) \end{vmatrix} \\ &= -2ta^2. \end{aligned}$$

As  $\|\mathbf{r}'(t)\| = a$ , we finally get

$$\kappa = \frac{\{\mathbf{e}_z \times \mathbf{r}'(t)\} \cdot \mathbf{r}''(t)}{\|\mathbf{r}'(t)\|^3} = -\frac{2ta^2}{a^3} = -2 \frac{t}{a} = -\frac{2s}{a^2}.$$


**Example 3.6** A space curve  $\mathcal{K}$  is given by the parametric description

$$\mathbf{r}(t) = \left( \frac{1}{2} t^2 - \ln t, 2 \sin t, 2 \cos t \right), \quad t \in [1, 2].$$

Prove that  $\|\mathbf{r}'(t)\| = t + \frac{1}{t}$ , and find the length of  $\mathcal{K}$ .

**A** Arc length.

**D** Compute  $\|\mathbf{r}'(t)\|$  og  $\ell = \int_0^1 \|\mathbf{r}'(t)\| dt$ .

Figure 45: The curve  $\mathcal{K}$ .

**I** It follows from

$$\mathbf{r}'(t) = \left( t - \frac{1}{t}, 2 \cos t, -2 \sin t \right), \quad t \in [1, 2],$$

that

$$\|\mathbf{r}'(t)\|^2 = \left( t - \frac{1}{t} \right)^2 + 4 \cos^2 t + 4 \sin^2 t = t^2 - 2 + \frac{1}{t^2} = \left( t + \frac{1}{t} \right)^2,$$

thus

$$\|\mathbf{r}'(t)\| = \left| t + \frac{1}{t} \right| = t + \frac{1}{t},$$

and accordingly,

$$\ell = \int_1^2 \|\mathbf{r}'(t)\| dt = \int_1^2 \left( t + \frac{1}{t} \right) dt = \left[ \frac{t^2}{2} + \ln t \right]_1^2 = \frac{3}{2} + \ln 2.$$

**Example 3.7** A space curve  $\mathcal{K}$  is given by the parametric description

$$\mathbf{r}(t) = (e^{3t}, e^{-3t}, \sqrt{18}t), \quad t \in [-1, 1].$$

Prove that  $\|\mathbf{r}'(t)\| = 3(e^{3t} + e^{-3t})$ , and find the arc length of  $\mathcal{K}$ .

**A** Arc length.


**D** Find  $\mathbf{r}'(t)$ .

**I** We get by differentiation

$$\mathbf{r}'(t) = (3e^{3t}, -3e^{-3t}, \sqrt{18}) = 3(e^{3t}, -e^{-3t}, \sqrt{2}),$$

thus

$$\|\mathbf{r}'(t)\| = 3\sqrt{(e^{3t})^2 + (-e^{-3t})^2 + 2} = 3\sqrt{(e^{3t} + e^{-3t})^2} = 3(e^{3t} + e^{-3t}),$$

Figure 46: The curve  $\mathcal{K}$ .

and we get the arc length

$$\begin{aligned}\ell(\mathcal{K}) &= \int_{-1}^1 \|\mathbf{r}'(t)\| dt = \int_{-1}^1 3(e^{3t} + e^{-3t}) dt \\ &= 2 \int_0^1 3 \cdot 2 \cosh 3t dt = 4[\sinh 3t]_0^1 = 4 \sinh 3.\end{aligned}$$

Please click the advert

# What do you want to do?

No matter what you want out of your future career, an employer with a broad range of operations in a load of countries will always be the ticket. Working within the Volvo Group means more than 100,000 friends and colleagues in more than 185 countries all over the world. We offer graduates great career opportunities – check out the Career section at our web site [www.volvogroup.com](http://www.volvogroup.com). We look forward to getting to know you!

**VOLVO**  
AB Volvo (publ)  
[www.volvogroup.com](http://www.volvogroup.com)

VOLVO TRUCKS | RENAULT TRUCKS | MACK TRUCKS | VOLVO BUSES | VOLVO CONSTRUCTION EQUIPMENT | VOLVO PENTA | VOLVO AERO | VOLVO IT  
VOLVO FINANCIAL SERVICES | VOLVO 3P | VOLVO POWERTRAIN | VOLVO PARTS | VOLVO TECHNOLOGY | VOLVO LOGISTICS | BUSINESS AREA ASIA

Download free ebooks at [bookboon.com](http://bookboon.com)

**Example 3.8** A space curve  $\mathcal{K}$  is given by the parametric description

$$\mathbf{r}(t) = \left( \frac{1}{3}t^3 - t, \frac{1}{3}t^3 + t, t^2 \right), \quad t \in [-1, 1].$$

Find the arc length of  $\mathcal{K}$ .

**A** Arc length.

**D** Find  $\|\mathbf{r}'(t)\|$ .




Figure 47: The curve  $\mathcal{K}$ .

**I** It follows from

$$\mathbf{r}'(t) = (t^2 - 1, t^2 + 1, 2t),$$

that

$$\|\mathbf{r}'(t)\|^2 = (t^2 - 1)^2 + (t^2 + 1)^2 + 4t^2 = 2t^4 + 2 + 4t^2 = 2(t^2 + 1)^2,$$

hence

$$\ell(\mathcal{K}) = \int_{-1}^1 \|\mathbf{r}'(t)\| dt = 2 \int_0^1 \sqrt{2}(t^2 + 1) dt = 2\sqrt{2} \left( \frac{1}{3} + 1 \right) = \frac{8\sqrt{2}}{3}.$$


**Example 3.9** A space curve  $\mathcal{K}$  is given by the parametric description

$$\mathbf{r}(t) = (6t^2, 4\sqrt{2}t^3, 3t^4), \quad t \in [-1, 1].$$

Explain why the curve is symmetric with respect to the  $(X, Z)$ -plane. Then find the arc length of  $\mathcal{K}$ .

**A** Arc length.

**D** Replace  $t$  by  $-t$ . Then find  $\mathbf{r}'(t)$ .

Figure 48: The curve  $\mathcal{K}$ .

**I** It follows from  $\mathbf{r}(-t) = (6t^2, -4\sqrt{2}t^3, 3t^4)$ , that the curve is symmetric with respect to the  $(X, Z)$ -plane.

From

$$\mathbf{r}'(t) = (12t, 12\sqrt{2}t^2, 12t^3) = 12t(1, \sqrt{2}t, t^2)$$

follows that

$$\|\mathbf{r}'(t)\| = 12|t| \cdot \sqrt{1 + 2t^2 + t^4} = 12|t| \cdot (1 + t^2).$$

Finally, when we exploit the symmetry above and put  $u = t^2$ , we find the arc length

$$\ell(\mathcal{K}) = 2 \int_0^1 \|\mathbf{r}'(t)\| dt = 2 \int_0^1 12t(1 + t^2) dt = 12 \int_0^1 (1 + u) du = 12 \left(1 + \frac{1}{2}\right) = 18.$$

**Example 3.10** A space curve  $\mathcal{K}$  is given by the parametric description

$$\mathbf{r}(t) = (t + \sin t, \sqrt{2} \cos t, t - \sin t), \quad t \in [-1, 1].$$

1) Find a parametric description of the tangent of  $\mathcal{K}$  at the point corresponding to  $t = 0$ .

2) Compute the arc length of  $\mathcal{K}$ .

**A** Space curve.


**D** Follow the standard method.

**I** 1) As  $\mathbf{r}(0) = (0, \sqrt{2}, 0)$ , and

$$\mathbf{r}'(t) = (1 + \cos t, -\sqrt{2} \sin t, 1 - \cos t), \quad \mathbf{r}'(0) = (2, 0, 0),$$

it follows that a parametric description of the tangent of  $\mathcal{K}$  at  $(0, \sqrt{2}, 0)$  is given by

$$\mathbf{x}(u) = (0, \sqrt{2}, 0) + (2u, 0, 0) = (2u, \sqrt{2}, 0), \quad u \in \mathbb{R}.$$

Figure 49: The curve  $\mathcal{K}$  and its tangent at  $(0, \sqrt{2}, 0)$ .

2) The arc length of  $\mathcal{K}$  is

$$\begin{aligned} \int_{-1}^1 \|\mathbf{r}'(t)\| dt &= \int_{-1}^1 \sqrt{(1+\cos t)^2 + 2\sin^2 t + (1-\cos t)^2} dt \\ &= \int_{-1}^1 \sqrt{2 + 2\cos^2 t + 2\sin^2 t} dt = \int_{-1}^1 \sqrt{4} dt = 2 \cdot 2 = 4. \end{aligned}$$

Please click the advert

*RedStarRésumé.*

## Do you want your Dream Job?

More customers get their dream job by using RedStarResume than any other resume service.

RedStarResume can help you with your job application and CV.

**Go to: [Redstarresume.com](http://Redstarresume.com)**  
**Use code "BOOKBOON" and save up to \$15**

(enter the discount code in the "Discount Code Box")





## 4 Tangential line integrals

**Example 4.1** Calculate in each of the following cases the tangential line integral

$$\int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x}$$

of the vector field  $\mathbf{V}$  along the den plane curve  $\mathcal{K}$ . This curve will either be given by a parametric description or by an equation. First sketch the curve.

- 1) The vector field  $\mathbf{V}(x, y) = (x^2 + y^2, x^2 - y^2)$  along the curve  $\mathcal{K}$  given by  $y = 1 - |1 - x|$  for  $x \in [0, 2]$ .
- 2) The vector field  $\mathbf{V}(x, y) = (x^2 - 2xy, y^2 - 2xy)$  along the curve  $\mathcal{K}$  given by  $y = x^2$  for  $x \in [-1, 1]$ .
- 3) The vector field  $\mathbf{V}(x, y) = (2a - y, x)$  along the curve  $\mathcal{K}$  given by  $\mathbf{r}(t) = a(t - \sin t, 1 - \cos t)$  for  $t \in [0, 2\pi]$ .
- 4) The vector field  $\mathbf{V}(x, y) = \left( \frac{x+y}{x^2+y^2}, \frac{y-x}{x^2+y^2} \right)$  along the curve  $\mathcal{K}$  given by  $x^2 + y^2 = a^2$  and run through in the positive orientation of the plane.
- 5) The vector field  $\mathbf{V}(x, y) = (x^2 - y^2, -(x+y))$  along the curve  $\mathcal{K}$  given by  $\mathbf{r}(t) = (a \cos t, b \sin t)$  for  $t \in \left[0, \frac{\pi}{2}\right]$ .
- 6) The vector field  $\mathbf{V}(x, y) = (x^2 - y^2, -(x+y))$  along the curve  $\mathcal{K}$  given by  $\mathbf{r}(t) = (a(1-t), bt)$  for  $t \in [0, 1]$ .
- 7) The vector field  $\mathbf{V}(x, y) = (-y^3, x^3)$  along the curve  $\mathcal{K}$  given by  $\mathbf{r}(t) = (1 + \cos t, \sin t)$  for  $t \in \left[\frac{\pi}{2}, \pi\right]$ .
- 8) The vector field  $\mathbf{V}(x, y) = \left(-y^2, a^2 \sinh \frac{x}{a}\right)$  along the curve  $\mathcal{K}$  given by  $y = a \cosh \frac{x}{a}$  for  $x \in [a, 2a]$ .

**A** Tangential line integrals.

**D** First sketch the curve. Then compute the tangential line integral.




Figure 50: The curve  $\mathcal{K}$  of **Example 4.1.1**.

I 1) Here the parametric description of the curve can also be written

$$y = \begin{cases} x & \text{for } x \in [0, 1], \\ 2 - x & \text{for } x \in [1, 2]. \end{cases}$$


**Try this...**



The sequence 2, 4, 6, 8, 10, 12, 14, 16, ... is the sequence of even whole numbers. The 100th place in this sequence is the number...?

Challenging? Not challenging? Try more >>

[www.alloptions.nl/life](http://www.alloptions.nl/life)



Figure 51: The curve  $\mathcal{K}$  of **Example 4.1.2**.

This gives the following computation of the tangential line integral

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{ (x^2 + y^2) dx + (x^2 - y^2) dy \} \\
 &= \int_0^1 \{ (x^2 + x^2) dx + (x^2 - x^2) dx \} \\
 &\quad + \int_1^2 \{ (x^2 + (2-x)^2) dx + (x^2 - (2-x)^2) (-dx) \} \\
 &= \int_0^1 2x^2 dx + \int_1^2 2(2-x)^2 dx = \frac{2}{3} [x^3]_0^1 + \frac{2}{3} [(x-2)^3]_1^2 \\
 &= \frac{2}{3} + \frac{2}{3} = \frac{4}{3}.
 \end{aligned}$$

2) Here

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{ (x^2 - 2xy) dx + (y^2 - 2xy) dy \} \\
 &= \int_{-1}^1 \{ (x^2 - 2x^3) dx + (x^4 - 2x^3) \cdot 2x dx \} \\
 &= \int_{-1}^1 (x^2 - 2x^3 + 2x^5 - 4x^4) dx \\
 &= \int_{-1}^1 (x^2 - 4x^4) dx + 0 = 2 \left[ \frac{1}{3} x^3 - \frac{4}{5} x^5 \right]_0^1 \\
 &= 2 \left( \frac{1}{3} - \frac{4}{5} \right) = \frac{2}{15} (5 - 12) = -\frac{14}{15}.
 \end{aligned}$$

Figure 52: The curve  $\mathcal{K}$  of **Example 4.1.3** for  $a = 1$ .Figure 53: The curve  $\mathcal{K}$  of **Example 4.1.4** for  $a = 1$ .

3) Similarly we get


$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{(2a - y) dx + x dy\} \\
 &= \int_0^{2\pi} \{(2a - a(1 - \cos t))a(1 - \cos t) + a(t - \sin t)a \sin t\} dt \\
 &= a^2 \int_0^{2\pi} \{(1 + \cos t)(1 - \cos t) + (t - \sin t) \sin t\} dt \\
 &= a^2 \int_0^{2\pi} \{1 - \cos^2 t + t \sin t - \sin^2 t\} dt = a^2 \int_0^{2\pi} t \sin t dt \\
 &= a^2 [-t \cos t + \sin t]_0^{2\pi} = -2\pi a^2.
 \end{aligned}$$

4) We split the curve  $\mathcal{K}$  into two pieces,  $\mathcal{K} = \mathcal{K}_1 + \mathcal{K}_2$ , where  $\mathcal{K}_1$  lies in the upper half plane, and  $\mathcal{K}_2$  lies in the lower half plane, i.e.  $y > 0$  inside  $\mathcal{K}_1$ , and  $y < 0$  inside  $\mathcal{K}_2$ . Then we get the

tangential line integral

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \left( \frac{x+y}{x^2+y^2} dx + \frac{y-x}{x^2+y^2} dy \right) \\
 &= \int_{\mathcal{K}} \frac{1}{x^2+y^2} \frac{1}{2} d(x^2+y^2) + \int_{\mathcal{K}} \frac{1}{x^2+y^2} (y dx - x dy) \\
 &= \int_{\mathcal{K}} \frac{1}{2} d \ln(x^2+y^2) + \int_{\mathcal{K}_1} \frac{1}{1+\left(\frac{x}{y}\right)^2} \left( \frac{1}{y} dx + x \left( -\frac{1}{y^2} \right) dy \right) \\
 &\quad + \int_{\mathcal{K}_2} \frac{1}{1+\left(\frac{x}{y}\right)^2} \left( \frac{1}{y} dx + x \left( -\frac{1}{y^2} \right) dy \right) \\
 &= 0 + \int_{\mathcal{K}_1} \frac{1}{1+\left(\frac{x}{y}\right)^2} d\left(\frac{x}{y}\right) + \int_{\mathcal{K}_2} \frac{1}{1+\left(\frac{x}{y}\right)^2} d\left(\frac{x}{y}\right) \\
 &= \int_{\mathcal{K}_1} d\text{Arctan}\left(\frac{x}{y}\right) + \int_{\mathcal{K}_2} d\text{Arctan}\left(\frac{x}{y}\right) \\
 &= [\text{Arctan } t]_{+\infty}^{-\infty} + [\text{Arctan } t]_{+\infty}^{-\infty} = -\pi - \pi = -2\pi.
 \end{aligned}$$

Please click the advert






Figure 54: The curve  $\mathcal{K}$  of **Example 4.1.5** for  $a = 1$  and  $b = 2$ .

ALTERNATIVELY we get by using the parametric description

$$(x, y) = a(\cos t, \sin t), \quad t \in [0, 2\pi],$$

that

$$\begin{aligned} \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \left( \frac{x+y}{x^2+y^2} dx + \frac{y-x}{x^2+y^2} dy \right) \\ &= \int_0^{2\pi} \frac{a^2}{a^2} \{ (\cos t + \sin t)(-\sin t) + (\sin t - \cos t) \cos t \} dt \\ &= \int_0^{2\pi} \{ -\cos t \cdot \sin t - \sin^2 t + \cos t \cdot \sin t - \cos^2 t \} dt \\ &= - \int_0^{2\pi} dt = -2\pi. \end{aligned}$$

5) Here

$$\begin{aligned} \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{ (x^2 - y^2) dx - (x + y) dy \} \\ &= \int_0^{\frac{\pi}{2}} \{ (a^2 \cos^2 t - b^2 \sin^2 t)(-a \sin t) - (a \cos t + b \sin t)b \cos t \} dt \\ &= \int_0^{\frac{\pi}{2}} \{ -a[(a^2 + b^2) \cos^2 t - b^2] \sin t - ab \cos^2 t - b \sin t \cos t \} dt \\ &= \left[ +a(a^2 + b^2) \frac{1}{3} \cos^3 t - ab^2 \cos t - \frac{ab}{2} \left( t + \frac{1}{2} \sin 2t \right) - \frac{1}{2} b^2 \sin^2 t \right]_0^{\frac{\pi}{2}} \\ &= -\frac{ab}{2} \cdot \frac{\pi}{2} - \frac{b^2}{2} - \frac{a(a^2 + b^2)}{3} + ab^2 = \frac{a}{3}(2b^2 - a^2) - \frac{b}{4}(2b + a\pi). \end{aligned}$$





Figure 55: The curve  $\mathcal{K}$  of **Example 4.1.6** for  $a = 1$  and  $b = 2$ .

6) Here

$$\begin{aligned}
 \int_{\mathcal{V}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{(x^2 - y^2)dx - (x + y)dy\} \\
 &= \int_0^1 \{[a^2(1-t)^2 - b^2t^2](-a) - [a - at + bt] \cdot b\} dt \\
 &= \int_0^1 \{-a^3(t-1)^2 + ab^2t^2 + b(a-b)t - ab\} dt \\
 &= \left[ -\frac{a^3}{3}(t-1)^3 + \frac{ab^2}{3}t^3 + \frac{1}{2}b(a-b)t^2 - abt \right]_0^1 \\
 &= \frac{ab^2}{3} + \frac{1}{2}(a-b)b - ab - \frac{a^3}{3} \\
 &= \frac{a}{3}(b^2 - a^2) - \frac{b}{2}(a+b).
 \end{aligned}$$

REMARK. The vector field  $\mathbf{V}(\mathbf{x})$  is identical to that in **Example 4.1.5** and in **Example 4.1.6**. Furthermore, the curves of these two examples have the same initial point and end point. Nevertheless the two tangential line integrals give different results. We shall later be interested in those vector fields  $\mathbf{V}(\mathbf{x})$ , for which the tangential line integral *only* depends on the initial and end points of the curve  $\mathcal{K}$ . (In Physics such vector fields correspond to the so-called conservative fields.) We have here an example in which this ideal property is *not* satisfied.  $\diamond$

Figure 56: The curve  $\mathcal{K}$  of **Example 4.1.7**.



7) We get

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{-y^3 dx + x^3 dy\} \\
 &= \int_{\frac{\pi}{2}}^{\pi} \{-\sin^3 t \cdot (-\sin t) + (1 + \cos t)^3 \cos t\} dt \\
 &= \int_{\frac{\pi}{2}}^{\pi} \{\sin^4 t + \cos t + 3\cos^2 t + 3\cos^3 t + \cos^4 t\} dt \\
 &= \int_{\frac{\pi}{2}}^{\pi} \left\{ \sin^4 t + \cos^4 t + \left( 2\cos^2 t \cdot \sin^2 t - \frac{1}{2} \sin^2 2t \right) + \cos t + \frac{3}{2} + \frac{3}{2} \cos 2t + 3\cos^3 t \right\} dt \\
 &= \int_{\frac{\pi}{2}}^{\pi} \left\{ (\sin^2 t + \cos^2 t)^2 - \frac{1}{4} + \frac{1}{4} \cos 4t + \cos t + \frac{3}{2} \cos 2t + 3\cos t - 3\sin^2 t \cos t \right\} dt \\
 &= \left[ t - \frac{t}{4} + \frac{1}{16} \sin 4t + \sin t + \frac{3}{2} t + \frac{3}{4} \sin 2t + 3\sin t - \sin^3 t \right]_{\frac{\pi}{2}}^{\pi} \\
 &= \left( 1 - \frac{1}{4} + \frac{3}{2} \right) \frac{\pi}{2} - 4 + 1 = \frac{9\pi}{8} - 3.
 \end{aligned}$$

8) We get

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \left\{ -y^2 dx + a^2 \sinh \frac{x}{a} dy \right\} \\
 &= \int_a^{2a} \left\{ -a^2 \cosh^2 \frac{x}{a} dx + a^2 \sinh \frac{x}{a} \cdot \sinh \frac{x}{a} dx \right\} \\
 &= -a^2 \int_a^{2a} \left\{ \cosh^2 \left( \frac{x}{a} \right) - \sinh^2 \left( \frac{x}{a} \right) \right\} dx = -a^3.
 \end{aligned}$$



Figure 57: The curve  $\mathcal{K}$  of **Example 4.1.8** for  $a = 1$ .Figure 58: The curves  $y = \sqrt{3x}$ ,  $y = \sqrt{3}x$  and  $y = \sqrt{3}x^2$ .

**Example 4.2** Compute the tangential line integral of the vector field

$$\mathbf{V}(x, y) = (2xy, x^6 y^2)$$

along the curve  $\mathcal{K}$  give  $n$  by  $y = ax^b$ ,  $x \in [0, 1]$ . Then find  $a$  such that the line integral becomes independent of  $b$ .

**A** Tangential line integral.

**D** Just use the standard method.

**I** We compute the line integral

$$\begin{aligned} \int_{\mathcal{K}} \mathbf{V}(x, y) \cdot d\mathbf{x} &= \int_{\mathcal{K}} (2xy \, dx + x^6 y^2 \, dy) = \int_0^1 \{2x a x^b + x^6 a^2 x^{2b} \cdot ab x^{b-1}\} \, dx \\ &= \int_0^1 \{2a x^{b+1} + a^3 b x^{3b+5}\} \, dx = \frac{2a}{b+2} + \frac{a^3 b}{3(b+2)} = \frac{a(a^2 b + 6)}{3(b+2)}. \end{aligned}$$


Assume that this result is independent of  $b$ . Then  $b+2$  must be proportional to  $a^2 b + 6$ , so  $a^2 = 3$ .

According to the convention  $a > 0$ , thus  $a = \sqrt{3}$ . By choosing this  $a$  we get

$$\int_{\mathcal{K}} \mathbf{V}(x, y) \cdot d\mathbf{x} = \frac{\sqrt{3}(3b+6)}{3(b+2)} = \sqrt{3},$$

which is independent of  $b$ .

Please click the advert



### Masters in Management



Designed for high-achieving graduates across all disciplines, London Business School's Masters in Management provides specific and tangible foundations for a successful career in business.

This 12-month, full-time programme is a business qualification with impact. In 2010, our MiM employment rate was 95% within 3 months of graduation\*; the majority of graduates choosing to work in consulting or financial services.

As well as a renowned qualification from a world-class business school, you also gain access to the School's network of more than 34,000 global alumni – a community that offers support and opportunities throughout your career.

For more information visit [www.london.edu/mm](http://www.london.edu/mm), email [mim@london.edu](mailto:mim@london.edu) or give us a call on **+44 (0)20 7000 7573**.

\* Figures taken from London Business School's Masters in Management 2010 employment report

**Example 4.3** Compute in each of the following cases the tangential line integral

$$\int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x}$$

of the vector field  $\mathbf{V}$  along the space curve  $\mathcal{K}$ , which is given by the parametric description

$$\mathcal{K} = \{\mathbf{x} \in \mathbb{R}^3 \mid \mathbf{x} = \mathbf{r}(t), t \in I\}.$$

- 1) The vector field is  $\mathbf{V}(x, y, z) = (y^2 - z^2, 2yz, -x^2)$ , and the curve  $\mathcal{K}$  is given by  $\mathbf{r}(t) = (t, t^2, t^3)$  for  $t \in I$ .
- 2) The vector field is  $\mathbf{V}(x, y, z) = \left(\frac{1}{x+z}, y+z, \frac{2}{x+y+z}\right)$ , and the curve  $\mathcal{K}$  is given by  $\mathbf{r}(t) = (t, t^2, t^3)$  for  $t \in [1, 2]$ .
- 3) The vector field is  $\mathbf{V}(x, y, z) = (3x^2 - 6yz, 2y + 3xz, 1 - 4xyz^2)$ , and the curve  $\mathcal{K}$  is given by  $\mathbf{r}(t) = (t, t^2, t^3)$  for  $t \in [0, 1]$ .
- 4) The vector field is  $\mathbf{V}(x, y, z) = (3x^2 - 6yz, 2y + 3xz, 1 - 4xyz^2)$ , and the curve  $\mathcal{K}$  is given by  $\mathbf{r}(t) = (t, t, t)$  for  $t \in [0, 1]$ .
- 5) The vector field is  $\mathbf{V}(x, y, z) = (3x^2 - 6yz, 2y + 3xz, 1 - 4xyz^2)$ , and the curve  $\mathcal{K}$  is given by

$$\mathbf{r}(t) = \begin{cases} (0, 0, t), & \text{for } t \in [0, 1], \\ (0, t-1, 1), & \text{for } t \in [1, 2], \\ (t-2, 1, 1), & \text{for } t \in [2, 3]. \end{cases}$$

- 6) The vector field is  $\mathbf{V}(x, y, z) = (x, y, xz - y)$ , and the curve  $\mathcal{K}$  is given by  $\mathbf{r}(t) = (t, 2t, 4t)$  for  $t \in [0, 1]$ .
- 7) The vector field is  $\mathbf{V}(x, y, z) = (2x + yz, 2y + xz, 2z + xy)$ , and the curve  $\mathcal{K}$  is given by
 
$$\mathbf{r}(t) = (a(\cosh t) \cos t, a(\cosh t) \sin t, at) \quad \text{for } t \in [0, 2\pi].$$
- 8) The vector field is  $\mathbf{V}(x, y, z) = (y^2 - z^2, 2yz, -x^2)$ , and the curve  $\mathcal{K}$  is given by  $\mathbf{r}(t) = (t, t, t)$  for  $t \in [0, 1]$ .

**A** Tangential line integrals in space.

**D** Insert the parametric descriptions and compute the tangential line integral. Notice that **Example 4.3.7** is a gradient field, so it is in this case possible to find the integral directly.

**I** 1) We get

$$\begin{aligned} \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{(y^2 - z^2)dx + 2yz dy - x^2 dz\} \\ &= \int_0^1 \{(t^4 - t^6) + 2t^2 \cdot t^3 \cdot 2t - t^3 \cdot 3t^2\} dt \\ &= \int_0^1 \{3t^6 - 2t^4\} dt = \frac{3}{7} - \frac{2}{5} = \frac{1}{35}. \end{aligned}$$

2) Here

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \left\{ \frac{1}{x+z} dx + (y+z) dy + \frac{2}{x+y+z} dz \right\} \\
 &= \int_1^2 \left\{ \frac{1}{t+t^3} + (t+t^3) + \frac{6t^2}{2t+t^3} \right\} dt \\
 &= \int_1^2 \left\{ \frac{1}{t} - \frac{t}{1+t^2} + \frac{6t}{2+t^2} + t+t^3 \right\} dt \\
 &= \left[ \ln t - \frac{1}{2} \ln(1+t^2) + 3 \ln(2+t^2) + \frac{t^2}{2} + \frac{t^4}{4} \right]_1^2 \\
 &= \ln 2 - \frac{1}{2} \ln 5 + 3 \ln 6 - \frac{1}{2} \ln 2 - 3 \ln 3 + \frac{4}{2} + \frac{16}{4} - \frac{1}{2} - \frac{1}{4} \\
 &= \frac{9}{2} \ln 2 + \frac{1}{2} \ln 5 + \frac{21}{4} = \frac{21}{4} + \frac{1}{2} \ln \frac{512}{5}.
 \end{aligned}$$

3) First notice that for any curve,

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{(3x^2 - 6yz)dx + (2y + 3xz)dy + (1 - 4xyz^2)dz\} \\
 (2) \quad &= \int_{\mathcal{K}} d(x^3 + y^2 + z) - \int_{\mathcal{K}} z\{6ydx - 3xdy + 4xyzdz\}.
 \end{aligned}$$

Such a rearrangement can also be used advantageously to **Example 4.3.3**, **Example 4.3.4** and **Example 4.3.5**.

When we apply (2), we get

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= [x^3 + y^2 + z]_{(0,0,0)}^{(1,1,1)} - \int_0^1 t^3 \{6t^2 - 3t \cdot 2t + t^6 \cdot 3t^2\} dt \\
 &= 3 - \int_0^1 12t^{11} dt = 3 - 1 = 2.
 \end{aligned}$$

ALTERNATIVELY, it follows by a direct insertion that

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{(3x^2 - 6yz)dx + (2y + 3xz)dy + (1 - 4xyz^2)dz\} \\
 &= \int_0^1 \{(3t^2 - 6t^2 \cdot t^3) + (2t^2 + 3t \cdot t^3)2t + (1 - 4t \cdot t^2 \cdot t^6)3t^2\} dt \\
 &= \int_0^1 \{3t^2 - 6t^5 + 4t^3 + 6t^5 + 3t^2 - 12t^{11}\} dt \\
 &= \int_0^1 (6t^2 + 4t^3 - 12t^{11}) dt = [2t^3 + t^4 - t^{12}]_0^1 = 2.
 \end{aligned}$$

4) The vector field is the same as in **Example 4.3.3**. Then we get by (2),

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= [x^3 + y^2 + z]_{(0,0,0)}^{(1,1,1)} - \int_0^1 (6t^2 - 3t^2 + 4t^4) dt \\
 &= 3 - \int_0^1 (3t^2 + 4t^4) dt = 3 - 1 - \frac{4}{5} = \frac{6}{5}.
 \end{aligned}$$

ALTERNATIVELY, it follows by a direct insertion that

$$\begin{aligned}\int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_0^1 \{(3t^2 - 6t^2) + (2t + 3t^2) + 1 - 4t^4\} dt \\ &= \int_0^1 (1 + 2t - 4t^4) dt = 1 + 1 - \frac{4}{5} = \frac{6}{5}.\end{aligned}$$

- 5) The vector field is the same as in **Example 4.3.3**. When we apply (2) and just check that  $\mathbf{r}(t)$  is a continuous curve, we get

$$\begin{aligned}\int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= [x^3 + y^2 + z]_{(0,0,0)}^{(1,1,1)} - \int_{\mathcal{K}} z\{6ydx - 3xdy + 4xyzdx\} \\ &= 3 - \int_0^1 0 dt - \int_1^2 +20 dt - \int_2^3 1 \cdot 6 dt = 3 - 6 = -3.\end{aligned}$$

ALTERNATIVELY, it follows by direct insertion that

$$\begin{aligned}\int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{(3x^2 - 6yz)dx + (2y + 3xz)dy + (1 - 4xyz^2)dz\} \\ &= \int_0^1 (1 - 4 \cdot 0)dt + \int_1^2 \{2(t-1) + 0\}dt + \int_2^3 \{3(t-2)^2 - 6\}dt \\ &= [t]_0^1 + 2 \left[ \frac{1}{2} (t-1)^2 \right]_1^2 + 3 \left[ \frac{1}{3} (t-2)^3 - 2t \right]_2^3 \\ &= 1 + 1 + 1 - 3 \cdot 2 \cdot 3 + 3 \cdot 2 \cdot 2 = 3(1 - 6 + 4) = -3.\end{aligned}$$

- 6) Here we get by insertion,

$$\begin{aligned}\int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{xdx + ydy + (xz - y)dz\} \\ &= \int_0^1 \{t + 2t \cdot 2 + (t \cdot 4t - 2t) \cdot 4\}dt \\ &= \int_0^1 (t + 4t + 16t^2 - 8t)dt = \int_0^1 (16t^2 - 3t)dt \\ &= \frac{16}{3} - \frac{3}{2} = \frac{32 - 9}{6} = \frac{23}{6}.\end{aligned}$$

- 7) It follows immediately that

$$\begin{aligned}\int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{(2x + yz)dx + (2y + xz)dy + (2xyz)dz\} \\ &= \int_{\mathcal{K}} \{d(x^2 + y^2 + z^2) + (yzdx + xzdy + xydz)\} \\ &= \int_{\mathcal{K}} d(x^2 + y^2 + z^2 + xyz) = [x^2 + y^2 + z^2 + xyz]_{(x,y,z)=(a,0,0)}^{a(\cosh 2\pi, 0, 2\pi)} \\ &= a^2 \cosh^2 2\pi + 4a^2\pi^2 - a^2 = a^2(4\pi^2 + \sinh^2 2\pi).\end{aligned}$$

ALTERNATIVELY, we get by the parametric description

$$\mathbf{r}(t) = a(\cosh t \cdot \cos t, \cosh t \cdot \sin t, t), \quad t \in [0, 2\pi],$$

that

$$\mathbf{r}'(t) = a(\sinh t \cdot \cos t - \cosh t \cdot \sin t, \sinh t \cdot \sin t + \cosh t \cdot \cos t, 1),$$

thus

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{(2x + yz)dx + (2y + xz)dy + (2z + xy)dz\} \\
 &= \int_0^{2\pi} (2a \cosh t \cos t + a^2 t \cosh t \sin t) a (\sinh t \cos t - \cosh t \sin t) dt \\
 &\quad + \int_0^{2\pi} (2a \cosh t \sin t + a^2 \cosh t \cos t) a (\sinh t \sin t + \cosh t \cos t) dt \\
 &\quad + \int_0^{2\pi} (2at + a^2 \cosh^2 t \cdot \cos t \cdot \sin t) a dt \\
 &= a^2 \cdot (\dots) + a^3 \cdot (\dots).
 \end{aligned}$$

Then the easiest method is to reduce and use that

$$\cos t = \frac{1}{2} (e^{it} + e^{-it}), \quad \sin t = \frac{1}{2i} (e^{it} - e^{-it}),$$

and similarly for  $\cosh t$  and  $\sinh t$ . We finally obtain the result by a partial integration.

Please click the advert



You're full of *energy*  
and *ideas*. And that's  
just what we are looking for.

Looking for a career where your ideas could really make a difference? UBS's Graduate Programme and internships are a chance for you to experience for yourself what it's like to be part of a global team that rewards your input and believes in succeeding together.

Wherever you are in your academic career, make your future a part of ours by visiting [www.ubs.com/graduates](http://www.ubs.com/graduates).

[www.ubs.com/graduates](http://www.ubs.com/graduates)



© UBS 2010. All rights reserved.

The variants above are somewhat sophisticated, so we proceed here by first calculating the coefficient of  $a^2$ :

$$\begin{aligned} & \int_0^{2\pi} 2 \cosh t \cdot t (\sinh t \cdot \cos t - \cosh t \cdot \sin t) dt \\ & \quad + \int_0^{2\pi} 2 \cosh t \cdot \sin t (\sinh t \cdot \sin t + \cosh t \cdot \cos t) dt + \int_0^{2\pi} 2t dt \\ & = 2 \int_0^{2\pi} \cosh t \cdot \sinh t dt + \int_0^{2\pi} 2t dt = [\sinh^2 t + t^2]_0^{2\pi} = 4\pi^2 + \sinh^2 2\pi. \end{aligned}$$

Then we find the coefficient of  $a^3$ :

$$\begin{aligned} & \int_0^{2\pi} t \{ \cosh t \sinh t \sin t \cos t - \cosh^2 t \sin^2 t \} dt \\ & \quad + \int_0^{2\pi} t \{ \cosh t \sinh t \sin t \cos t + \cosh^2 t \cos^2 t \} dt + \int_0^{2\pi} \cosh^2 t \cos t \sin t dt \\ & = \int_0^{2\pi} t (\cosh t \sinh t \sin 2t + \cosh^2 t \cos 2t) dt + \frac{1}{2} \int_0^{2\pi} \cosh^2 t \sin 2t dt. \end{aligned}$$

Notice that

$$\frac{d}{dt} \left\{ \frac{1}{2} \cosh^2 t \cdot \sin 2t \right\} = \cosh t \cdot \sinh t \cdot \sin 2t + \cosh^2 t \cdot \cos 2t,$$

so the whole expression can then be written

$$\begin{aligned} & \int_0^{2\pi} \left\{ t \frac{d}{dt} \left( \frac{1}{2} \cosh^2 t \sin 2t \right) + \frac{dt}{dt} \cdot \left( \frac{1}{2} \cosh^2 t \sin 2t \right) \right\} dt \\ & = \int_0^{2\pi} \frac{d}{dy} \left( \frac{t}{2} \cosh^2 t \sin 2t \right) dt = \left[ \frac{t}{2} \cosh^2 t \cdot \sin 2t \right]_0^{2\pi} = 0. \end{aligned}$$

As a conclusion we get

$$\int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} = a^2(4\pi^2 + \sinh^2 2\pi) + 0 \cdot a^3 = a^2(4\pi^2 + \sinh^2 2\pi).$$

8) Here we get [cf. also **Example 4.3.1**, where the vector field is the same]

$$\begin{aligned} \int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{ (y^2 - z^2) dx + 2yz dy - x^2 dz \} \\ &= \int_0^1 \{ (t^2 - t^2) + 2t^2 - t^2 \} dt = \int_0^1 t^2 dt = \frac{1}{3}. \end{aligned}$$

**Example 4.4** Compute in each of the following cases the tangential line integral of the given vector field  $\mathbf{V}$  along the given curve  $\mathcal{K}$ .

1) The vector field is  $\mathbf{V}(x, y) = (x + y, x - y)$ , and the curve  $\mathcal{K}$  is the ellipse of centrum  $(0, 0)$  and half axes  $a, b$ , run through in the positive orientation of the plane.

2) The vector field is  $\mathbf{V}(x, y) = \left( \frac{1}{|x| + |y|}, \frac{1}{|x| + |y|} \right)$ , and the curve  $\mathcal{K}$  is the square defined by its vertices

$$(1, 0), \quad (0, 1), \quad (-1, 0), \quad (0, -1),$$

in the positive orientation of the plane.

3) The vector field is  $\mathbf{V}(x, y) = (x^2 - y, y^2 + x)$ , and the curve  $\mathcal{K}$  is the line segment from  $(0, 1)$  to  $(1, 2)$ .

4) The vector field is  $\mathbf{V}(x, y) = (x^2 - y^2, y^2 + x)$ , and the curve  $\mathcal{K}$  is the broken line from  $(0, 1)$  over  $(1, 1)$  to  $(1, 2)$ .

5) The vector field is  $\mathbf{V}(x, y) = (x^2 - y, y^2 + x)$ , and the curve  $\mathcal{K}$  is that part of the parabola of equation  $y = 1 + x^2$ , which has the initial point  $(0, 1)$  and the end point  $(1, 2)$ .

6) The vector field is  $\mathbf{V}(x, y, z) = (yz, xz, x(y + 1))$ , and the curve  $\mathcal{K}$  is the triangle given by its vertices

$$(0, 0, 0), \quad (1, 1, 1), \quad (-1, 1, -1),$$

and run through in given sequence.

7) The vector field is  $\mathbf{V}(x, y, z) = (\sin y, \sin z, \sin x)$ , and the curve  $\mathcal{K}$  is the line segment from  $(0, 0, 0)$  to  $(\pi, \pi, \pi)$ .

8) The vector field is  $\mathbf{V}(x, y, z) = (z, x, -y)$ , and the curve  $\mathcal{K}$  is the quarter circle from  $(a, 0, 0)$  to  $(0, 0, a)$  followed by another quarter circle from  $(0, 0, a)$  to  $(0, a, 0)$ , both of centrum  $(0, 0, 0)$ .

**A** Tangential line integrals in the 2-dimensional and the 3-dimensional space.

**D** Sketch in the 2-dimensional case the curve  $\mathcal{K}$ . Then check if any part of  $\mathbf{V}(\mathbf{x}) \cdot d\mathbf{x}$  can be sorted out as a total differential. Finally, insert the parametric description and compute.

**I** 1) As  $\mathcal{K}$  is a closed curve, we get

$$\int_{\mathcal{K}} \mathbf{V}(\mathbf{x}) \cdot d\mathbf{x} = \int_{\mathcal{K}} \{(x + y)dx + (x - y)dy\} = \int_{\mathcal{K}} d\left(\frac{1}{2}x^2 + xy - \frac{1}{2}y^2\right) = 0,$$

because  $\mathbf{V} \cdot d\mathbf{x}$  is a total differential.



ALTERNATIVELY,  $\mathcal{K}$  has e.g. the parametric description

$$(x, y) = \mathbf{r}(t) = (a \cos t, b \sin t), \quad t \in [0, 2\pi],$$

thus

$$\mathbf{r}'(t) = (-a \sin t, b \cos t).$$




Figure 59: A possible curve  $\mathcal{K}$  in **Example 4.4.1**.Figure 60: The curve of **Example 4.4.2**.

Then by insertion,

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{(x+y)dx + (x-y)dy\} \\
 &= \int_0^{2\pi} \{(a \cos t + b \sin t)(-a \sin t) + (a \cos t - b \sin t)b \cos t\} dt \\
 &= \int_0^{2\pi} \{-a^2 \cos t \sin t - ab \sin^2 t + ab \cos^2 t - b^2 \sin t \cos t\} dt \\
 &= \int_0^{2\pi} \left\{ ab \cos 2t - \frac{1}{2}(a^2 + b^2) \sin 2t \right\} dt = 0.
 \end{aligned}$$

2) Since  $|x| + |y| = 1$  on  $\mathcal{K}$ , we have

$$\int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} = \int_{\mathcal{K}} \frac{1}{|x| + |y|} (dx + dy) = \int_{\mathcal{K}} 1 d(x+y) = 0.$$

Figure 61: The curve  $\mathcal{K}$  of **Example 4.4.3**

ALTERNATIVELY, and more difficult we can use the parametric description of  $\mathcal{K}$  given by

$$\mathbf{r}(t) = \begin{cases} (1-t, t), & t \in [0, 1], \\ (1-t, 2-t), & t \in [1, 2], \\ (t-3, 2-t), & t \in [2, 3], \\ (t-3, t-4), & t \in [3, 4], \end{cases}$$

hence

$$\mathbf{r}'(t) = \begin{cases} (-1, 1), & t \in ]0, 1[, \\ (-1, -1), & t \in ]1, 2[, \\ (1, -1), & t \in ]2, 3[, \\ (1, 1), & t \in ]3, 4[. \end{cases}$$

Since  $|x| + |y| = 1$  on  $\mathcal{K}$ , we get

$$\begin{aligned} \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} &= \int_{\mathcal{K}} (dx + dy) = \int_0^1 (-1 + 1)dt + \int_1^2 (-1 - 1)dt + \int_2^3 (1 - 1)dt + \int_3^4 (1 + 1)dt \\ &= 0 - 2 + 0 + 2 = 0. \end{aligned}$$

3) First notice that

$$\begin{aligned} \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{(x^2 - y)dx + (y^2 + x)dy\} = \frac{1}{3} \int_{\mathcal{K}} d(x^3 + y^3) + \int_{\mathcal{K}} (-y dx + x dy) \\ &= \frac{1}{3}(8 + 1 - 1) + \int_{\mathcal{K}} (-y dx + x dy), \end{aligned}$$

so

$$(3) \quad \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} = \frac{8}{3} + \int_{\mathcal{K}} (-y dx + x dy)$$

$$(4) \quad = \int_{\mathcal{K}} \{(x^2 - y)dx + (y^2 + x)dy\}.$$

Now we compute **Example 4.4.3**, **Example 4.4.4** and **Example 4.4.5** in the two variants corresponding to (3) and (4), respectively.

A parametric description of  $\mathcal{K}$  is e.g.

$$\mathbf{r}(t) = (t, 1+t), \quad t \in [0, 1],$$

and accordingly,  $\mathbf{r}'(t) = (1, 1)$ .

Please click the advert




Discover the truth at [www.deloitte.ca/careers](http://www.deloitte.ca/careers)

**Deloitte.**

© Deloitte & Touche LLP and affiliated entities.

Download free ebooks at [bookboon.com](http://bookboon.com)

Figure 62: The curve  $\mathcal{K}$  of **Example 4.4.4**.

Then by (3),

$$\int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} = \frac{8}{3} + \int_0^1 \{-(1+t) + t\} dt = \frac{8}{3} - 1 = \frac{5}{3}.$$

ALTERNATIVELY, we get by (4) that

$$\begin{aligned} \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} &= \int_0^1 \{t^2 - (1+t) + (1+t)^2 + t\} dt = \int_0^1 \{(1+t)^2 + t^2 - 1\} dt \\ &= \left[ \frac{1}{3}(1+t)^3 + \frac{1}{3}t^3 - t \right]_0^1 = \frac{8+1-1}{3} - 1 = \frac{5}{3}. \end{aligned}$$

4) It follows from (3) that


$$\int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} = \frac{8}{3} + \int_0^1 (-1) dx + \int_1^2 1 dy = \frac{8}{3} - 1 + 1 = \frac{8}{3}.$$

ALTERNATIVELY, we get by (4),

$$\begin{aligned} \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} &= \int_0^1 (x^2 - 1) dx + \int_1^2 (y^2 + 1) dy = \left[ \frac{1}{3}x^3 - x \right]_0^1 + \left[ \frac{1}{3}y^3 + y \right]_1^2 \\ &= \frac{1}{3} - 1 + \frac{8}{3} + 2 - \frac{1}{3} - 1 = \frac{8}{3}. \end{aligned}$$

5) By (3),

$$\begin{aligned} \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} &= \frac{8}{3} + \int_0^1 \{(-1 - x^2) + x \cdot 2x\} dx \\ &= \frac{8}{3} + \int_0^1 (x^2 - 1) dx = \frac{8}{3} + \left[ \frac{1}{3}x^3 - x \right]_0^1 = \frac{8}{3} + \frac{1}{3} - 1 = 2. \end{aligned}$$

Figure 63: The curve  $\mathcal{K}$  of **Example 4.4.5**.

ALTERNATIVELY, by (4),

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{x^2 - x^2 - 1 + [(x^2 + 1)^2 + x] \cdot 2x\} dx \\
 &= \int_{\mathcal{K}} \{2x^5 + 4x^3 + 2x^2 + 2x - 1\} dx \\
 &= \left[ \frac{1}{3} x^6 + x^3 + \frac{2}{3} x^3 + x^2 - x \right]_0^1 = \frac{1}{3} + 1 + \frac{2}{3} + 1 - 1 = 2.
 \end{aligned}$$

6) Here a parametric description is e.g. given by

$$\mathbf{r}(t) = \begin{cases} (t, t, t), & t \in [0, 1], \\ (3 - 2t, 1, 3 - 2t), & t \in [1, 2], \\ (t - 3, -t + 3, t - 3), & t \in [2, 3], \end{cases}$$

hence

$$\mathbf{r}'(t) = \begin{cases} (1, 1, 1), & t \in ]0, 1[, \\ (-2, 0, -2), & t \in ]1, 2[, \\ (1, -1, 1), & t \in ]2, 3[. \end{cases}$$

FIRST VARIANT. We get by direct insertion,

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} &= \int_{\mathcal{K}} \{yz \, dx + xz \, dy + x(y + 1) \, dz\} \\
 &= \int_0^1 (t^2 + t^2 + t^2 + t) dt + \int_1^2 \{1 \cdot (3 - 2t) \cdot (-2) + (3 - 2t) \cdot 2 \cdot (-2)\} dt \\
 &\quad + \int_2^3 \{(-t + 3)(t - 3) \cdot 1 + (t - 3)^2 \cdot (-1) + (t - 3)(-t + 3) \cdot 1 + t - 3\} dt \\
 &= \int_0^1 (3t^2 + t) dt - 6 \int_1^2 (3 - 2t) dt - \int_2^3 \{3(t - 3)^2 - (t - 3)\} dt \\
 &= \left[ t^3 + \frac{1}{2} t^2 \right]_0^1 + 6 \left[ t^2 - 3t \right]_1^2 - \left[ (t - 3)^3 - \frac{1}{2} (t - 3)^2 \right]_2^3 \\
 &= 1 + \frac{1}{2} + 6(4 - 6 - 1 + 3) + \left( -1 - \frac{1}{2} \right) = 0.
 \end{aligned}$$

**2. variant.** Reduction by removing a total differential.

As

$$\mathbf{V} \cdot d\mathbf{x} = yz \, dx + xz \, dy + xy \, dz + x \, dz = d(xyz) + x \, dz,$$

and as  $\mathcal{K}$  is a closed curve, we have  $\int_{\mathcal{K}} d(xyz) = 0$ , so the calculations are simplified by removing  $d(xyz)$ :

$$\begin{aligned} \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} &= \int_{\mathcal{K}} d(xyz) + \int_{\mathcal{K}} x \, dz = 0 + \int_{\mathcal{K}} x \, dz \\ &= \int_0^1 t \, dt + \int_1^2 (3-2t) \cdot (-2) \, dt + \int_2^3 (t-3) \, dt \\ &= \left[ \frac{1}{2} t^2 \right]_0^1 + \int_1^2 (4t-6) \, dt + \left[ \frac{1}{2} (t-3)^2 \right]_2^3 \\ &= \frac{1}{2} + [2t^2 - 6t]_1^2 - \frac{1}{2} = 8 - 12 - 2 + 6 = 0. \end{aligned}$$

REMARK. The expressions will be even simpler, if we do not insist on that the parametric intervals  $[0, 1]$ ,  $[1, 2]$ ,  $[2, 3]$  should follow each other. Instead one can split  $\mathcal{K}$  into three subcurves

$$\begin{aligned} \mathcal{K}_1 : \quad \mathbf{r}_1(t) &= (t, t, t), & t &\in [0, 1], \\ \mathcal{K}_2 : \quad \mathbf{r}_2(t) &= (1-2t, 1, 1-2t), & t &\in [0, 1], \\ \mathcal{K}_3 : \quad \mathbf{r}_3(t) &= (t-1, 1-t, t-1), & t &\in [0, 1], \end{aligned}$$

where

$$\begin{aligned} \mathcal{K}_1 : \quad \mathbf{r}'_1(t) &= (1, 1, 1), & t &\in ]0, 1[; \\ \mathcal{K}_2 : \quad \mathbf{r}'_2(t) &= (-2, 0, 2), & t &\in ]0, 1[; \\ \mathcal{K}_3 : \quad \mathbf{r}'_3(t) &= (1, -1, 1), & t &\in ]0, 1[. \end{aligned}$$

We obtain that the three line integrals can be joined like in the second variant above:

$$\begin{aligned} \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} &= \dots = \int_{\mathcal{K}} x \, dz = \int_{\mathcal{K}_1} x \, dz + \int_{\mathcal{K}_2} x \, dz + \int_{\mathcal{K}_3} x \, dz \\ &= \int_0^1 t \, dt + \int_0^1 (1-2t) \cdot (-2) \, dt + \int_0^1 (t-1) \, dt \\ &= 3 \int_0^1 (2t-1) \, dt = 3 [t^2 - t]_0^1 = 0. \quad \diamond \end{aligned}$$

7) The most obvious parametric description is here

$$\mathbf{r}(t) = t(1, 1, 1), \quad \text{med } \mathbf{r}'(t) = (1, 1, 1), \quad t \in [0, \pi].$$

Thus we can put  $x = y = z = t$  everywhere. Then

$$\int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} = \int_{\mathcal{K}} \{\sin y \, dx + \sin z \, dy + \sin x \, dz\} = \int_0^\pi 3 \sin t \, dt = [-3 \cos t]_0^\pi = 6.$$

8) If we call the two curve segments for  $\mathcal{K}_1$  and  $\mathcal{K}_2$ , then the most obvious parametric description is

$$\begin{aligned} \mathcal{K}_1 : \quad a(\cos t, 0, \sin t), & \quad t \in \left[0, \frac{\pi}{2}\right], \\ \mathcal{K}_2 : \quad a(0, \sin t, \cos t), & \quad t \in \left[0, \frac{\pi}{2}\right]. \end{aligned}$$

Then by insertion,

$$\begin{aligned}
 \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} &= \int_{\mathcal{K}} (z \, dx + x \, dy - y \, dz) \\
 &= a^2 \int_0^{\frac{\pi}{2}} (\sin t \cdot (-\sin t)) \, dt + a^2 \int_0^{\frac{\pi}{2}} (-\sin t) \cdot (-\sin t) \, dt \\
 &= -a^2 \int_0^{\frac{\pi}{2}} \sin^2 t \, dt + a^2 \int_0^{\frac{\pi}{2}} \sin^2 t \, dt = 0.
 \end{aligned}$$

Please click the advert

It's only an opportunity if you act on it

[IKEA.SE/STUDENT](http://IKEA.SE/STUDENT)

**IKEA**

© Inter IKEA Systems B.V. 2009

**Example 4.5** Find in each of the following cases a function

$$\Phi(x, y) = \int_{\mathcal{K}} \mathbf{V}(\tilde{\mathbf{x}}) \cdot d\tilde{\mathbf{x}},$$

to the given vector field  $\mathbf{V} : A \rightarrow \mathbb{R}^2$ , where  $\mathcal{K}$  is the broken line which runs from  $(0, 0)$  over  $(x, 0)$  to  $(x, y)$ . Check if  $\Phi$  is defined in all of  $A$ , and compute finally the gradient  $\nabla\Phi$ .

1) The vector field  $\mathbf{V}(x, y) = (y^2 - 2xy, -x^2 + 2xy)$  is defined in  $A = \mathbb{R}^2$ .

2) The vector field  $\mathbf{V}(x, y) = \left( \frac{1}{\sqrt{y^2 - x^2 + 1}}, x \right)$  is defined in

$$A = \{(x, y) \mid -\sqrt{1 + y^2} < x < \sqrt{1 + y^2}\}.$$

3) The vector field  $\mathbf{V}(x, y) = \left( \frac{x}{\sqrt{1 - x^2 - y^2}}, \frac{y}{\sqrt{1 - x^2 - y^2}} \right)$  is defined in the disc  $A$  given by  $x^2 + y^2 < 1$ .

4) The vector field  $\mathbf{V}(x, y) = \left( \frac{x - 1}{\sqrt{(x - 1)^2 + y^2}}, \frac{y}{\sqrt{(x - 1)^2 + y^2}} \right)$  in the set  $A$  given by  $(x, y) \neq (1, 0)$ .

5) The vector field  $\mathbf{V}(x, y) = (\cos y, \cos x)$  is defined in  $A = \mathbb{R}^2$ .

6) The vector field  $\mathbf{V}(x, y) = (\cos(xy), 0)$  is defined in  $A = \mathbb{R}^2$ .

7) The vector field  $\mathbf{V}(x, y) = (x^2 + y^2, xy)$  is defined in  $A = \mathbb{R}^2$ .

8) The vector field  $\mathbf{V}(x, y) = (x^2 + y^2, 2xy)$  is defined in  $A = \mathbb{R}^2$ .

**A** Tangential line integrals.

**D** Remove whenever possible total differentials. Integrate along a broken line. Finally, compute the gradient  $\nabla\Phi$ .

**I** 1) We get by inspection,

$$\begin{aligned} \Phi(x, y) &= \int_{\mathcal{K}} \mathbf{V}(\tilde{\mathbf{x}}) \cdot d\tilde{\mathbf{x}} = \int_{\mathcal{K}} \{(\tilde{y}^2 - 2\tilde{x}\tilde{y})d\tilde{x} + (-\tilde{x}^2 + 2\tilde{x}\tilde{y})d\tilde{y}\} \\ &= \int_{\mathcal{K}} d(\tilde{x}\tilde{y}^2 - \tilde{x}^2\tilde{y}) = xy^2 - x^2y \quad (= xy(y - x)). \end{aligned}$$

ALTERNATIVELY,


$$\Phi(x, y) = \int_{\mathcal{K}} \{(\tilde{y}^2 - 2\tilde{x}\tilde{y})d\tilde{x} + (-\tilde{x}^2 + 2\tilde{x}\tilde{y})d\tilde{y}\} = \int_0^x 0 dt + \int_0^y (-x^2 + 2xt)dt = xy^2 - x^2y.$$

Finally,

$$\nabla\Phi = (y^2 - 2xy, 2xy - x^2) = \mathbf{V}(x, y),$$

and  $\Phi$  is defined in all of  $A = \mathbb{R}^2$ .



Figure 64: The domains of  $\mathbf{V}(x, y)$  and  $\Phi(x, y)$ .

- 2) The domain  $A$  for  $\mathbf{V}(x, y)$  lies between the two hyperbolic branches given by  $x^2 - y^2 = 1$ . The domain  $\tilde{A}$  of  $\Phi(x, y)$  is smaller, in fact the points lying between the two vertical lines  $x = \pm 1$ , because we can only reach these by curves of the type  $\mathcal{K}$ . (The curve  $\mathcal{K}$  must never leave  $A$ , because we require that  $\mathbf{V}$  is defined).

We get for  $(x, y) \in \tilde{A}$ ,

$$(5) \quad \Phi(x, y) = \int_0^x \frac{1}{\sqrt{1-t^2}} dt + \int_0^y x dt = \text{Arcsin } x + xy.$$

The function  $\Phi$  is only defined in  $\tilde{A}$ . In this subset of  $A$  we get

$$\nabla \Phi(x, y) = \left( \frac{1}{\sqrt{1-x^2}} + y, x \right) \neq \mathbf{V}(x, y).$$

In particular,  $\mathbf{V}(x, y)$  is not a gradient field.

REMARK. Formula (5) is a mindless insertion into one of the solution formulæ for this type of problems. It cannot be applied here because the *assumptions* of it are *not* fulfilled.  $\diamond$

- 3) Here we get

$$\begin{aligned} \Phi(x, y) &= \int_{\mathcal{K}} \left\{ \frac{x}{\sqrt{1-x^2-y^2}} dx + \frac{y}{\sqrt{1-x^2-y^2}} dy \right\} \\ &= \int_{\mathcal{K}} d\left(-\sqrt{1-x^2-y^2}\right) = 1 - \sqrt{1-x^2-y^2}. \end{aligned}$$

ALTERNATIVELY we get for  $x^2 + y^2 < 1$  by integration along the broken line that

$$\begin{aligned} \Phi(x, y) &= \int_0^x \frac{t}{\sqrt{1-t^2}} dt + \int_0^y \frac{t}{\sqrt{1-x^2-t^2}} dt = \left[-\sqrt{1-t^2}\right]_0^x + \left[-\sqrt{1-x^2-t^2}\right]_0^y \\ &= 1 - \sqrt{1-x^2} + \sqrt{1-x^2} - \sqrt{1-x^2-y^2} = 1 - \sqrt{1-x^2-y^2}. \end{aligned}$$

It follows immediately that

$$\nabla \Phi(x, y) = \left( \frac{x}{\sqrt{1-x^2-y^2}}, \frac{y}{\sqrt{1-x^2-y^2}} \right) = \mathbf{V}(x, y),$$




Figure 65: The domain  $\tilde{A}$  of  $\Phi$  lies to the left of the dotted line  $x = 1$ .

and that  $\Phi$  is defined in all of  $A$ .

4) In this case we have for any curve  $\mathcal{K}$  from  $(0,0)$  in  $A$  that

$$\begin{aligned}\Phi(x, y) &= \int_{\mathcal{K}} \left\{ \frac{x-1}{\sqrt{(x-1)^2 + y^2}} dx + \frac{y}{\sqrt{(x-1)^2 + y^2}} dy \right\} \\ &= \int_{\mathcal{K}} d\left(\sqrt{(x-1)^2 + y^2}\right) = \sqrt{(x-1)^2 + y^2} - 1.\end{aligned}$$

If we only integrate along curves  $\mathcal{K}$  of this type, then we can only reach points in

$$\tilde{A} = \{(x, y) \mid x < 1, y \in \mathbb{R}\}.$$

By integration along a broken line in this domain,

$$\begin{aligned}\Phi(x, y) &= \int_0^x \frac{t-1}{\sqrt{(t-1)^2 + 0^2}} dt + \int_0^y \frac{t}{\sqrt{(x-1)^2 + t^2}} dt \\ &= \int_0^x \frac{t-1}{|t-1|} dt + \left[ \sqrt{(x-1)^2 + t^2} \right]_0^y \\ &= \int_0^x (-1) dt + \left[ \sqrt{(x-1)^2 + y^2} - \sqrt{(x-1)^2} \right] \quad (\text{because } t < x < 1) \\ &= -x - |x-1| + \sqrt{(x-1)^2 + y^2} = x - (1-x) + \sqrt{(x-1)^2 + y^2} \\ &= \sqrt{(x-1)^2 + y^2} - 1.\end{aligned}$$

It follows that  $\nabla \Phi = \mathbf{V}$  and that  $\Phi$  can be extended to all of  $A$ .

5) When we integrate along the broken line

$$(0, 0) \longrightarrow (x, 0) \longrightarrow (x, y)$$

we get

$$\Phi(x, y) = \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} = \int_0^x \cos 0 dt + \int_0^y \cos x dt = x + y \cos x,$$

which is defined in all of  $\mathbb{R}^2$ . Here,

$$\nabla \Phi(x, y) = (1 - y \sin x, \cos x) \neq \mathbf{V}.$$

It is seen that  $\mathbf{V}$  is not a gradient field.

6) When we integrate along the broken line

$$(0, 0) \longrightarrow (x, 0) \longrightarrow (x, y)$$

we get in all of  $\mathbb{R}^2$ ,

$$\Phi(x, y) = \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} = \int_0^x \cos(t \cdot 0) dt + 0 = x,$$

where  $\nabla \Phi = (1, 0) \neq \mathbf{V}$ , so  $\mathbf{V}$  is not a gradient field.

7) When we integrate along the broken line

$$(0, 0) \longrightarrow (x, 0) \longrightarrow (x, y)$$

we get in all of  $\mathbb{R}^2$ ,

$$\Phi(x, y) = \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} = \int_0^x (t^2 + 0^2) dt + \int_0^y xt dt = x^3 + \frac{1}{2} xy^2,$$

Please click the advert

## YOUR CHANCE TO CHANGE THE WORLD

Here at Ericsson we have a deep rooted belief that the innovations we make on a daily basis can have a profound effect on making the world a better place for people, business and society. Join us.

In Germany we are especially looking for graduates as Integration Engineers for

- Radio Access and IP Networks
- IMS and IPTV

We are looking forward to getting your application!  
To apply and for all current job openings please visit our web page: [www.ericsson.com/careers](http://www.ericsson.com/careers)

ericsson.  
com



where

$$\nabla\Phi = \left(3x^2 + \frac{1}{2}y^2, xy\right) \neq \mathbf{V}(x, y).$$

It follows that  $\mathbf{V}$  is not a gradient field.

8) When we integrate along the broken line

$$(0, 0) \longrightarrow (x, 0) \longrightarrow (x, y)$$

we get in  $\mathbb{R}^2$

$$\Phi(x, y) = \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} = \int_0^x (t^2 + 0^2) dt + \int_0^y 2xt dt = \frac{x^3}{3} + xy^2,$$

where

$$\nabla\Phi = (x^2 + y^2, 2xy) = \mathbf{V}(x, y).$$

In this case  $\mathbf{V}(x, y)$  is a gradient field.

**Example 4.6** Compute in each of the following cases the tangential line integral of the given vector field  $\mathbf{V} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  along the described curve  $\mathcal{K}$ .

- 1) The vector field  $\mathbf{V}(x, y) = \nabla \left( \frac{1}{2}x^2 + xy - \frac{1}{2}y^2 \right)$  along the ellipse  $\mathcal{K}$  of centrum  $(0, 0)$  and half axes  $a, b$ , in the positive orientation of the plane.
- 2) The vector field  $\mathbf{V}(x, y) = \nabla (x^4 + \ln(1 + y))$  along the arc of the parabola  $\mathcal{K}$  given by  $y = x^2$ ,  $x \in [-1, 3]$ .
- 3) The vector field  $\mathbf{V}(x, y) = \nabla(x + 2y - \exp(xy))$  along the broken line  $\mathcal{K}$ , which goes from  $(2, 0)$  over  $(1, 2)$  to  $(0, 1)$ .

**A** Line integral of a gradient field.

**D** As  $\mathbf{V}(x, y) = \nabla F$ , the tangential line integral is only depending on the initial point and the end point,

$$\int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{s} = F(\mathbf{x}_s) - F(\mathbf{x}_b).$$

**I** 1) The ellipse is a closed curve, so

$$\int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} = 0.$$

2) The initial point is  $(-1, 1)$ , and the end point is  $(3, 9)$ , hence

$$\int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} = [x^4 + \ln(1 + y)]_{(-1, 1)}^{(3, 9)} = 81 + \ln 10 - 1 - \ln 2 = 80 + \ln 5.$$

3) The initial point is  $(2, 0)$ , and the end point is  $(0, 1)$ , thus

$$\int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} = [x + 2y - \exp(xy)]_{(2,0)}^{(0,1)} = 0 + 2 - 1 - 2 - 0 + 1 = 0.$$

**Example 4.7** Compute in each of the following cases the tangential line integral of the given vector field  $\mathbf{V} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  along the described curve  $\mathcal{K}$ .

1) The vector field  $\nabla(x^2 + yz)$  along the curve  $\mathcal{K}$ , given by

$$\mathbf{r}(t) = (\cos t, \sin t, \sin(2t)), \quad t \in [0, 2\pi].$$

2) The vector field  $\nabla(\cos(xyz))$  along the line segment  $\mathcal{K}$  from  $\left(\pi, \frac{1}{2}, 0\right)$  to  $\left(\frac{1}{2}, \pi, -1\right)$ .

3) The vector field  $\nabla(\exp x + \ln(1 + |yz|))$  along the broken line  $\mathcal{K}$ , which goes from  $(0, 1, 1)$ , via  $(\pi, -3, 2)$  to  $(1, \sqrt{3}, -\sqrt{3})$ .

**A** Tangential line integrals of gradient fields.

**D** Use that

$$\int_{\mathcal{K}} \nabla F \cdot d\mathbf{x} = F(\text{end point}) - F(\text{initial point}),$$

is independent of the path of integration.

Since the absolute value occurs in **Example 4.7.3**, we shall here be very careful.

**I** 1) As  $\mathcal{K}$  is a closed curve (i.e. the initial point  $(1, 0, 0)$  is equal to the end point), it follows that

$$\int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} = 0.$$

2) Since  $F(x, y, z) = \cos(xyz)$ , and the initial point and the end point are given, we have

$$\int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} = \cos\left(\frac{1}{2} \cdot \pi \cdot (-1)\right) - \cos\left(\pi \cdot \frac{1}{2} \cdot 0\right) = -1.$$

3) First notice that

$$F(x, y, z) = \begin{cases} \exp x + \ln(1 + yz) & \text{for } yz > 0, \\ \exp x + \ln(1 - yz) & \text{for } yz < 0, \end{cases}$$

so we must be very careful, whenever the curve  $\mathcal{K}$  intersects one of the planes  $y = 0$  or  $z = 0$ . In case of the first curves this can occur, because the parametric description is

$$t(0, 1, 1) + (1 - t)(\pi, -3, 2) = ((1 - t)\pi, 4t - 3, 2 - t), \quad t \in [0, 1],$$

and the same is true for the second curve, because it has the parametric description

$$t(\pi, -3, 2) + (1 - t)(1, \sqrt{3}, -\sqrt{3}) = (1 + t(\pi - 1), \sqrt{3} - t(3 + \sqrt{3}), -\sqrt{3} + t(2 + \sqrt{3})),$$

for  $t \in [0, 1]$ .

The former curve intersects the plane  $y = 0$  for  $t = \frac{3}{4}$ , and the latter curve intersects both the plane  $y = 0$  and the plane  $z = 0$ . The point is, however, that in everyone of these intersection points the dubious term  $\ln(1 + |xy|) = 0$ , so they are of no importance. Hence we can conclude that

$$\begin{aligned} \int_{\mathcal{K}} \mathbf{V} \cdot d\mathbf{x} &= [\exp x + \ln(1 + |yz|)]_{(0,1,1)}^{(1,\sqrt{3},-\sqrt{3})} \\ &= e + \ln 4 - 1 - \ln 2 = e - 1 + \ln 2. \end{aligned}$$

REMARK. Always be very careful when either the absolute value or the square root occur. One should at least give a note on them.  $\diamond$

Please click the advert

SIMPLY CLEVER

ŠKODA



We will turn your CV into  
an opportunity of a lifetime



Do you like cars? Would you like to be a part of a successful brand?  
We will appreciate and reward both your enthusiasm and talent.  
Send us your CV. You will be surprised where it can take you.

Send us your CV on  
[www.employerforlife.com](http://www.employerforlife.com)



**Example 4.8** Given the vector field

$$\mathbf{V}(x, y) = \left( \frac{2y}{2x+y}, \frac{y}{2x+y} + \ln(2x+y) \right).$$

1. Sketch the domain of  $\mathbf{V}$ , and explain why  $\mathbf{V}$  is a gradient field.
2. Find every integral of  $\mathbf{V}$ .

Let  $\mathcal{K}$  be the curve given by

$$(x, y) = (2t^2, t), \quad 1 \leq t \leq 2.$$

3. Compute the value of the tangential line integral

$$\int_{\mathcal{K}} \mathbf{V} \cdot \mathbf{t} \, ds.$$

Let  $F$  be the integral of  $\mathbf{V}$ , for which  $F(1, 1) = 0$ .

4. Find an equation of the tangent at the en point  $(1, 1)$  of that level curve for  $F$ , which goes through the point  $(1, 1)$ .

**A** Gradient field, integrals, tangential line integral, level curve.

**D** Follow the guidelines.




Figure 66: The domain is the open half plane above the oblique line.

- I** 1) Clearly,  $\mathbf{V}(x, y)$  is defined in the domain where  $2x + y > 0$ , cf. the figure.

As

$$\frac{\partial V_1}{\partial y} = \frac{2}{2x+y} - \frac{2y}{(2x+y)^2},$$

and

$$\frac{\partial V_2}{\partial x} = -\frac{2y}{(2x+y)^2} + \frac{2}{2x+y} = \frac{\partial V_1}{\partial y},$$

it follows that  $V_1 dx + V_2 dy$  is a closed differential form. Since the domain is simply connected, the differential form is even exact, and  $\mathbf{V}$  is a gradient field.

2) Since

$$F_1(x, y) = \int \frac{2y}{2x+y} dx = y \int \frac{2 dx}{2x+y} = y \ln(2x+y), \quad 2x+y > 0,$$

where

$$\nabla F_1 = \left( \frac{2y}{2x+y}, \frac{y}{2x+y} + \ln(2x+y) \right) = \mathbf{V}(x, y),$$

all integrals are given by

$$F(x, y) = y \ln(2x+y) + C, \quad C \in \mathbb{R}.$$




Figure 67: The curve  $\mathcal{K}$ .

3) We get by the reduction theorem for tangential line integrals that

$$\begin{aligned} \int_{\mathcal{K}} \mathbf{V} \cdot \mathbf{t} ds &= \int_1^2 \mathbf{V}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt \\ &= \int_1^2 \left( \frac{2t}{4t^2+t}, \frac{t}{4t^2+t} + \ln(4t^2+t) \right) \cdot (4t, 1) dt \\ &= \int_1^2 \left\{ \frac{8t}{4t+1} + \frac{1}{4t+1} + \ln(4t^2+t) \right\} dt \\ &= \int_1^2 \left\{ 2 - \frac{1}{4t+1} + \ln t + \ln(4t+1) \right\} dt \\ &= 2 - \frac{1}{4} [\ln(4t+1)]_1^2 + [t \ln t - t]_1^2 + [t \ln(4t+1)]_1^2 - \int_1^2 \frac{4t}{4t+1} dt \\ &= 2 - \frac{1}{4} [\ln(4t+1)]_1^2 + 2 \ln 2 - 1 + 2 \ln 9 - \ln 5 - 1 + \frac{1}{4} [\ln(4t+1)]_1^2 \\ &= 2 \ln 2 + 4 \ln 3 - \ln 5 = \ln \frac{324}{5}. \end{aligned}$$

4) It follows from  $F(1, 1) = \ln 3 + C = 0$  that  $C = -\ln 3$ , so

$$F(x, y) = y \ln(2x+y) - \ln 3.$$



However, we shall not need the exact value of  $C = -\ln 3$  in the following.

The normal of the level curve is  $\nabla F = \mathbf{V}$ , hence

$$\mathbf{V}(1,1) = \left( \frac{2}{3}, \frac{1}{3} + \ln 3 \right),$$

and the direction of the tangent is e.g.

$$\mathbf{v} = \left( \frac{1}{3} + \ln 3, -\frac{2}{3} \right),$$

and we get a parametric description of the tangent,

$$(x(t), y(t)) = (1, 1) + t \left( \frac{1}{3} + \ln 3, -\frac{2}{3} \right), \quad t \in \mathbb{R}.$$

If we instead want an equation of the tangent, then one possibility is given by

$$0 = \mathbf{V} \cdot (x - 1, y - 1) = \frac{2}{3}x + \left( \frac{1}{3} + \ln 3 \right)y - 1 - \ln 3.$$

Please click the advert

With us you can  
shape the future.  
Every single day.

For more information go to:  
[www.eon-career.com](http://www.eon-career.com)

Your energy shapes the future.

**e.on**