

Integral Operators

Functional Analysis Examples c-5

Leif Mejlbro



Download free books at

bookboon.com

Leif Mejlbro

Integral Operators

Integral Operators

© 2009 Leif Mejlbro & Ventus Publishing ApS


ISBN 978-87-7681-529-5

Disclaimer: The texts of the advertisements are the sole responsibility of Ventus Publishing, no endorsement of them by the author is either stated or implied.

Contents

1.	Hilbert-Smith operators	5
2.	Other types of integral operators	47
	Index	66

Please click the advert



Masters in Management



Designed for high-achieving graduates across all disciplines, London Business School's Masters in Management provides specific and tangible foundations for a successful career in business.

This 12-month, full-time programme is a business qualification with impact. In 2010, our MiM employment rate was 95% within 3 months of graduation*; the majority of graduates choosing to work in consulting or financial services.

As well as a renowned qualification from a world-class business school, you also gain access to the School's network of more than 34,000 global alumni – a community that offers support and opportunities throughout your career.

For more information visit www.london.edu/mm, email mim@london.edu or give us a call on **+44 (0)20 7000 7573**.

* Figures taken from London Business School's Masters in Management 2010 employment report

1 Hilbert-Schmidt operators

Example 1.1 Let (e_k) denote an orthonormal basis in a Hilbert space H , and assume that the operator T has the matrix representation (t_{jk}) with respect to the basis (e_k) . Show that

$$\sum_{j=1}^{\infty} \sum_{k=1}^{\infty} |t_{jk}|^2 < \infty$$

implies that T is compact.

Let (f_k) denote another orthonormal basis in H , and let

$$s_{jk} = (Tf_j, f_k)$$

so that (s_{jk}) is the matrix representation of T with respect to the basis (f_k) .

Show that

$$\sum_{j=1}^{\infty} \sum_{k=1}^{\infty} |t_{jk}|^2 = \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} |s_{jk}|^2.$$

An operator satisfying

$$\sum_{j=1}^{\infty} \sum_{k=1}^{\infty} |t_{jk}|^2 < \infty$$

is called a general Hilbert-Schmidt operator.

Write $t_{jk} = (Te_j, e_j)$. It follows from VENTUS, HILBERT SPACES, ETC., EXAMPLE 2.7 that

$$Tx = T \left(\sum_{j=1}^{+\infty} x_j e_j \right) = \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} x_j t_{jk} e_k.$$

Define the sequence (T_n) of operators by

$$T_n x = T_n \left(\sum_{j=1}^{+\infty} x_j e_j \right) = \sum_{j=1}^{+\infty} \sum_{k=1}^n x_j t_{jk} e_k.$$

The range of T_n is finite dimensional, so T_n is compact. Then we conclude from

$$\|(T - T_n)x\|^2 = \left\| \sum_{j=1}^{+\infty} \sum_{n=1}^{+\infty} x_j t_{jk} e_k \right\|^2 = \sum_{k=n+1}^{+\infty} \left| \sum_{j=1}^{+\infty} x_j t_{jk} \right|^2,$$

where

$$\left| \sum_{j=1}^{+\infty} x_j t_{jk} \right|^2 \leq \left\{ \sum_{j=1}^{+\infty} |x_j|^2 \right\} \cdot \left\{ \sum_{j=1}^{+\infty} |t_{jk}|^2 \right\},$$

that

$$\|(T - T_n)x\|^2 \leq \left\{ \sum_{k=n+1}^{+\infty} \sum_{j=1}^{+\infty} |t_{jk}|^2 \right\} \cdot \|x\|^2.$$

It follows that

$$\|T - T_n\|^2 \leq \sum_{k=n+1}^{+\infty} \sum_{j=1}^{+\infty} |t_{jk}|^2.$$

Putting

$$a_k = \sum_{j=1}^{+\infty} |t_{jk}|^2 \geq 0,$$

it follows from the assumption that

$$\sum_{k=1}^{+\infty} a_k = \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |t_{jk}|^2 < +\infty.$$

Hence, to every $\varepsilon > 0$ there is an $n \in \mathbb{N}$, such that

$$\sum_{k=n+1}^{+\infty} a_k < \varepsilon^2,$$

Please click the advert



You're full of *energy*
and *ideas*. And that's
just what we are looking for.

Looking for a career where your ideas could really make a difference? UBS's Graduate Programme and internships are a chance for you to experience for yourself what it's like to be part of a global team that rewards your input and believes in succeeding together.

Wherever you are in your academic career, make your future a part of ours by visiting www.ubs.com/graduates.

www.ubs.com/graduates



© UBS 2010. All rights reserved.

from which

$$\|T - T_n\|^2 \leq \sum_{k=n+1}^{+\infty} \sum_{j=1}^{+\infty} |t_{jk}|^2 = \sum_{k=n+1}^{+\infty} a_k < \varepsilon^2,$$

thus $\|T - T_n\| < \varepsilon$, and we have proved that $T_n \rightarrow T$. Because all the T_n are compact, we conclude that T is also compact.

Given another orthonormal basis (f_k) of H , and let $s_{jk} = (Tf_j, f_k)$. Then an application of Parseval's equation gives that

$$\sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |(Te_k, f_j)|^2 = \sum_{k=1}^{+\infty} \|Te_k\|^2 = \sum_{k=1}^{+\infty} \sum_{j=1}^{+\infty} |(Te_k, e_j)|^2 = \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |t_{kj}|^2$$

and

$$\begin{aligned} \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |(Te_k, f_j)|^2 &= \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |(e_k, T^* f_j)|^2 = \sum_{j=1}^{+\infty} \|T^* f_j\|^2 = \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |(T^* f_j, f_k)|^2 \\ &= \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |(f_j, T f_k)|^2 = \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |(T f_j, f_k)|^2 = \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |s_{jk}|^2, \end{aligned}$$

hence,

$$\sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |t_{jk}|^2 = \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |t_{kj}|^2 = \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |s_{jk}|^2.$$

Example 1.2 For a general Hilbert-Schmidt operator we define the Hilbert-Schmidt norm $\|\cdot\|_{\text{HS}}$ by

$$\|T\|_{\text{HS}} = \left\{ \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |t_{jk}|^2 \right\}^{\frac{1}{2}}.$$

Show that this is a norm, and show that

$$\|T\| \leq \|T\|_{\text{HS}}$$

for a general Hilbert-Schmidt operator T .

Write $t_{jk} = (Te_j, e_k)$, and let

$$\|T\|_{\text{HS}} = \left\{ \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |t_{jk}|^2 \right\}^{\frac{1}{2}}.$$

Then $\|T\|_{\text{HS}} \geq 0$, and if $\|T\|_{\text{HS}} = 0$, then $t_{jk} = (Te_j, e_k) = 0$ for all $j, k \in \mathbb{N}$, thus

$$Te_j = \sum_{k=1}^{+\infty} (Te_j, e_k) e_k = \sum_{k=1}^{+\infty} t_{jk} e_k = 0 \quad \text{for every } j \in \mathbb{N}.$$

It follows that $T = 0$ as required.

We infer from $(\alpha T e_j, e_k) = \alpha (T e_j, e_k) = \alpha t_{jk}$ that

$$\|\alpha T\|_{\text{HS}} = \left\{ |\alpha|^2 \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |t_{jk}|^2 \right\}^{\frac{1}{2}} = |\alpha| \cdot \|T\|_{\text{HS}}.$$

Finally, if $\mathbf{S} = (s_{jk})$ and $\mathbf{T} = (t_{jk})$, then

$$\begin{aligned} \|S + T\|_{\text{HS}}^2 &= \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |s_{jk} + t_{jk}|^2 \leq \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} \left\{ |s_{jk}|^2 + 2 |s_{jk}| \cdot |t_{jk}| + |t_{jk}|^2 \right\} \\ &= \|S\|_{\text{HS}}^2 + \|T\|_{\text{HS}}^2 + 2 \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |s_{jk}| \cdot |t_{jk}| \\ &\leq \|S\|_{\text{HS}}^2 + \|T\|_{\text{HS}}^2 + 2 \left\{ \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |s_{jk}|^2 \right\}^{\frac{1}{2}} \cdot \left\{ \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |t_{jk}|^2 \right\}^{\frac{1}{2}} \\ &= \|S\|_{\text{HS}}^2 + \|T\|_{\text{HS}}^2 + 2 \|S\|_{\text{HS}} \cdot \|T\|_{\text{HS}} = \{\|S\|_{\text{HS}} + \|T\|_{\text{HS}}\}^2, \end{aligned}$$

and we have proved the triangle inequality,

$$\|S + T\|_{\text{HS}} \leq \|S\|_{\text{HS}} + \|T\|_{\text{HS}}.$$

We have proved that $\|\cdot\|_{\text{HS}}$ is a norm.

Please click the advert



Discover the truth at www.deloitte.ca/careers

Deloitte.

© Deloitte & Touche LLP and affiliated entities.

Finally,

$$\begin{aligned}
\|Tx\|^2 &= \left\| \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} x_j t_{jk} e_k \right\|^2 = \sum_{k=1}^{+\infty} \left| \sum_{j=1}^{+\infty} x_j t_{jk} \right|^2 \leq \sum_{k=1}^{+\infty} \sum_{j=1}^{+\infty} \sum_{\ell=1}^{+\infty} |x_j| \cdot |t_{jk}| \cdot |x_\ell| \cdot |t_{\ell k}| \\
&= \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} \sum_{\ell=1}^{+\infty} \{|x_j| \cdot |t_{\ell k}|\} \cdot \{|x_\ell| \cdot |t_{jk}|\} \\
&\leq \left\{ \sum_{j,k,\ell=1}^{+\infty} |x_j|^2 |t_{\ell j}|^2 \right\}^{\frac{1}{2}} \cdot \left\{ \sum_{j,k,\ell=1}^{+\infty} |x_\ell|^2 |t_{jk}|^2 \right\}^{\frac{1}{2}} = \|T\|_{\text{HS}}^2 \cdot \|x\|^2,
\end{aligned}$$

hence $\|Tx\| \leq \|T\|_{\text{HS}} \cdot \|x\|$ for every x , and we find that $\|T\| \leq \|T\|_{\text{HS}}$.

Example 1.3 Define for $f \in L^2(\mathbb{R})$, the operator K by

$$Kf(x) = \int_{-\infty}^{\infty} \frac{1}{2} \exp(-|x-t|) f(t) dt.$$

Show that $Kf \in L^2(\mathbb{R})$ and that K is linear and bounded, with norm ≤ 1 .

Show that the function $\frac{1}{2} \exp(-|x-t|)$ does not belong to $L^2(\mathbb{R}^2)$, so that K is not a Hilbert-Schmidt operator.

First we see that

$$\begin{aligned}
Kf(x) &= \int_{-\infty}^{+\infty} \frac{1}{2} \exp(-|x-t|) f(t) dt = \int_{-\infty}^x \frac{1}{2} e^{-x} e^t f(t) dt + \int_x^{+\infty} \frac{1}{2} e^x e^{-t} f(t) dt \\
&= \frac{1}{2} e^{-x} \int_{-\infty}^x e^t f(t) dt + \frac{1}{2} e^x \int_x^{+\infty} e^{-t} f(t) dt.
\end{aligned}$$

Then

$$\begin{aligned}
|Kf(x)|^2 &= \left\{ \int_{-\infty}^{+\infty} \frac{1}{2} \exp(-|x-t|) f(t) dt \right\}^2 \\
&\leq \int_{-\infty}^{+\infty} \frac{1}{2} \exp(-|x-t|) |f(t)| dt \cdot \int_{-\infty}^{+\infty} \frac{1}{2} \exp(-|x-u|) |f(u)| du \\
&= \frac{1}{4} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \exp(-|x-t|) \exp(-|x-u|) \cdot |f(t)| \cdot |f(u)| dt du \\
&= \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{1}{4} \exp(-|x-t| - |x-u|) \cdot |f(t)| \cdot |f(u)| dt du.
\end{aligned}$$

Hence

$$\int_{-\infty}^{+\infty} |Kf(x)|^2 dx \leq \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \left\{ \int_{-\infty}^{+\infty} \frac{1}{4} \exp(-|x-t| - |x-u|) dx \right\} |f(t)| \cdot |f(u)| dt du.$$

If $t \leq u$, then

$$|x-t| + |x-u| = \begin{cases} t-x+u-x = t+u-2x, & \text{for } x \leq t, \\ x-t+u-x = u-t, & \text{for } t \leq x \leq u, \\ x-t+x-u = 2x-t-u, & \text{for } x \geq u. \end{cases}$$

This gives the inspiration to the following rearrangement

$$\int_{-\infty}^{+\infty} |Kf(x)|^2 dx \leq 2 \int_{-\infty}^{+\infty} \left(\int_t^{+\infty} \left\{ \int_{-\infty}^{+\infty} \frac{1}{4} \exp(-|x-t| - |x-u|) dx \right\} |f(u)| du \right) |f(t)| dt,$$

where

$$\begin{aligned} \int_{-\infty}^{+\infty} e^{-|x-t|-|x-u|} dx &= \int_{-\infty}^t e^{2x-t-u} dx + \int_t^{+\infty} e^{-u+t} dx + \int_u^{+\infty} e^{-2x+t+u} dx \\ &= \left[\frac{1}{2} e^{2x-t-u} \right]_{x=-\infty}^t + (u-t)e^{-u+t} + \left[-\frac{1}{2} e^{-2x+t+u} \right]_{x=u}^{+\infty} \\ &= \frac{1}{2} e^{t-u} + (u-t)e^{t-u} + \frac{1}{2} e^{t-u} = (u-t+1)e^{t-u}, \end{aligned}$$

and where we have assumed that $t \leq u$.

By insertion,

$$\int_{-\infty}^{+\infty} |Kf(x)|^2 dx \leq \frac{1}{2} \int_{-\infty}^{+\infty} \left\{ \int_t^{+\infty} (u-t+1)e^{t-u} |f(u)| du \right\} |f(t)| dt.$$

Then we change variables $y = u - t$ and $z = t + u$, thus

$$t = \frac{y+z}{2} \quad \text{og} \quad u = \frac{y-z}{2},$$

where $y \in [0, +\infty[$ and $z \in \mathbb{R}$. We get

$$\begin{aligned} \int_{-\infty}^{+\infty} |Kf(x)|^2 dx &\leq \frac{1}{4} \int_{-\infty}^{+\infty} \int_0^{+\infty} (y+1)e^{-y} \left| f\left(\frac{y-z}{2}\right) \right| \cdot \left| f\left(\frac{y+z}{2}\right) \right| dy dz \\ &= \frac{1}{4} \int_0^{+\infty} \left\{ \int_{-\infty}^{+\infty} \left| f\left(\frac{y-z}{2}\right) \right| \cdot \left| f\left(\frac{y+z}{2}\right) \right| dz \right\} (y+1)e^{-y} dy. \end{aligned}$$

Then for every fixed y it follows by the Cauchy-Schwarz inequality,

$$\begin{aligned} &\int_{-\infty}^{+\infty} \left| f\left(\frac{y-z}{2}\right) \right| \cdot \left| f\left(\frac{y+z}{2}\right) \right| dz \\ &\leq \left\{ \int_{-\infty}^{+\infty} \left| f\left(\frac{y-z}{2}\right) \right|^2 dz \right\}^{\frac{1}{2}} \cdot \left\{ \int_{-\infty}^{+\infty} \left| f\left(\frac{y+z}{2}\right) \right|^2 dz \right\}^{\frac{1}{2}} \\ &= \left\{ 2 \int_{-\infty}^{+\infty} \left| f\left(\frac{y-z}{2}\right) \right|^2 d\left(\frac{y-z}{2}\right) \right\}^{\frac{1}{2}} \cdot \left\{ 2 \int_{-\infty}^{+\infty} \left| f\left(\frac{y-z}{2}\right) \right|^2 d\left(\frac{y+z}{2}\right) \right\}^{\frac{1}{2}} \\ &= 2\|f\|_2 \cdot \|f\|_2 = 2\|f\|_2^2, \end{aligned}$$

and we get by insertion the estimate

$$\begin{aligned} \int_{-\infty}^{+\infty} |Kf(x)|^2 dx &\leq \frac{1}{2} \int_0^{+\infty} (y+1)e^{-y} dy \cdot \|f\|_2^2 \\ &= \frac{1}{2} \left[-e^{-y}(y+1) + \int e^{-y} dy \right]_0^{+\infty} \cdot \|f\|_2^2 \\ &= \frac{1}{2} [-e^{-y}(y+2)]_0^{+\infty} \cdot \|f\|_2^2 = \|f\|_2^2, \end{aligned}$$

so we have proved that $Kf \in L^2(\mathbb{R})$ and that

$$\|Kf\|_2 \leq \|f\|_2 \quad \text{for every } f \in L^2(\mathbb{R}),$$

hence $\|K\| \leq 1$.

On the other hand, the kernel $\frac{1}{2}e^{-|x-t|}$ does not belong to $L^2(\mathbb{R})$, because we get by a formal computation that

$$\begin{aligned} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{1}{4} e^{-2|x-t|} dx dt &= \frac{1}{4} \int_{-\infty}^{+\infty} \left\{ 2 \int_t^{+\infty} e^{-2(x-t)} dx \right\} dt \\ &= \frac{1}{4} \int_{-\infty}^{+\infty} \left\{ \int_0^{+\infty} e^{-x} dx \right\} dt = \frac{1}{4} \int_{-\infty}^{+\infty} 1 dt = +\infty. \end{aligned}$$

Example 1.4 Let K denote the Hilbert-Schmidt operator with kernel

$$k(x, y) = \sin(x) \cos(t), \quad 0 \leq x, t \leq 2\pi.$$

Show that the only eigenvalue for K is 0.
Find an orthonormal basis for $\ker(K)$.

First notice that

$$Kf(x) = \int_0^{2\pi} k(x, t) f(t) dt = \sin(x) \cdot \int_0^{2\pi} \cos(t) \cdot f(t) dt,$$

hence $Kf(x) = a(f) \cdot \sin(x)$, where

$$a(f) = \int_0^{2\pi} \cos(t) \cdot f(t) dt \in \mathbb{C}.$$

If $\lambda \in \sigma_p(K)$, then the corresponding eigenfunction must be $f(x) = \sin(x)$. Then by insertion,

$$(K \sin)(x) = \sin(x) \int_0^{2\pi} \cos(t) \cdot \sin(t) dt = 0,$$

proving that $\lambda = 0$ is the only eigenvalue.

Now,

$$\frac{1}{\sqrt{2\pi}}, \frac{1}{\sqrt{\pi}} \cos(x), \frac{1}{\sqrt{\pi}} \sin(x), \dots, \frac{1}{\sqrt{\pi}} \cos(nx), \frac{1}{\sqrt{\pi}} \sin(nx), \dots,$$

is an orthonormal basis for $L^2([0, 2\pi])$, so $\ker(K)$ is spanned by all these with the exception of $\frac{1}{\sqrt{\pi}} \cos(x)$, in which case

$$\begin{aligned} K \left(\frac{1}{\sqrt{\pi}} \cos \right) (x) &= \sqrt{\pi} \int_0^{2\pi} \frac{1}{\sqrt{\pi}} \cos(t) \cdot \frac{1}{\sqrt{\pi}} \cos(t) dt \cdot \sin(x) \\ &= \sqrt{\pi} \cdot \sin(x) = \pi \cdot \frac{1}{\sqrt{\pi}} \sin(x), \end{aligned}$$

and we get in particular, $K^2 \equiv 0$.

Note that

$$\begin{aligned} k_2(x, t) &= \int_0^{2\pi} k(x, s)k(s, t) ds = \int_0^{2\pi} \sin(x) \cdot \cos(s) \cdot \sin(s) \cdot \cos(t) ds \\ &= \sin(x) \cdot \cos(t) \cdot \int_0^{2\pi} \sin(s) \cdot \cos(s) ds = 0, \end{aligned}$$

which agrees with $K^2 \equiv 0$.

Please click the advert

It's only an opportunity if you act on it

IKEA.SE/STUDENT

© Inter IKEA Systems B.V. 2009

Example 1.5 Let K denote the Hilbert-Schmidt operator with continuous kernel k on $L^2(I)$, where I is a closed and bounded interval. Show that all the iterated kernels K_n are continuous on I^2 and show that

$$\|k_n\|_2 \leq \|k\|_2^n.$$

Show that if $|\lambda| \|k\|_2 < 1$, then the series

$$\sum_{n=1}^{\infty} \lambda^n k_n$$

is convergent in $L^2(I)$.

Write $I = [a, b]$. It is well-known that

$$k_n(x, t) = \int_a^b f(x, s) k_{n-1}(s, t) ds.$$

The first claim is proved by induction. Assume that both $k(x, s)$ and $k_{n-1}(s, t)$ are continuous. By subtracting something and then adding it again we get

$$\begin{aligned} k_n(x, t) - k_n(x_0, t_0) &= \int_a^b \{k(x, s)k_{n-1}(s, t) - k(x_0, s)k_{n-1}(s, t)\} ds \\ &\quad + \int_a^b \{k(x_0, s)k_{n-1}(s, t) - k(x_0, s)k_{n-1}(s, t_0)\} ds \\ &= \int_a^b \{k(x, s) - k(x_0, s)\} k_{n-1}(s, t) ds \\ &\quad + \int_a^b k(x_0, s) \cdot \{k_{n-1}(s, t) - k_{n-1}(s, t_0)\} ds. \end{aligned}$$

To every $\varepsilon > 0$ there is a $\delta > 0$, such that

$$|k(x, s) - k(x_0, s)| < \varepsilon \quad \text{for } |x - x_0| < \delta \text{ and all } s \in [a, b],$$

and

$$|k_{n-1}(s, t) - k_{n-1}(s, t_0)| < \varepsilon \quad \text{for } |t - t_0| < \delta \text{ and all } s \in [a, b].$$

If therefore $|x - x_0| < \delta$ and $|t - t_0| < \delta$, then we get the following estimate,

$$\begin{aligned} |k_n(x, t) - k_n(x_0, t_0)| &\leq \int_a^b \varepsilon \cdot \|k_{n-1}\|_{\infty} dx + \int_a^b \|k\|_{\infty} \cdot \varepsilon ds \\ &= (b - a) \{\|k\|_{\infty} + \|k_{n-1}\|_{\infty}\} \varepsilon, \end{aligned}$$

and we conclude that $k_n(x, t)$ is continuous, and the claim follows by induction.

Furthermore,

$$\begin{aligned}
 \|k_n\|_2^2 &= \int_a^b \int_a^b |k_n(x, t)|^2 dx dt \\
 &= \int_a^b \int_a^b \left| \int_a^b k(x, s) k_{n-1}(s, t) ds \right| \cdot \left| \int_a^b k(x, r) k_{n-1}(r, t) dr \right| dx dt \\
 &\leq \int_a^b \int_a^b \int_a^b \int_a^b |k(x, s)| \cdot |k_{n-1}(s, t)| \cdot |k(x, r)| \cdot |k_{n-1}(r, t)| ds dr dx dt \\
 &\leq \frac{1}{2} \int_a^b \int_a^b \int_a^b \int_a^b \{ |k(x, s)|^2 |k_{n-1}(r, t)|^2 + |k_{n-1}(s, t)|^2 |k(x, r)|^2 \} ds dr dx dt \\
 &= \frac{1}{2} \{ \|k\|_2^2 \|k_{n-1}\|_2^2 + \|k_{n-1}\|_2^2 \|k\|_2^2 \} = \|k\|_2^2 \|k_{n-1}\|_2^2,
 \end{aligned}$$

and we have proved that

$$\|k_n\|_2 \leq \|k\|_2 \|k_{n-1}\|_2.$$

Hence we get for $n = 2$ that $\|k_2\|_2 \leq \|k\|_2^2$.

Assume that $\|k_{n-1}\|_2 \leq \|k\|_2^{n-1}$. Then

$$\|k_n\|_2 \leq \|k\|_2 \|k_{n-1}\|_2 \leq \|k\|_2 \cdot \|k\|_2^{n-1} = \|k\|_2^n,$$

and the claim follows by induction.

Please click the advert

YOUR CHANCE TO CHANGE THE WORLD

Here at Ericsson we have a deep rooted belief that the innovations we make on a daily basis can have a profound effect on making the world a better place for people, business and society. Join us.

In Germany we are especially looking for graduates as Integration Engineers for

- Radio Access and IP Networks
- IMS and IPTV

We are looking forward to getting your application!
To apply and for all current job openings please visit our web page: www.ericsson.com/careers

ericsson.
com



The remaining claim is now trivial, because

$$\left\| \sum_{n=1}^{+\infty} \lambda^n k_n(x, t) \right\|_2 \leq \sum_{n=1}^{+\infty} |\lambda|^n \|k_n\|_2 \leq \sum_{n=1}^{+\infty} |\lambda|^n \|k\|_2^n = \sum_{n=1}^{+\infty} \{|\lambda| \cdot \|k\|_2\}^n = \frac{1}{1 - |\lambda| \cdot \|k\|_2},$$

where we have used that the geometric series is convergent for $|\lambda| \cdot \|k\|_2 < 1$.

Example 1.6 Let K and L denote the Hilbert-Schmidt operators with continuous kernels k and ℓ on $L^2(I)$, where I is a closed and bounded interval. We define the trace of K , $\text{tr}(K)$ by

$$\text{tr}(K) = \int_I k(x, x) dx,$$

and similarly for L .

Show that

$$|\text{tr}(KL)| \leq \|K\|_{\text{HS}} \|L\|_{\text{HS}},$$

and

$$|\text{tr}(K^n)| \leq \|K\|_{\text{HS}}^n, \quad n \geq 2.$$

Moreover, if (K_n) , (L_n) denote sequences of Hilbert-Schmidt operators like above, where

$$\|K_n - K\|_{\text{HS}} \rightarrow 0 \quad \text{and} \quad \|L_n - L\|_{\text{HS}} \rightarrow 0,$$

then

$$\text{tr}(K_n L_n) \rightarrow \text{tr}(KL).$$

Remark 1.1 We first show that the claim is not true, if we replace the Hilbert-Schmidt norm $\|\cdot\|_{\text{HS}}$ by the operator norm.

Let

$$k(x, t) = \ell(x, t) = x + t$$

be the kernel of self adjoint Hilbert-Schmidt operators K and L on $L^2([0, 1])$. It follows from Example 1.7 below that $\frac{1}{2} \pm \frac{1}{\sqrt{3}}$ are the two eigenvalues different from zero of both K and L , and the norm of K (and L) is given by the absolute value of the numerically largest eigenvalue,

$$\|K\| = \|L\| = \frac{1}{2} + \frac{1}{\sqrt{3}}.$$

Furthermore,¹

$$\begin{aligned} \|k\|_2^2 &= \|\ell\|_2^2 = \int_0^1 \int_0^1 (x+t)^2 dx dt = \int_0^1 \int_0^1 (x^2 + 2xt + t^2) dx dt = \int_0^1 \left[\frac{x^3}{3} + x^2 t + x t^2 \right]_{x=0}^1 dt \\ &= \int_0^1 \left\{ \frac{1}{3} + t + t^2 \right\} dt = \frac{1}{3} + \frac{1}{2} + \frac{1}{3} = \frac{7}{6}. \end{aligned}$$

Finally,

$$\operatorname{tr}(KL) = \int_0^1 \left\{ \int_0^1 (x+s)(s+x) ds \right\} dx = \int_0^1 \left\{ \int_0^1 (x+s)^2 ds \right\} dx = \|k\|_2^2 = \frac{7}{6}.$$

Thus, in this example,

$$\operatorname{tr}(KT) = \frac{7}{6} = \|k\|_2^k > \|K\|^2 = \|K\| \cdot \|L\| = \left\{ \frac{1}{2} + \frac{1}{\sqrt{3}} \right\}^2 = \frac{1}{4} + \frac{1}{3} + \frac{\sqrt{3}}{3},$$

which either can be shown numerically, or of course must follow from the theory, because we always have that $\|K\| \leq \|k\|_2$. Here we cannot have equality, if $\sigma_p(K)$ contains at least two different points $\neq 0$. \diamond

Then we turn to the example itself.

Write $I = [a, b]$, and let

$$Ku(x) = \int_a^b k(x, t)u(t) dt \quad \text{and} \quad Lu(x) = \int_a^b \ell(x, t)u(t) dt$$

for $u \in L^2([a, b])$. Then

$$\begin{aligned} ((KL)u)(x) &= K(Lu)(x) = \int_a^b k(x, t) Lu(t) dt = \int_a^b k(x, t) \left\{ \int_a^b \ell(t, s)u(s) ds \right\} dt \\ &= \int_a^b \left\{ \int_a^b k(x, t)\ell(t, s) dt \right\} u(s) ds, \end{aligned}$$

and it follows that the composition KL has the kernel

$$m(x, t) = \int_a^b k(x, s)\ell(s, t) ds.$$

Then

$$\begin{aligned} |\operatorname{tr}(KL)| &= \left| \int_a^b m(x, x) dx \right| = \left| \int_a^b \left\{ \int_a^b k(x, t)\ell(t, x) dt \right\} dx \right| \\ &\leq \int_a^b \left\{ \int_a^b |k(x, t)|^2 dt \right\}^{\frac{1}{2}} \cdot \left\{ \int_a^b |\ell(t, x)|^2 dt \right\}^{\frac{1}{2}} dx. \end{aligned}$$

Putting

$$k_1(x) = \left\{ \int_a^b |k(x, t)|^2 dt \right\}^{\frac{1}{2}} \quad \text{og} \quad \ell_1(x) = \left\{ \int_a^b |\ell(t, x)|^2 dt \right\}^{\frac{1}{2}},$$

we get $k_1, \ell_1 \in L^2([a, b])$, and it follows from the Cauchy-Schwarz inequality that

$$\begin{aligned} |\operatorname{tr}(KL)| &\leq \int_a^b k_1(x)\ell_1(x) dx \leq \{k_1(x)^2 dx\}^{\frac{1}{2}} \left\{ \int_a^b \ell_1(x)^2 dx \right\}^{\frac{1}{2}} \\ &= \left\{ \int_a^b \left(\int_a^b |k(x, t)|^2 dt \right) dx \right\}^{\frac{1}{2}} \left\{ \int_a^b \left(\int_a^b |\ell(t, x)|^2 dt \right) dx \right\}^{\frac{1}{2}} \\ &= \|k\|_2 \cdot \|\ell\|_2 = \|K\|_{\text{HS}} \cdot \|L\|_{\text{HS}}, \end{aligned}$$

and the first claim is proved.

We note that since KL has the kernel

$$m(x, t) = \int_a^b k(x, s)\ell(s, t) ds,$$

we have

$$\begin{aligned} \|KL\|_{\text{HS}}^2 &\leq \int_a^b \int_a^b |m(x, t)|^2 dx dt = \int_a^b \left\{ \int_a^b \left| \int_a^b k(x, s)\ell(s, t) ds \right|^2 dx \right\} dt \\ &\leq \int_a^b \left(\int_a^b \left\{ \left(\int_a^b |k(x, s)|^2 ds \right)^{\frac{1}{2}} \left(\int_a^b |\ell(s, t)|^2 ds \right)^{\frac{1}{2}} \right\}^2 dx \right) dt \\ &= \int_a^b \left(\int_a^b \left\{ \left(\int_a^b |k(x, s)|^2 ds \right) \cdot \left(\int_a^b |\ell(s, t)|^2 ds \right) \right\} dx \right) dt \\ &= \int_a^b \int_a^b |k(x, s)|^2 ds dx \cdot \int_a^b \int_a^b |\ell(s, t)|^2 ds dt = \|k\|_2^2 \cdot \|\ell\|_2^2 = \|K\|_{\text{HS}}^2 \cdot \|L\|_{\text{HS}}^2. \end{aligned}$$

This proves that we always have

$$(1) \quad \|KL\|_{\text{HS}} \leq \|K\|_{\text{HS}} \cdot \|L\|_{\text{HS}}.$$

Recall for $n = 1$ that

$$\text{tr}(K) = \int_a^b k(x, x) dx.$$

Choosing $k(x, x) = 1$ and $k(x, t)$ continuous, such that $\|k\|_2 < \varepsilon$, we get

$$\text{tr}(K) = b - a \quad \text{and} \quad \|K\|_{\text{HS}}^2 < \varepsilon,$$

which shows that the formula is not true for $n = 1$.

On the other hand, if $n \geq 2$, then it follows from the first question and (1) that

$$|\text{tr}(K^n)| = |\text{tr}(K K^{n-1})| \leq \|K\|_{\text{HS}} \|K^{n-1}\|_{\text{HS}} \leq \|K\|_{\text{HS}} \|K\|_{\text{HS}}^{n-1} = \|K\|_{\text{HS}}^n.$$

Finally, we note that for any scalar λ and any Hilbert-Schmidt operators,

$$\text{tr}(K + \lambda L) = \int_a^b \{k(x, x) + \lambda \ell(x, x)\} dx = \text{tr}(K) + \lambda \text{tr}(L),$$

proving that the *trace* is linear on the vector space of all Hilbert-Schmidt operators. Then we get

$$\begin{aligned} \text{tr}(KL) - \text{tr}(K_n L_n) &= \text{tr}(KL - K_n L_n) = \text{tr}(KL - K L_n + K L_n - K_n L_n) \\ &= \text{tr}(K(L - L_n)) + \text{tr}((K - K_n)L_n) \\ &= \text{tr}(K(L - L_n)) + \text{tr}((K - K_n)(L_n - L)) + \text{tr}((K - K_n)L), \end{aligned}$$

and it follows from the assumptions and the first part of the example that

$$\begin{aligned} |\text{tr}(KL) - \text{tr}(K_n L_n)| \\ \leq \|K\|_{\text{HS}} \|L - L_n\|_{\text{HS}} + \|K - K_n\|_{\text{HS}} \|L - L_n\|_{\text{HS}} + \|K - K_n\|_{\text{HS}} \|L\|_{\text{HS}} \rightarrow 0 \quad \text{for } n \rightarrow +\infty. \end{aligned}$$

Example 1.7 Let K denote the Hilbert-Schmidt operator on $L^2([0, 1])$ with kernel

$$k(x, t) = x + t.$$

Find all eigenvalues and eigenfunctions for K .
Solve the equation

$$Ku = \mu u + f, \quad f \in L^2([0, 1]),$$

when μ is not in the spectrum for K .

It follows from

$$(2) \quad Kf(x) = x \int_0^1 f(t) dt + \int_0^1 t \cdot f(t) dt,$$

that every eigenfunction corresponding to an eigenvalue $\lambda \neq 0$ must have the form $f(x) = ax + b$. By insertion into (2) we get

$$Kf(x) = x \int_0^1 (at + b) dt + \int_0^1 (at^2 + bt) dt = \left\{ \frac{a}{2} + b \right\} x + \left\{ \frac{a}{3} + \frac{b}{2} \right\}.$$

Please click the advert

SIMPLY CLEVER



We will turn your CV into an opportunity of a lifetime



Do you like cars? Would you like to be a part of a successful brand?
We will appreciate and reward both your enthusiasm and talent.
Send us your CV. You will be surprised where it can take you.

Send us your CV on
www.employerforlife.com



This expression is equal to $\lambda(ax + b)$, if and only if (a, b) and $\left(\frac{a}{2} + b, \frac{a}{3} + \frac{b}{2}\right)$ are proportion, thus if and only if

$$0 = \begin{vmatrix} \frac{a}{2} + b & \frac{a}{3} + \frac{b}{2} \\ a & b \end{vmatrix} = \frac{ab}{2} + b^2 - \frac{a^3}{3} - \frac{ab}{2} = b^2 - \frac{a^2}{3},$$

hence if and only if $b = \pm \frac{1}{\sqrt{3}} a$. Since

$$\lambda a = \frac{a}{2} + b = \left\{ \frac{1}{2} \pm \frac{1}{\sqrt{3}} \right\} a,$$

the corresponding eigenvalues are $\lambda = \frac{1}{2} \pm \frac{1}{\sqrt{3}}$.

For $\lambda_1 = \frac{1}{2} + \frac{1}{\sqrt{3}}$ we get the eigenfunction $f_1(x) = x + \frac{1}{\sqrt{3}}$.

For $\lambda_2 = \frac{1}{2} - \frac{1}{\sqrt{3}}$ we get the eigenfunction $f_2(x) = x - \frac{1}{\sqrt{3}}$.

Finally, K is trivially self adjoint, thus $\lambda = 0$ is an eigenvalue for every function

$$f \in \left\{ \text{span} \left(x + \frac{1}{\sqrt{3}}, x - \frac{1}{\sqrt{3}} \right) \right\}^\perp = \{ \text{span}(1, x) \}^\perp,$$

hence for every function $f \in L^2([0, 1])$, for which

$$\int_0^1 f(t) dt = 0 \quad \text{og} \quad \int_0^1 t f(t) dt = 0.$$

Now, $k(x, t) = \overline{k(t, x)}$, so K is self adjoint. Therefore, if we put

$$\varphi_1(x) = \frac{f_1}{\|f_1\|_2} \quad \text{and} \quad \varphi_2 = \frac{f_2}{\|f_2\|_2},$$

then the operator K is described by

$$(3) \quad Ku = \lambda_1 (u, \varphi_1) \varphi_1 + \lambda_2 (u, \varphi_2) \varphi_2.$$

If $(f, \varphi_1) = (f, \varphi_2) = 0$, then it follows by a simple check that the solution of the equation

$$Ku = \mu u + f, \quad \text{hvor } \mu \notin \left\{ 0, \frac{1}{2} + \frac{1}{\sqrt{3}}, \frac{1}{2} - \frac{1}{\sqrt{3}} \right\},$$

is given by $u = -\frac{1}{\mu} f$.

Then assume that $f = a \varphi_1 + b \varphi_2$. The equation $Ku = \mu u + f$ can now be written in the form

$$\lambda_1 (u, \varphi_1) \varphi_1 + \lambda_2 (u, \varphi_2) \varphi_2 = \mu \sum_{n=1}^{\infty} (u, \varphi_n) \varphi_n + a \varphi_1 + b \varphi_2,$$

which implies that

$$u = c_1 \varphi_1 + c_2 \varphi_2,$$

where

$$c_1 = (u, \varphi_1) = \frac{a}{\lambda_1 - \mu} = \frac{1}{\lambda_1 - \mu} (f, \varphi_1),$$

and

$$c_2 = (u, \varphi_2) = \frac{b}{\lambda_2 - \mu} = \frac{1}{\lambda_2 - \mu} (f, \varphi_2).$$

The equation being linear, it follows in general from the rewriting

$$Ku - \mu u = f = (f, \varphi_1) \varphi_1 + (f, \varphi_2) \varphi_2 + \{f - (f, \varphi_1) \varphi_1 - (f, \varphi_2) \varphi_2\},$$

that

$$\begin{aligned} u &= \frac{1}{\lambda_1 - \mu} (f, \varphi_1) \varphi_1 + \frac{1}{\lambda_2 - \mu} (f, \varphi_2) \varphi_2 - \frac{1}{\mu} f + \frac{1}{\mu} (f, \varphi_1) \varphi_1 + \frac{1}{\mu} (f, \varphi_2) \varphi_2 \\ &= \frac{\lambda_1}{\mu(\lambda_1 - \mu)} (f, \varphi_1) \varphi_1 + \frac{\lambda_2}{\mu(\lambda_2 - \mu)} (f, \varphi_2) \varphi_2 - \frac{1}{\mu} f = A \varphi_1 + B \varphi_2 - \frac{1}{\mu} f, \end{aligned}$$

which *in principle* can be written explicitly by means of the functions $f_i(x)$, $i = 1, 2$. We shall, however, not waste our time on that, because the result will look extremely nasty.

Example 1.8 Let K denote the Hilbert-Schmidt operator on $L^2\left(\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]\right)$ with kernel

$$k(x, t) = \cos(x - t).$$

Find all eigenvalues and eigenfunctions for K .

Solve the equation

$$Ku = \mu u + f, \quad f \in L^2\left(\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]\right),$$

when μ is not in the spectrum for K .

Obviously, K is self adjoint.

It follows in general from

$$\cos(x - t) = \cos(x) \cdot \cos(t) + \sin(x) \cdot \sin(t),$$

that

$$(4) \quad Kf(x) = \cos(x) \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} f(t) \cos(t) dt + \sin(x) \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} f(t) \sin(t) dt.$$

Then any eigenfunction corresponding to some eigenvalue $\lambda \neq 0$ must be of the structure

$$f(x) = a \cdot \cos(x) + b \cdot \sin(x).$$

By insertion into (4),

$$\begin{aligned} Kf(x) &= \cos(x) \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \{a \cdot \cos^2 t + b \cdot \sin t \cos t\} dt + \sin(x) \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \{a \cdot \sin t \cos t + b \cdot \sin^2 t\} dt \\ &= \left\{ \frac{a\pi}{2} + 0 \right\} \cos(x) + \left\{ 0 + \frac{b\pi}{2} \right\} \sin(x) = \frac{\pi}{2} \{a \cos(x) + b \sin(x)\} = \frac{\pi}{2} f(x), \end{aligned}$$

hence $f(x) = a \cdot \cos(x) + b \cdot \sin(x)$ is for every pair $(a, b) \neq (0, 0)$ an eigenfunction corresponding to the eigenvalue $\lambda = \frac{\pi}{2}$.

For $\lambda = 0$ we get the eigenspace $\{\cos(x), \sin(x)\}^\perp$ in $L^2\left(\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]\right)$.

ALTERNATIVELY, we see that

$$\cos(x-t) = \frac{1}{2} e^{ix} e^{-it} + \frac{1}{2} e^{-ix} e^{it}.$$

We get from

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} |e^{\pm ix}|^2 dx = \pi,$$

the normed functions

$$\varphi_1(x) = \frac{1}{\sqrt{\pi}} e^{ix} \quad \text{and} \quad \varphi_{-1} = \frac{1}{\sqrt{\pi}} e^{-ix},$$

where

$$(\varphi_1, \varphi_{-1}) = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \varphi_1(x) \overline{\varphi_{-1}(x)} dx = \frac{1}{\pi} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{2ix} dx = \frac{1}{2i\pi} \{e^{i\pi} - e^{-i\pi}\} = 0,$$

hence

$$k(x, t) = \cos(x-t) = \frac{\pi}{2} \varphi_1(x) \overline{\varphi_1(t)} + \frac{\pi}{2} \varphi_{-1}(x) \overline{\varphi_{-1}(t)}.$$

We obtain directly that $\lambda = \frac{\pi}{2}$ is the only eigenvalue $\neq 0$, thus $\|K\| = \frac{\pi}{2}$, and the eigenfunctions are φ_1 and φ_{-1} .

Remark 1.2 A basis for $L^2\left(\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]\right)$ is e.g.

$$\frac{1}{\sqrt{2\pi}}, \frac{1}{\sqrt{\pi}} \cos 2x, \frac{1}{\sqrt{\pi}} \sin 2x, \frac{1}{\sqrt{\pi}} \cos 4x, \frac{1}{\sqrt{\pi}} \sin 4x, \dots,$$

from which it follows that $\{\cos(x), \sin(x)\}^\perp$ may be difficult to describe. \diamond

It follows from $\overline{k(t, x)} = k(x, t)$ that K is self adjoint, which also was noted previously. We may therefore apply the standard method where we expand after the eigenfunctions.

First choose f , such that

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} f(t) \cos t \, dt = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} f(t) \sin t \, dt = 0.$$

Then $Kf = 0$, and we conclude that $u = -\frac{1}{\mu} f$ is the only solution.

We get in the general case that

$$\begin{aligned} u &= \sum_{n=1}^{+\infty} (u, \varphi_n) \varphi_n = \frac{1}{\frac{\pi}{2} - \mu} \{(f, \varphi_1) \varphi_1 + (f, \varphi_2) \varphi_2\} - \frac{1}{\mu} f + \frac{1}{\mu} (f, \varphi_1) \varphi_1 + \frac{1}{\pi} (f, \varphi_2) \varphi_2 \\ &= \frac{\frac{\pi}{2}}{\mu (\frac{\pi}{2} - \mu)} \{(f, \varphi_1) \varphi_1 + (f, \varphi_2) \varphi_2\} - \frac{1}{\mu} f. \end{aligned}$$

Please click the advert

With us you can
shape the future.
Every single day.

For more information go to:
www.eon-career.com

Your energy shapes the future.

e.on

Now,

$$\varphi_i = \frac{f_i}{\|f_i\|_2}, \quad i = 1, 2,$$

where $f_1(x) = \cos x$ and $f_2(x) = \sin x$, and $\|f_1\|_2^2 = \|f_2\|_2^2 = \frac{\pi}{2}$, hence

$$\begin{aligned} u &= \frac{\frac{\pi}{2}}{\mu(\frac{\pi}{2} - \mu)} \cdot \frac{1}{\frac{\pi}{2}} \{(f, \cos t) \cos(x) + (f, \sin t) \sin(x)\} - \frac{1}{\mu} f \\ &= \frac{1}{\mu(\frac{\pi}{2} - \mu)} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} f(t) \cos t \, dt \cdot \cos(x) + \frac{1}{\mu(\frac{\pi}{2} - \mu)} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} f(t) \sin t \, dt \cdot \sin(x) - \frac{1}{\mu} f(x). \end{aligned}$$

Notice that this expression can be written as

$$u = \frac{1}{\mu(\frac{\pi}{2} - \mu)} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos(x - t) f(t) \, dt - \frac{1}{\mu} f(x) = \frac{1}{\mu(\frac{\pi}{2} - \mu)} Kf - \frac{1}{\mu} f.$$

We have assumed that

$$\mu \notin \sigma(K) = \sigma_p(K) = \left\{0, \frac{\pi}{2}\right\}.$$

Example 1.9 Let K denote the Hilbert-Schmidt operator on $L^2([-\pi, \pi])$ with kernel

$$k(x, t) = \{\cos(x) + \cos(t)\}^2.$$

Find all eigenvalues and eigenfunctions for K , and find an orthonormal basis for $\ker(K)$.

By a simple computation,

$$\begin{aligned} k(x, t) &= (\cos x + \cos t)^2 = \cos^2 x + 2 \cos x \cos t + \cos^2 t \\ &= \frac{1}{2} \cos 2x + 2 \cos x \cos t + \frac{1}{2} \cos 2t + \frac{1}{2} \\ &= \frac{1}{2} \cos 2x + 2 \cos x \cos t + \left\{1 + \frac{1}{2} \cos 2t\right\} \cdot 1. \end{aligned}$$

Hence

$$\begin{aligned} (5) \quad Kf(x) &= \cos 2x \int_{-\pi}^{\pi} \frac{1}{2} f(t) \, dt + \cos x \int_{-\pi}^{\pi} 2 f(t) \cos t \, dt \\ &\quad + \int_{-\pi}^{\pi} f(t) \, dt + \int_{-\pi}^{\pi} \frac{1}{2} f(t) \cos 2t \, dt. \end{aligned}$$

Therefore, any eigenfunction corresponding to an eigenvalue $\lambda \neq 0$ must be of the form

$$f(x) = a \cdot \cos 2x + b \cdot \cos x + c,$$

where we shall find the constants a , b and c . We get by insertion into (5) that

$$\begin{aligned} Kf(x) &= \cos 2x \int_{-\pi}^{\pi} \frac{1}{2} (a \cdot \cos 2t + b \cdot \cos t + c) dt + \cos x \int_{-\pi}^{\pi} 2(a \cos 2t + b \cos t + c) \cos t dt \\ &\quad + \int_{-\pi}^{\pi} (a \cdot \cos 2t + b \cdot \cos t + c) dt \\ &\quad + \int_{-\pi}^{\pi} \frac{1}{2} (a \cdot \cos 2t + b \cdot \cos t + c) \cdot \cos 2t dt \\ &= c\pi \cdot \cos 2x + 2b\pi \cos x + 2\pi c + \frac{a\pi}{2}. \end{aligned}$$

This expression is equal to $\lambda a \cdot \cos 2x + \lambda b \cdot \cos x + \lambda c$, if and only if

$$\lambda a = c\pi, \quad \lambda b = 2\pi b, \quad \lambda c = 2\pi c + \frac{a\pi}{2}.$$

We immediately get the eigenvalue $\lambda = 2\pi$ with its corresponding eigenfunction $\cos x$.

The other eigenfunctions are found in the following way: The vectors (a, c) and $(c\pi, 2c\pi + \frac{a\pi}{2})$ must be proportional, so

$$0 = \begin{vmatrix} c & 2c + \frac{a}{2} \\ a & c \end{vmatrix} = c^2 - 2ac - \frac{a^2}{2} = (c - a)^2 - \frac{3}{2}a^2,$$

hence

$$c = a \pm \sqrt{\frac{3}{2}}a = \left\{ 1 \pm \sqrt{\frac{3}{2}} \right\} a,$$

corresponding to

$$\lambda = \frac{c\pi}{a} = \left\{ 1 \pm \sqrt{\frac{3}{2}} \right\} \pi.$$

For $\lambda_1 = \left\{ 1 + \sqrt{\frac{3}{2}} \right\} \pi$ we get the eigenfunction

$$f_1(x) = \cos 2x + 1 + \sqrt{\frac{3}{2}} \quad \left[= 2 \cos^2 x + \sqrt{\frac{3}{2}} \right].$$

For $\lambda_2 = \left\{ 1 - \sqrt{\frac{3}{2}} \right\} \pi$ we get the eigenfunction

$$f_2(x) = \cos 2x + 1 - \sqrt{\frac{3}{2}} \quad \left[= 2 \cos^2 x - \sqrt{\frac{3}{2}} \right].$$

For $\lambda = 2\pi$ we get the eigenfunction $f_3(x) = \cos x$.

There is no reason here to norm these eigenfunctions. We only notice that they span the same subspace of $L^2([-\pi, \pi])$ as 1 , $\cos x$, and $\cos 2x$ do.

It follows from $\overline{k(t, x)} = k(x, t)$ that K is self adjoint, so the null-space is simply the orthogonal complement of the subspace mentioned above. Thus we conclude that $\ker(K)$ is spanned by

$$\sin x, \sin 2x, \cos 3x, \sin 3x, \cos 4x, \sin 4x, \dots,$$

i.e. of the usual trigonometric basis with the exception of $1, \cos x$ and $\cos 2x$.

Example 1.10 Let K denote a self adjoint Hilbert-Schmidt operator on $L^2(I)$ with kernel k . Show that $\|K\| = \|k\|_2$ if and only if the spectrum for K consists of at most two points.

It follows from K being self adjoint that $\overline{k(t, x)} = k(x, t)$ and there exist an ortonormal sequence (φ_n) in $L^2(I)$ and a sequence (λ_n) of real numbers with $|\lambda_1| \geq |\lambda_2| \geq \dots$, where either $\lambda_n = 0$ eventually, or $\lambda_n \rightarrow 0$, such that

$$(6) \quad Ku = \sum_{n=1}^{+\infty} \lambda_n (u, \varphi_n) \varphi_n \quad \text{for } u \in L^2(I),$$

where every φ_n is an eigenfunction of the corresponding $\lambda_n \in \sigma_p(K)$, and where 0 is either an eigenvalue or belongs to the continuous spectrum $\sigma_c(K)$, and where

$$\sigma(K) = \{0\} \cup \sigma_p(K).$$

We shall prove that $\|K\| = \|k\|_2$, if and only if $\sigma(K)$ contains at most two points.

- 1) If $\sigma(K)$ only consists of one point, then $\sigma(K) = \{0\}$, and $Ku \equiv 0$, thus $k(x, t) = 0$ almost everywhere, and it follows trivially that $\|K\| = \|k\|_1 = 0$.
- 2) If $\sigma(K)$ contains two points, then it follows from the introducing argument that we necessarily must have

$$\sigma(M) = \{0, \lambda\},$$

so the operator is described by

$$Ku = (u, \varphi) \varphi = \lambda \int_a^b \varphi(x) \overline{\varphi(t)} u(t) dt,$$

from which we derive that

$$k(x, t) = \lambda \varphi(t) \varphi(x).$$

Clearly, $\|K\| = \lambda$. Because $\|\varphi\|_2 = 1$, we get

$$\|k\|_2^2 = \int_a^b \int_a^b |k(x, t)|^2 dx dt = |\lambda|^2 \int_a^b \int_a^b |\varphi(x)|^2 |\varphi(t)|^2 dx dt = |\lambda|^2.$$

Hence $\|k\|_2 = |\lambda| = \|K\|$ in this case.

- 3) If $\sigma(K)$ contains more than two points, then

$$\|K\| = \max |\lambda_n| = |\lambda_1|.$$

Furthermore, we get by the computation

$$Ku(x) = \int_I k(x, y) u(t) dt = \sum_{n=1}^{+\infty} \lambda_n (u, \varphi_n) \varphi_n(x) = \int_I \sum_{n=1}^{+\infty} \lambda_n \varphi_n(x) \overline{\varphi_n(t)} u(t) dt,$$

that

$$\|k\|_2^2 = \sum_{n=1}^{+\infty} \lambda_n^2 > \lambda_1^2 = \|K\|^2,$$

and the claim is proved.

Please click the advert



Nido

Luxurious accommodation

Central zone 1 & 2 locations

Meet hundreds of international students

BOOK NOW and get a £100 voucher from voucherexpress

Nido Student Living - London

Visit www.NidoStudentLiving.com/Bookboon for more info.

+44 (0)20 3102 1060

Example 1.11 Let $\{e_1, e_2, \dots, e_p\}$ denote a finite orthonormal set in $L^2(I)$, and let the Hilbert-Schmidt operator K be given by the kernel

$$k(x, y) = \sum_{i=1}^p \sum_{j=1}^p k_{ij} e_i(x) e_j(y).$$

Find the trace $\text{tr}(K)$.

We say that the operator K has a canonical kernel of finite rank.

This example is trivial,

$$\text{tr}(K) = \int_I k(x, x) dx = \int_I \sum_{i=1}^p \sum_{j=1}^p k_{ij} e_i(x) e_j(x) dx = \sum_{i=1}^p \sum_{j=1}^p k_{ij} \delta_{ij} = \sum_{i=1}^p k_{ii}.$$

Note that this corresponds to the trace of matrix (k_{ij}) .

Example 1.12 Denote by K a self adjoint Hilbert-Schmidt operator on $L^2(I)$ of kernel k . Prove that K is a general Hilbert-Schmidt operator (cf. the definition in EXAMPLE 1.1), and find the Hilbert-Schmidt norm $\|K\|_{\text{HS}}$.

Put

$$Ku = \sum_{n=1}^{+\infty} \lambda_n (u, \varphi_n) \varphi_n.$$

It follows from VENTUS, HILBERT SPACES ETC., EXAMPLE 2.7 that

$$t_{jk} = (K\varphi_j, \varphi_k) = \left(\sum_{n=1}^{+\infty} \lambda_n (\varphi_j, \varphi_n) \varphi_n, \varphi_k \right) = (\lambda_j \varphi_j, \varphi_k) = \lambda_j \delta_{jk},$$

thus $t_{jj} = \lambda_j$ and $t_{jk} = 0$ for $j \neq k$.

Then by EXAMPLE 1.1, K is a general Hilbert-Schmidt operator, if

$$\sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |t_{jk}|^2 < +\infty,$$

because it was proved that this number is independent of the choice of orthonormal basis. Furthermore, it follows from EXAMPLE 1.2 that

$$\|K\|_{\text{HS}} = \left\{ \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |t_{jk}|^2 \right\}^{\frac{1}{2}}.$$

In the present case we get

$$\|K\|_{\text{HS}} = \left\{ \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} |\lambda_j|^2 \delta_{jk} \right\}^{\frac{1}{2}} = \left\{ \sum_{j=1}^{+\infty} |\lambda_j|^2 \right\}^{\frac{1}{2}} = \|k\|_2.$$

Example 1.13 *Let*

$$k(x, t) = \{\sin(x) + \sin(t)\}^2 - \frac{1}{8}$$

be the kernel for a Hilbert-Schmidt operator K on the complex Hilbert space $L^2([-\pi, \pi])$.

Show that K is self adjoint and express the range $K(L^2([-\pi, \pi]))$ of K with the help of the non-normalized basis

$$1, \cos(x), \sin(x), \cos(2x), \sin(2x), \dots$$

Find all non-zero eigenvalues and corresponding eigenfunctions for K , and determine $\sigma(K)$.

Solve the equation $Ku = \pi u - \frac{5\pi}{4}$ in $L^2([-\pi, \pi])$.

1) Clearly, $k(x, t) \in L^2([-\pi, \pi] \times [-\pi, \pi])$, and

$$\overline{k(t, x)} = (\sin t + \sin x)^2 - \frac{1}{8} = k(x, t),$$

thus $k(x, t)$ is Hermitian, and K is a self adjoint Hilbert-Schmidt-operator. It follows from

$$\begin{aligned} k(x, t) &= (\sin x + \sin t)^2 - \frac{1}{8} = \sin^2 x + 2 \sin x \cdot \sin t + \sin^2 t - \frac{1}{8} \\ &= -\frac{1}{2} \cos 2x + 2 \sin x \cdot \sin t - \frac{1}{2} \cos 2t + \frac{7}{8}, \end{aligned}$$

that

$$\begin{aligned} (7) \quad Kf(x) &= \left\{ -\frac{1}{2} \int_{-\pi}^{\pi} f(t) dt \right\} \cos 2x + \left\{ 2 \int_{-\pi}^{\pi} f(t) \sin t dt \right\} \sin x \\ &\quad + \left\{ -\frac{1}{2} \int_{-\pi}^{\pi} f(t) \cos 2t dt + \frac{7}{8} \int_{-\pi}^{\pi} f(t) dt \right\} \cdot 1, \end{aligned}$$

and we conclude that the range $K(L^2([-\pi, \pi]))$ is spanned by 1 , $\sin x$ and $\cos 2x$.

(Choose e.g. suitable linear combinations of these three functions in order to conclude that the dimension is 3).

2) An eigenfunction f corresponding to an eigenvalue $\lambda \neq 0$ must necessarily lie in the range, thus it is of the form

$$f(x) = a \cdot \cos 2x + b \cdot \sin x + c, \quad a, b, c \in \mathbb{C}.$$

When we insert this expression into (7) and then apply that 1 , $\sin x$ and $\cos 2x$ are mutually orthogonal, we get

$$\begin{aligned} Kf(x) &= \left\{ -\frac{1}{2} c \cdot 2\pi \right\} \cos 2x + \left\{ 2b \cdot \frac{2\pi}{2} \right\} \sin x + \left\{ -\frac{1}{2} a \cdot \frac{2\pi}{2} + \frac{7}{8} c \cdot 2\pi \right\} \cdot 1 \\ &= -c\pi \cdot \cos 2x + 2b\pi \cdot \sin x + \left\{ \frac{7\pi}{4} c - \frac{\pi}{2} a \right\} \cdot 1. \end{aligned}$$

We have for comparison,

$$\lambda f(x) = \lambda a \cdot \cos 2x + \lambda b \cdot \sin x + \lambda c \cdot 1.$$

The coefficient b occurs only in connection with $\sin x$, hence we conclude that $\sin x$ is an eigenfunction corresponding to the eigenvalue $\lambda = 2\pi$.

Assume that $b = 0$. If $a \cdot \cos 2x + c$ is an eigenfunction, then the vectors

$$\left(-c\pi, \frac{7\pi}{4}c - \frac{\pi}{2}a\right) = \pi \left(-c, \frac{7}{4}c - \frac{1}{2}a\right) \quad \text{og} \quad (a, c)$$

must be proportional with the eigenvalue $\lambda = -\frac{c}{a}\pi$ as the factor of proportion. Thus we get the condition

$$\begin{vmatrix} a & -c \\ c & \frac{7}{4}c - \frac{1}{2}a \end{vmatrix} = c^2 + \frac{7}{4}ac - \frac{1}{2}a^2 = 0.$$

By solving this equation with respect to c we get

$$c = -\frac{7}{8}a \pm \sqrt{\frac{49}{64}a^2 + \frac{1}{2}a^2} = -\frac{7}{8}a \pm \sqrt{\frac{81}{64}a^2} = -\frac{7}{8}a \pm \frac{9}{8}a.$$

We have now two possibilities:

- a) For $c = -\frac{7}{8}a - \frac{9}{8}a = -2a$ we get $\lambda = -\frac{c}{a}\pi = 2\pi$, corresponding to the eigenfunction $\cos 2x - 2$.
- b) For $c = -\frac{7}{8}a + \frac{9}{8}a = \frac{1}{4}a$ we get $\lambda = -\frac{c}{a}\pi = -\frac{\pi}{4}$, corresponding to the eigenfunction $\cos 2x + \frac{1}{4}$.

Summing up,

$$\begin{aligned} \lambda_1 &= 2\pi, & \varphi_1(x) &= \sin x, \\ \lambda_2 &= 2\pi, & \varphi_2(x) &= \cos 2x - 2, \\ \lambda_3 &= -\frac{\pi}{4}, & \varphi_3(x) &= \cos 2x + \frac{1}{4}. \end{aligned}$$

Notice that $\lambda_1 = \lambda_2$, and that the eigenfunctions are not normed.

It follows e.g. from $(K \cos)(x) = 0$ that $\ker(K) \neq \emptyset$, thus

$$\sigma(K) = \sigma_p = \left\{0, -\frac{\pi}{2}, 2\pi\right\}.$$

- 3) The equation $Ku = \pi u - \frac{5\pi}{4}$ can be solved in several ways:

First method. The coefficient π of u on the right hand side of the equation does not belong to the spectrum, $\pi \notin \sigma(K)$, hence the solution is unique. Because

$$-\frac{5\pi}{4} = \frac{5\pi}{9}(\cos 2x - 2) - \frac{5\pi}{9}\left(\cos 2x + \frac{1}{4}\right),$$

we see that $-\frac{5\pi}{4}$ lies in the subspace spanned by the eigenvectors

$$\varphi_2(x) = \cos 2x - 2 \quad \text{and} \quad \varphi_3(x) = \cos 2x + \frac{1}{4}.$$

Thus we *guess* a solution of the structure

$$u(x) = a \cdot (\cos 2x - 2) + b \cdot \left(\cos 2x + \frac{1}{4} \right).$$

We get by insertion of this structure that

$$\begin{aligned} Ku(x) - \pi u(x) &= 2\pi a \cdot (\cos 2x - 2) - \frac{\pi}{4} b \cdot \left(\cos 2x + \frac{1}{4} \right) \\ &\quad - \pi a (\cos 2x - 2) - \pi b \left(\cos 2x + \frac{1}{4} \right) \\ &= \pi a (\cos 2x - 2) - \frac{5\pi}{4} b \left(\cos 2x + \frac{1}{4} \right) \\ &= \pi \left(a - \frac{5}{4} b \right) \cos 2x - \pi \left(2a + \frac{5}{16} - b \right). \end{aligned}$$

Please click the advert

I joined MITAS because
I wanted **real responsibility**

The Graduate Programme
for Engineers and Geoscientists
Maersk.com/Mitas



Month 16

I was a construction
supervisor in
the North Sea
advising and
helping foremen
solve problems

Real work
International opportunities
Three work placements



 **MAERSK**

Download free ebooks at bookboon.com

This expression is equal to $-\frac{5\pi}{4}$, if

$$a = \frac{5}{4}b \quad \text{and} \quad 2 \cdot \frac{5}{4}b + \frac{1}{4} \cdot \frac{5}{4}b = \frac{5}{4},$$

hence $\frac{9}{4}b = 1$ and $b = \frac{4}{9}$, $a = \frac{5}{9}$. Finally, we get by insertion,

$$u(x) = \frac{5}{9}(\cos 2x - 2) + \frac{4}{9}\left(\cos 2x + \frac{1}{4}\right) = \cos 2x - 1 = -2\sin^2 x.$$

Method 1a. A variant of the FIRST METHOD is to guess a solution of the form

$$u(x) = a \cdot \cos 2x + c.$$

Then apply the previous computation from (2) to get

$$Ku(x) = -c\pi \cdot \cos 2x + \left\{ \frac{7\pi}{4}c - \frac{\pi}{2}a \right\},$$

and

$$-\pi u(x) = -a\pi \cdot \cos 2x - c\pi,$$

hence

$$Ku(x) - \pi u(x) = -(a + c)\cos 2x + \frac{3\pi}{4}c - \frac{\pi}{2}a.$$

This expression is equal to $-\frac{5\pi}{4}$, if and only if

$$c = -a \quad \text{and} \quad -\frac{5\pi}{4} = \frac{3\pi}{4}c - \frac{\pi}{2}a = -\frac{5\pi}{4}a,$$

thus $a = 1$ and $c = -1$, and the unique solution is given by

$$u(x) = \cos 2x - 1 = -2\sin^2 x.$$

Second method. It is also possible to apply the standard method. A straightforward computation where we explicitly use the previously found eigenfunctions (these should then be normed), would demand a lot of energy, although one at different stages could apply one of the two variants above.

We shall show below how this might be carried out. First put

$$\varphi_1(x) = \sin x, \quad \varphi_2(x) = \cos 2x - 2, \quad \varphi_3(x) = \cos 2x + \frac{1}{4}.$$

Let $\{\varphi_n \mid n \geq 4\}$ denote an orthonormal basis of the null-space $\ker(K)$. Then a solution of the equation

$$Ku = \pi u - \frac{5\pi}{4},$$

has the structure

$$u = \sum_{n=1}^{+\infty} a_n \varphi_n, \quad \text{where } \sum_{n=4}^{+\infty} |a_n|^2 < +\infty.$$

Put $f(x) = -\frac{5\pi}{4}$. It follows from

$$(f, \varphi_n) = \left(-\frac{5\pi}{4}, \varphi_n\right) = 0 \quad \text{for } n \in \mathbb{N} \setminus \{2, 3\},$$

and

$$f(x) = -\frac{5\pi}{4} = c_2(\cos 2x - 2) + c_3 \left(\cos 2x + \frac{1}{4}\right) = (c_2 + c_3) \cos 2x - \left(2c_2 - \frac{1}{4}c_3\right),$$

that $c_3 = -c_2$, and

$$2c_2 - \frac{1}{4}c_3 = 2c_2 + \frac{1}{4}c_2 = \frac{9}{4}c_2 = \frac{5\pi}{4},$$

thus

$$c_2 = \frac{5\pi}{9} \quad \text{and} \quad c_3 = -\frac{5\pi}{9}.$$

Then we get by insertion into the equation

$$Ku - \pi u = -\frac{5\pi}{4}$$

that

$$\begin{aligned} Ku - \pi u &= \lambda_1 a_1 \varphi_1 + \lambda_2 a_2 \varphi_2 + \lambda_3 a_3 \varphi_3 - \sum_{n=1}^{+\infty} a_n \varphi_n \\ &= (2\pi - \pi)a_1 \varphi_1 + (2\pi - \pi)a_2 \varphi_2 - \left(\frac{\pi}{4} + \pi\right)a_3 \varphi_3 - \pi \sum_{n=4}^{+\infty} a_n \varphi_n \\ &= \pi a_1 \varphi_1 + \pi a_2 \varphi_2 - \frac{5\pi}{4} a_3 \varphi_3 - \pi \sum_{n=4}^{+\infty} a_n \varphi_n \\ &= -\frac{5\pi}{4} = c_2 \varphi_2 + c_3 \varphi_3, \end{aligned}$$

and we derive that

$$a_1 = 0, \quad a_2 = \frac{1}{\pi} c_2 = \frac{5}{9}, \quad a_3 = -\frac{4}{5\pi} c_3 = \frac{4}{9}, \quad a_n = 0 \text{ for } n \geq 4,$$

hence

$$u(x) = \frac{5}{9}(\cos 2x - 2) + \frac{4}{9} \left(\cos 2x + \frac{1}{4}\right) = \cos 2x - 1 = -2 \sin^2 x.$$

Example 1.14 Let $k(x, t) = x + t + 2xt$ be the kernel for the Hilbert-Schmidt operator K on the Hilbert space $H = L^2([-1, 1])$.

Show that K is self adjoint and determine the range $K(H)$.

Find all non-zero eigenvalues and corresponding eigenfunctions for K , and determine $\sigma(K)$ as well as $\|K\|$.

Express Kf , $f \in H$, with the help of the Legendre polynomials (P_n) .

Let $f(x) = \cosh(1) \cosh(x) - \cosh(2x)$. Show that $(f, P_0) = (f, P_1) = 0$ and solve the equation

$$Ku(x) + u(x) = f(x).$$

1) It follows from

$$\overline{k(t, x)} = \overline{t + x + 2tx} = x + t + 2xt = k(x, t),$$

that the kernel is Hermitian, thus K is self adjoint. We conclude from

$$Kf(x) = \int_{-1}^1 (x + t + 2xt)f(t) dt = x \int_{-1}^1 (1 + 2t)f(t) dt + \int_{-1}^1 t f(t) dt,$$

that the range is $K(L^2([-1, 1])) = \text{span}\{1, x\}$.

2) The only possible eigenfunctions must be of the form $f(x) = ax + b$. We get by insertion the condition

$$\lambda f(x) = \lambda ax + \lambda b = Kf(x) = x \int_{-1}^1 (1 + 2t)(at + b) dt + \int_{-1}^1 t(at + b) dt,$$

hence

$$\lambda a = \int_{-1}^1 (1 + 2t)(at + b) dt = \int_{-1}^1 \{2at^2 + (a + 2b)t + b\} dt = \frac{4}{3}a + 2b$$

and

$$\lambda b = \int_{-1}^1 (at^2 + bt) dt = \frac{2a}{3}.$$

Hence,

$$\lambda^2 a = \frac{4}{3}a\lambda + 2\lambda b = \frac{4}{3}\lambda a + \frac{4}{3}a.$$

If $a = 0$, then $2b = \left(\lambda - \frac{4}{3}\right)a = 0$, which leads to nothing, so we may assume that $a \neq 0$, e.g. $a = 1$. Then

$$\lambda^2 - \frac{4}{3}\lambda - \frac{4}{3} = 0,$$

i.e.

$$\lambda = \frac{2}{3} \pm \sqrt{\frac{4}{9} + \frac{4}{3}} = \frac{2}{3} \pm \sqrt{\frac{16}{9}} = \frac{2}{3} \pm \frac{4}{3} = \begin{cases} 2, \\ -\frac{2}{3}. \end{cases}$$

If $\lambda_1 = 2$ and $a = 1$, then $b = \frac{1}{\lambda_1} \cdot \frac{2a}{3} = \frac{1}{3}$, and the corresponding eigenfunction is

$$\varphi_1(x) = x + \frac{1}{3}, \quad \lambda_1 = 2.$$

If $\lambda_2 = -\frac{2}{3}$ and $a = 1$, then $b = \frac{1}{\lambda_2} \cdot \frac{2a}{3} = -\frac{3}{2} \cdot \frac{2}{3} = -1$, and the corresponding eigenfunction is

$$\varphi_2(x) = x - 1, \quad \lambda_2 = -\frac{2}{3}.$$

Since K is self adjoint and of Hilbert-Schmidt-type, $\|K\|$ is the absolute value of the eigenvalue of largest absolute value,

$$\|K\| = 2.$$

Finally,

$$\sigma(K) = \sigma_p(K) = \left\{ -\frac{2}{3}, 0, 2 \right\},$$

and every function, which is orthogonal on both φ_1 and φ_2 , i.e. on both 1 and x by a change of basis, must lie in the eigenspace corresponding to $\lambda = 0$.

Please click the advert



Brain power

By 2020, wind could provide one-tenth of our planet's electricity needs. Already today, SKF's innovative know-how is crucial to running a large proportion of the world's wind turbines.

Up to 25 % of the generating costs relate to maintenance. These can be reduced dramatically thanks to our systems for on-line condition monitoring and automatic lubrication. We help make it more economical to create cleaner, cheaper energy out of thin air.

By sharing our experience, expertise, and creativity, industries can boost performance beyond expectations. Therefore we need the best employees who can meet this challenge!

The Power of Knowledge Engineering

Plug into The Power of Knowledge Engineering.
Visit us at www.skf.com/knowledge

SKF

- 3) It is well-known that the Legendre polynomials form an orthogonal system on $L^2([-1, 1])$. We have in particular,

$$P_0(t) = 1 \quad \text{and} \quad P_1(t) = t,$$

and since $\text{span}\{P_0, P_1\} = K(L^2([-1, 1]))$, we infer that

$$KP_n = 0 \quad \text{for every } n \geq 2.$$

It follows that if $f = \sum_{n=0}^{+\infty} a_n P_n$, then

$$\begin{aligned} Kf(x) &= K\left(\sum_{n=0}^{+\infty} a_n P_n\right)(x) = K\left(\sum_{n=0}^1 a_n P_n\right)(x) \\ &= K(a_0 + a_1 t)(x) = \int_{-1}^1 (a_0 + a_1 t)(x + t + 2xt) dt \\ &= \int_{-1}^1 \{a_0 x + a_0 t + 2a_0 x \cdot t + a_1 x \cdot t + a_1(1 + 2x)t^2\} dt \\ &= 2a_0 x + \frac{2}{3} a_1(1 + 2x) = \left(2a_0 + \frac{4}{3} a_1\right)x + \frac{2}{3} a_1 \\ &= \left(2a_0 + \frac{4}{3} a_1\right) P_1(x) + \frac{2}{3} a_1 P_0(x). \end{aligned}$$

- 4) Let $f(x) = \cosh 1 \cdot \cosh x - \cosh 2x$. Then

$$\begin{aligned} (f, P_0) &= \int_{-1}^1 \{\cosh 1 \cdot \cosh x - \cosh 2x\} dx = \cosh 1 \cdot [\sinh x]_{-1}^1 - \left[\frac{1}{2} \sinh 2x\right]_{-1}^1 \\ &= \cosh 1 \cdot 2 \sinh 1 - \frac{1}{2} \cdot 2 \sinh 2 = \sinh 2 - \sinh 2 = 0, \end{aligned}$$

and

$$(f, P_1) = \int_{-1}^1 \{\cosh 1 \cdot \cosh x - \cosh 2x\} \cdot x dx = 0,$$

because the integrand is an odd function, and because we integrate over a finite symmetric interval.

Finally, we shall solve the equation

$$Ku(x) + u(x) = \cosh 1 \cdot \cosh x - \cosh 2x.$$

If

$$u = \sum_{n=0}^{+\infty} a_n P_n \quad \text{and} \quad \cosh 1 \cdot \cosh x - \cosh 2x = \sum_{n=2}^{+\infty} b_n P_n,$$

then it follows from the above that

$$\begin{aligned} \frac{2}{3} a_1 P_0 + \left(2a_0 + \frac{4}{3} a_1\right) P_1 + a_0 P_0 + a_1 P_1 + \sum_{n=2}^{+\infty} a_n P_n \\ = \sum_{n=2}^{+\infty} b_n P_n = \cosh 1 \cdot \cosh x - \cosh 2x, \end{aligned}$$


and we conclude that $a_n = b_n$ for $n \geq 2$ and that

$$\begin{cases} a_0 + \frac{2}{3}a_1 = 0, \\ 2a_0 + \frac{1}{3}a_1 = 0, \end{cases} \quad \text{hence } a_0 = a_1 = 0,$$

and whence

$$u = \sum_{n=2}^{+\infty} a_n P_n = \sum_{n=2}^{+\infty} b_n P_n = \cosh 1 \cdot \cosh x - \cosh 2x.$$

Please click the advert



Are you considering a European business degree?

LEARN BUSINESS at university level. We mix cases with cutting edge research working individually or in teams and everyone speaks English. Bring back valuable knowledge and experience to boost your career.

MEET a culture of new foods, music and traditions and a new way of studying business in a safe, clean environment – in the middle of Copenhagen, Denmark.

ENGAGE in extra-curricular activities such as case competitions, sports, etc. – make new friends among CBS' 18,000 students from more than 80 countries.

Copenhagen Business School
HANDELSHØJSKOLEN

EFMD
EQUIS

AACSB
ACCREDITED

Accredited by
Association
of MBAs

CEMS

EFMD
EQUIS

AACSB
ACCREDITED

Accredited by
Association
of MBAs

CEMS

EFMD
EQUIS

AACSB
ACCREDITED

Accredited by
Association
of MBAs

CEMS

See what we look like
and how we work on cbs.dk

Download free ebooks at bookboon.com

Example 1.15 In $L^2([-\pi, \pi])$ we consider the orthonormal basis (e_n) , $n \in \mathbb{Z}$, where

$$e_n(t) = \frac{1}{\sqrt{2\pi}} e^{int}.$$

1. Let $\varphi : \mathbb{R} \rightarrow \mathbb{C}$ denote a continuous function with period 2π , and assume that $\varphi(-x) = \overline{\varphi(x)}$ for all $x \in \mathbb{R}$. Show that

$$Ku(x) = \int_{-\pi}^{\pi} \varphi(x-t) u(t) dt$$

defines a selfadjoint Hilbert-Schmidt operator on $L^2([-\pi, \pi])$.

2. Show that all e_n are eigenfunctions for K .

From now on we assume that φ is the periodic extension from $[-\pi, \pi]$ to \mathbb{R} of the function

$$\varphi(x) = 1 - \frac{|x|}{\pi}.$$

3. Calculate the spectrum of K .

4. Solve the equation

$$Ku = \frac{2}{\pi} u + f \quad \text{in } L^2([-\pi, \pi]),$$

where $f(x) = \sin^2(x) + \sin(x)$.

5. Solve the equation

$$Ku = \frac{4}{\pi} u + 1 \quad \text{in } L^2([-\pi, \pi]).$$

- 1) The kernel is

$$k(x, t) = \varphi(x-t), \quad x, t \in [-\pi, \pi],$$

where

$$\begin{aligned} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} |\varphi(x-t)|^2 dt dx &= \int_{-\pi}^{\pi} \left\{ \int_{-\pi-t}^{\pi-t} |\varphi(u)|^2 du \right\} dx = \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} |\varphi(u)|^2 du dx \\ &= 2\pi \|\varphi\|_2^2 < +\infty, \end{aligned}$$

proving that K is a Hilbert-Schmidt operator.

ALTERNATIVELY, φ is continuous on a compact set, hence $|\varphi(x)| \leq c$ for $x \in [-\pi, \pi]$. Then apply the periodicity to get the estimate

$$\int_{-\pi}^{\pi} \int_{-\pi}^{\pi} |\varphi(x-t)|^2 dt dx \leq c^2 (2\pi)^2 = 4\pi^2 c^2 < +\infty. \quad \diamond$$

From $\varphi(-x) = \overline{\varphi(x)}$ follows that

$$\overline{k(t, x)} = \overline{\varphi(t-x)} = \varphi(x-t) = k(x, t),$$

which shows that the kernel is Hermitian, thus K is self adjoint.

2) By insertion of $e_n(x)$ follows by a change of variable,

$$\begin{aligned} K e_n(x) &= \int_{-\pi}^{\pi} \varphi(x-t) e_n(t) dt = \int_{x-\pi}^{x+\pi} \varphi(u) e_n(x-u) du \\ &= \int_{x-\pi}^{x+\pi} \varphi(u) \cdot e^{-inu} du \cdot \frac{1}{\sqrt{2\pi}} e^{inx} = \int_{-\pi}^{\pi} \varphi(u) e^{-inu} du \cdot e_n(x), \end{aligned}$$

from which follows that every $e_n(x)$, $n \in \mathbb{Z}$, is an eigenfunction for K .

Conversely, if ψ is an eigenfunction, then $\psi = \sum c_n e_n$, hence ψ must lie in the subspace corresponding to the e_n , which have the same eigenvalue. This means that the eigenvalues are

$$\int_{-\pi}^{\pi} \varphi(u) e^{-inu} du, \quad n \in \mathbb{Z},$$

and it suffices only to look at the eigenfunctions $e_n(x)$, $n \in \mathbb{Z}$, in the following.




Figure 1: The graph of the function φ .

3) If $\varphi(x) = 1 - \frac{|x|}{\pi}$ for $x \in [-\pi, \pi]$, then we have in particular that $\varphi(-x) = \overline{\varphi(x)}$, and that φ is continuous – also after a periodic extension. Therefore, we are again in the situation above. If $n \neq 0$, then the eigenvalues are given by

$$\begin{aligned} \int_{-\pi}^{\pi} \left(1 - \frac{|x|}{\pi}\right) e^{-inx} dx &= - \int_{-\pi}^{\pi} \frac{|x|}{\pi} e^{-inx} dx = -\frac{2}{\pi} \int_0^{\pi} x \cos(nx) dx \\ &= 0 + \frac{2}{n\pi} \int_0^{\pi} \sin(nx) dx = \frac{2\{1 - (-1)^n\}}{\pi n^2}. \end{aligned}$$

For $n = 0$ we instead get by considering an area on the figure,

$$\int_{-\pi}^{\pi} \left(-\frac{|x|}{\pi}\right) dx = \pi.$$

ALTERNATIVELY,

$$\int_{-\pi}^{\pi} \left(a - \frac{|x|}{\pi} \right) dx = 2\pi - \frac{2}{\pi} \int_0^{\pi} x dx = 2\pi - \frac{2\pi^2}{2\pi} = \pi.$$

Summing up,

$$\lambda_0 = \pi, \quad \begin{cases} \lambda_{2n} = 0, & n \in \mathbb{Z} \setminus \{0\}, \\ \lambda_{2n+1} = \frac{4}{\pi(2n+1)^2}, & n \in \mathbb{Z}, \end{cases}$$

and we conclude that the spectrum is

$$\sigma(K) = \sigma_p(K) = \{0, \pi\} \cup \left\{ \frac{4}{\pi(2n+1)^2} \mid n \in \mathbb{N}_0 \right\}.$$

Notice that the eigenspace corresponding to each eigenvalue of the form $\frac{4}{\pi(2n+1)^2}$ is of dimension 2, while the eigenspace corresponding to $\lambda_0 = \pi$ is only of dimension 1.

4) Let

$$u = \sum c_n e_n = c_0 e_0 + \sum_{n \neq 0} c_{2n} e_{2n} + \sum_{n \in \mathbb{Z}} c_{2n+1} e_{2n+1}.$$

Then

$$\begin{aligned} f(x) &= \sin^2 x + \sin x = \frac{1 - \cos 2x}{2} + \sin x = \frac{1}{2} + \frac{e^{ix} - e^{-ix}}{2i} - \frac{e^{2ix} + e^{-2ix}}{4} \\ &= \frac{\sqrt{2\pi}}{2} e_0(x) + i \frac{\sqrt{2\pi}}{2} e_{-1}(x) - i \frac{\sqrt{2\pi}}{2} e_1(x) - \frac{\sqrt{2\pi}}{4} e_2(x) - \frac{\sqrt{2\pi}}{4} e_{-2}(x) \\ &= Ku - \frac{2}{\pi} u \\ &= \left(\pi - \frac{2}{\pi} \right) c_0 e_0(x) + \sum_{n \in \mathbb{Z} \setminus \{0\}} \left(-\frac{2}{\pi} \right) c_{2n} e_{2n}(x) + \sum_{n \in \mathbb{Z}} \left\{ \frac{4}{(2n+1)^2 \pi} - \frac{2}{\pi} \right\} c_{2n+1} e_{2n+1}(x). \end{aligned}$$

It follows from $\frac{2}{\pi} \notin \sigma_p(K) = \sigma(K)$ by identification that

$$c_0 = \frac{\sqrt{2\pi}}{2} \cdot \frac{1}{\pi - \frac{2}{\pi}} = \sqrt{2\pi} \cdot \frac{\pi}{2(\pi^2 - 2)},$$

and

$$c_{-1} = i \frac{\sqrt{2\pi}}{2} \cdot \frac{1}{\frac{4}{\pi} - \frac{2}{\pi}} = i \sqrt{2\pi} \cdot \frac{\pi}{4}, \quad c_1 = \overline{c_{-1}} = -i \sqrt{2\pi} \cdot \frac{\pi}{4},$$

and

$$c_{-2} = c_2 = -\frac{\sqrt{2\pi}}{4} \cdot \frac{1}{-\frac{2}{\pi}} = \sqrt{2\pi} \cdot \frac{\pi}{8}, \quad \text{and } c_n = 0 \quad \text{otherwise.}$$

This implies that

$$\begin{aligned} u(x) &= \frac{\pi}{2(\pi^2 - 2)} \sqrt{2\pi} e_0(x) + \frac{\pi}{2} \cdot \frac{\sqrt{2\pi}}{2i} \{e_1(x) - e_{-1}(x)\} + \frac{\pi}{4} \frac{\sqrt{2\pi}}{2} \{e_2(x) + e_{-2}(x)\} \\ &= \frac{\pi}{2(\pi^2 - 2)} + \frac{\pi}{2} \sin x + \frac{\pi}{4} \cos 2x. \end{aligned}$$

5) In this case, $\frac{4}{\pi}$ is an eigenvalue corresponding to the eigenvectors $e_1(x)$ and $e_{-1}(x)$. Since $1 = \sqrt{2\pi} e_0$ is orthogonal to e_1 and e_{-1} , we get

$$u = c_{-1}e_{-1} + c_1e_1 + c_0e_0,$$

where c_{-1} and c_1 are arbitrary constants, and

$$1 = K(c_0e_0) - \frac{4}{\pi}c_0e_0 = \left(\pi - \frac{4}{\pi}\right)c_0e_0 = \left(\pi - \frac{4}{\pi}\right)c_0 \cdot \frac{1}{\sqrt{2\pi}},$$

hence

$$c_0 = \frac{\sqrt{2\pi}}{\pi - \frac{4}{\pi}} = \frac{\pi\sqrt{2\pi}}{\pi^2 - 4},$$

Please click the advert



The financial industry needs a strong software platform

That's why we need you

SimCorp is a leading provider of software solutions for the financial industry. We work together to reach a common goal: to help our clients succeed by providing a strong, scalable IT platform that enables growth, while mitigating risk and reducing cost. At SimCorp, we value commitment and enable you to make the most of your ambitions and potential.

Are you among the best qualified in finance, economics, IT or mathematics?

Find your next challenge at
www.simcorp.com/careers



www.simcorp.com

MITIGATE RISK | REDUCE COST | ENABLE GROWTH

and we get the solutions

$$u(x) = \frac{\pi\sqrt{2\pi}}{\pi^2 - 4} + \tilde{c}_1 e^{ix} + \tilde{c}_{-1} e^{-ix},$$

where \tilde{c}_1 and $\tilde{c}_{-1} \in \mathbb{C}$ are arbitrary constants.

Example 1.16 Let H denote the Hilbert space $L^2([0, 2\pi])$ with the subspace $F = C([0, 2\pi])$, and let K denote the integral operator on H with the kernel

$$k(x, t) = \begin{cases} \frac{i}{2} \exp\left(\frac{i}{2}(x-t)\right), & \text{if } 0 \leq t < x \leq 2\pi, \\ 0 & \text{if } 0 \leq t = x \leq 2\pi, \\ -\frac{i}{2} \exp\left(\frac{i}{2}(x-t)\right), & \text{if } 0 \leq x < t \leq 2\pi. \end{cases}$$

- 1) Show that K is a self adjoint Hilbert-Schmidt operator.
- 2) Assume that F is equipped with the sup-norm. Show that $K : H \rightarrow F$ is continuous.
- 3) Now let S denote the restriction of K to F (considered as a subspace of H). Show that S is injective and that S^{-1} is given by

$$D(S^{-1}) = \{g \in C^1([0, 2\pi]) \mid g(0) = g(2\pi)\},$$

and

$$S^{-1}g = -i g' - \frac{1}{2} g \quad \text{for } g \in D(S^{-1}).$$

- 4) Find all normalized eigenfunctions and associated eigenvalues for S^{-1} . Show that all eigenvalues are simple and that the set of normalized eigenfunctions is an orthonormal system in H .
- 5) Show that the eigenfunctions for S^{-1} are also eigenfunctions for K and find the associated eigenvalues. Justify that all eigenfunctions for K are given this way, and write the kernel for K using the normalized eigenfunctions.
- 6) Let $f \in H$ be given by the Fourier expansion

$$f = \sum_{n=-\infty}^{\infty} c_n e^{inx}.$$

Expand Kf using the Fourier coefficients c_n instead of f .

- 1) The kernel $k(x, t)$ is bounded and continuous for $t \neq x$ in the compact set $[0, 2\pi]^2$, hence $k \in L^2([0, 2\pi]^2)$ with

$$\|k\|_2^2 = \int_0^{2\pi} \left\{ \int_0^{2\pi} |k(x, t)|^2 dt \right\} dx = \frac{1}{4} \cdot (2\pi)^2 = \pi^2,$$

i.e. $\|k\|_2 = \pi$. This shows that K is a Hilbert-Schmidt operator.

We see from

$$\begin{aligned} \overline{k(t, x)} &= \begin{cases} -\frac{i}{2} \exp\left(-\frac{i}{2}(t-x)\right), & \text{for } 0 \leq x < t \leq 2\pi, \\ 0 & \text{for } 0 \leq x = t \leq 2\pi, \\ \frac{i}{2} \exp\left(-\frac{i}{2}(t-x)\right), & \text{for } 0 \leq t < x \leq 2\pi, \end{cases} \\ &= \begin{cases} \frac{i}{2} \exp\left(\frac{i}{2}(x-t)\right), & \text{for } 0 \leq t < x \leq 2\pi, \\ 0 & \text{for } 0 \leq t = x \leq 2\pi, \\ -\frac{i}{2} \exp\left(\frac{i}{2}(x-t)\right), & \text{for } 0 \leq x < t \leq 2\pi, \end{cases} \\ &= k(x, t), \end{aligned}$$

that $k(x, t)$ is Hermitian,, thus K is a self adjoint Hilbert-Schmidt operator.

2) The operator K is described by

$$\begin{aligned} Kf(x) &= \int_0^{2\pi} k(x, t) f(t) dt = \frac{i}{2} \int_0^x \exp\left(\frac{i}{2}(x-t)\right) f(t) dt - \frac{i}{2} \int_x^{2\pi} \exp\left(\frac{i}{2}(x-t)\right) f(t) dt \\ &= \frac{i}{2} \exp\left(i \frac{x}{2}\right) \int_0^x \exp\left(-i \frac{t}{2}\right) f(t) dt - \frac{i}{2} \exp\left(i \frac{x}{2}\right) \int_x^{2\pi} \exp\left(-i \frac{t}{2}\right) f(t) dt \\ &= \frac{i}{2} \exp\left(i \frac{x}{2}\right) \left\{ \int_0^x \exp\left(-i \frac{t}{2}\right) f(t) dt + \int_{2\pi}^x \exp\left(-i \frac{t}{2}\right) f(t) dt \right\}. \end{aligned}$$

Applying the Cauchy-Schwarz inequality over $[x, x + \Delta x]$ we get

$$\left| \int_x^{x+\Delta x} \exp\left(-i \frac{t}{2}\right) f(t) dt \right| \leq \|f\|_2 \cdot \sqrt{\Delta x},$$

where obviously the latter factor in the expression for $Kf(x)$ is continuous. The former factor is also continuous, so $K : H \rightarrow F$ is a mapping of H into F .

Then we get the estimate

$$\begin{aligned} |Kf(x)| &\leq \frac{1}{2} \cdot 1 \cdot \left\{ \int_0^x 1 \cdot |f(t)| dt + \int_x^{2\pi} 1 \cdot |f(t)| dt \right\} \\ &\leq \frac{1}{2} \|f\|_2 \{ \sqrt{x} + \sqrt{2\pi - x} \} \leq \frac{1}{2} \|f\|_2 \cdot \{ \sqrt{\pi} + \sqrt{\pi} \} = \sqrt{\pi} \cdot \|f\|_2, \end{aligned}$$

because $\sqrt{x} + \sqrt{2\pi - x}$ has its maximum in the interval $[0, 2\pi]$ at $x = \pi$. Then

$$\|Kf\|_\infty \leq \sqrt{\pi} \cdot \|f\|_2, \quad \text{hence} \quad \|K\| \leq \sqrt{\pi},$$

and the linear operator $K : H \rightarrow F$ is continuous.

3) Assume that $f \in F$ with $Kf \equiv 0$. Then by (2),

$$\int_0^x \exp\left(-i \frac{t}{2}\right) f(t) dt + \int_{2\pi}^x \exp\left(-i \frac{t}{2}\right) f(t) dt = 0,$$

for all $x \in [0, 2\pi]$. Both integrands are continuous, and the sum of the integrals are C^1 and constant, hence by differentiation,

$$0 = \exp\left(-i \frac{x}{2}\right) f(x) + \exp\left(-i \frac{x}{2}\right) f(x) = 2 \exp\left(-i \frac{x}{2}\right) f(x),$$

and we get $f \equiv 0$, so $S = K|_F$ is injective.

It was mentioned above that $Kf \in C^1$, if $f \in C$. Furthermore,

$$Kf(0) = \frac{i}{2} \cdot 1 \left\{ 0 - \int_0^{2\pi} \exp\left(-i \frac{t}{2}\right) f(t) dt \right\} = -\frac{i}{2} \int_0^{2\pi} \exp\left(-i \frac{t}{2}\right) f(t) dt,$$

and

$$\begin{aligned} Kf(2\pi) &= \frac{i}{2} \exp\left(i \cdot \frac{2\pi}{2}\right) \left\{ \int_0^{2\pi} \exp\left(-i \frac{t}{2}\right) f(t) dt + 0 \right\} \\ &= -\frac{i}{2} \int_0^{2\pi} \exp\left(-i \frac{t}{2}\right) f(t) dt = Kf(0), \end{aligned}$$

so we infer that

$$D(S^{-1}) = KF \subseteq \{g \in C^1([0, 2\pi]) \mid g(0) = g(2\pi)\}.$$

If on the other hand $g \in C^1([0, 2\pi])$ satisfies $g(0) = g(2\pi)$, then we shall check if the equation

$$Kf(x) = \frac{i}{2} \exp\left(i \frac{x}{2}\right) \left\{ \int_0^x \exp\left(-i \frac{t}{2}\right) f(t) dt + \int_{2\pi}^x \exp\left(-i \frac{t}{2}\right) f(t) dt \right\} = g(x)$$

has a solution $f \in F$. This equation is equivalent to

$$(8) \quad \int_0^x \exp\left(-i \frac{t}{2}\right) f(t) dt + \int_{2\pi}^x \exp\left(-i \frac{t}{2}\right) f(t) dt = -2i \exp\left(-i \frac{x}{2}\right) g(x),$$

so we get by differentiation,

$$(9) \quad 2 \exp\left(-i \frac{x}{2}\right) f(x) = -2i \exp\left(-i \frac{x}{2}\right) \left\{ -\frac{i}{2} g(x) + g'(x) \right\},$$

where (9) is equivalent to that the candidate $f(x)$ must have the structure

$$f(x) = -\frac{1}{2} g(x) - i g'(x).$$

It is obvious that f given in this way is continuous, when $g \in C^1$. The proof will be concluded, if we can prove that the additional condition $g(0) = g(2\pi)$ combined with (9) implies (8). The trick is that we write

$$2 \exp\left(-i \frac{x}{2}\right) f(x) = \exp\left(-i \frac{x}{2}\right) f(x) + \exp\left(-i \frac{x}{2}\right) f(x),$$

where we integrate the former term on the right hand side from 0 to x , and the latter from 2π to x . This construction is guaranteed by the assumption $g(0) = g(2\pi)$.

ALTERNATIVELY one may compute explicitly,

$$Kf(x) = -i K(g')(x) - \frac{1}{2} K(g)(x),$$

and then convince oneself by some partial integration that the result is $g(x)$. \diamond

4) The equation $S^{-1}g(x) = \lambda g(x)$ for $g \in D(S^{-1})$ is rewritten as

$$-i g'(x) - \frac{1}{2} g(x) = \lambda g(x), \quad g(0) = g(2\pi), \quad g \in C^1([0, 2\pi]),$$

i.e.

$$g'(x) = i \left\{ \lambda + \frac{1}{2} \right\} g(x), \quad g(0) = g(2\pi).$$

Please click the advert



What do you want to do?

No matter what you want out of your future career, an employer with a broad range of operations in a load of countries will always be the ticket. Working within the Volvo Group means more than 100,000 friends and colleagues in more than 185 countries all over the world. We offer graduates great career opportunities – check out the Career section at our web site www.volvogroup.com. We look forward to getting to know you!

VOLVO
AB Volvo (publ)
www.volvogroup.com

VOLVO TRUCKS | RENAULT TRUCKS | MACK TRUCKS | VOLVO BUSES | VOLVO CONSTRUCTION EQUIPMENT | VOLVO PENTA | VOLVO AERO | VOLVO IT
VOLVO FINANCIAL SERVICES | VOLVO 3P | VOLVO POWERTRAIN | VOLVO PARTS | VOLVO TECHNOLOGY | VOLVO LOGISTICS | BUSINESS AREA ASIA

The complete solution without the boundary condition is

$$g(x) = c \cdot \exp \left(i \left(\lambda + \frac{1}{2} \right) x \right).$$

Choosing $c = 1$ and inserting into the boundary condition, we get

$$\exp \left(i \left(\lambda + \frac{1}{2} \right) 0 \right) = 1 = \exp \left(i \left(\lambda + \frac{1}{2} \right) \cdot 2\pi \right),$$

the solutions of which are $\lambda_n + \frac{1}{2} = n \in \mathbb{Z}$.

The eigenvalues are

$$\sigma_p(S^{-1}) = \left\{ \lambda_n = n - \frac{1}{2} \mid n \in \mathbb{Z} \right\},$$

with the corresponding normalized eigenfunctions

$$e_n(x) = \frac{1}{\sqrt{2\pi}} e^{in\pi}, \quad n \in \mathbb{Z}.$$

5) It follows from $S^{-1}e_n(x) = \lambda_n e_n(x)$ that

$$\lambda_n K e_n(x) = e_n(x), \quad \text{thus} \quad K e_n(x) = \frac{1}{\lambda_n} e_n(x),$$

and K has the same eigenfunctions as S^{-1} , and the corresponding eigenvalues are

$$\left\{ \frac{1}{\lambda_n} = \frac{1}{n - \frac{1}{2}} = \frac{2}{2n - 1} \mid n \in \mathbb{Z} \right\} \subseteq \sigma_p(K).$$

Using that K is a self adjoint Hilbert-Schmidt operator, we get that the spectrum is given by

$$\sigma(K) = \{0\} \cup \left\{ \frac{2}{2n - 1} \mid n \in \mathbb{Z} \right\},$$

where each $\frac{2}{2n - 1}$ is an eigenvalue. Now, K is injective according to (3), so 0 is not an eigenvalue, thus

$$\sigma_c(K) = \{0\} \quad \text{and} \quad \sigma_p(K) = \left\{ \frac{2}{2n - 1} \mid n \in \mathbb{Z} \right\}.$$

Finally,

$$k(x, t) = \sum_{n=-\infty}^{+\infty} \frac{1}{\lambda_n} e_n(x) \cdot \overline{e_n(t)} = \frac{1}{\pi} \sum_{n=-\infty}^{+\infty} \frac{1}{2n - 1} e^{in(x-t)}.$$


6) Let $f \in H$ be given by the Fourier expansion

$$f = \sum_{n=-\infty}^{+\infty} c_n e^{inx}.$$

Since e^{inx} is an eigenfunction for K corresponding to the eigenvalue $\frac{1}{\lambda_n} = \frac{2}{2n-1}$, it follows by a termwise application of K that

$$Kf = \sum_{n=-\infty}^{+\infty} c_n K(e^{inx}) = \sum_{n=-\infty}^{+\infty} \frac{2}{2n-1} c_n e^{inx}.$$

Please click the advert



Do you want your Dream Job?


More customers get their dream job by using RedStarResume than any other resume service.

RedStarResume can help you with your job application and CV.

Go to: Redstarresume.com

Use code "BOOKBOON" and save up to \$15

(enter the discount code in the "Discount Code Box")



2 Other types of integral operators

Example 2.1 We shall consider $H = L^2([0, 1])$ as a real Hilbert space, and define $T : H \rightarrow H$ by

$$Tf(x) = \int_0^x f(t) dt.$$

Show that

$$|Tf(x)| \leq \sqrt{x} \|f\|_2,$$

and use this to show that $\|T\| < 1$.

Show that

$$T^n f(x) = \int_0^x \frac{(x-t)^{n-1}}{(n-1)!} f(t) dt.$$

Show that $\log(I+T)$ is a well-defined operator of Volterra type, and find an explicit expression for the kernel of this operator, using only known functions, that is, find k such that

$$\log(I+T)f(x) = \int_0^x k(x, t) f(t) dt.$$

1) It follows from the Cauchy-Schwarz inequality that

$$\begin{aligned} |Tf(x)| &= \left| \int_0^x f(t) dt \right| = \left| \int_0^1 1_{[0,x]}(t) f(t) dt \right| \leq \|1_{[0,x]}\|_2 \|f\|_2 \\ &= \left(\int_0^1 \{1_{[0,x]}(t)\}^2 dt \right)^{\frac{1}{2}} \|f\|_2 = \left\{ \int_0^x dt \right\}^{\frac{1}{2}} \|f\|_2 = \sqrt{x} \cdot \|f\|_2. \end{aligned}$$

(There are more variants of this computation).

2) It follows from the estimate above that

$$\|Tf\|_2^2 = \int_0^1 |Tf(x)|^2 dx \leq \int_0^1 x \|f\|_2^2 dx = \left[\frac{x^2}{2} \right]_0^1 \|f\|_2^2 = \frac{1}{2} \|f\|_2^2,$$

and we conclude that

$$\|T\| \leq \frac{1}{\sqrt{2}} < 1.$$

3) The formula clearly holds for $n = 1$. Assume that for some $n \in \mathbb{N}$,

$$T^n f(x) = \int_0^x \frac{(x-t)^{n-1}}{(n-1)!} f(t) dt, \quad f \in L^2([0, 1]).$$

Interchanging the order of integration in the computation below we get

$$\begin{aligned} T^{n+1}f(x) &= T^n(Tf)(x) = \int_0^x \frac{(x-t)^{n-1}}{(n-1)!} Tf(t) dt = \int_0^x \frac{(x-t)^{n-1}}{(n-1)!} \int_0^t f(s) ds dt \\ &= \int_0^x \left\{ \int_s^x \frac{(x-t)^{n-1}}{(n-1)!} dt \right\} f(s) ds = \int_0^x \left[-\frac{(x-t)^n}{n!} \right]_{t=s}^{t=x} f(s) ds \\ &= \int_0^x \frac{(x-s)^n}{n!} f(s) ds, \end{aligned}$$

and it follows that the formula also holds, when n is replaced by $n + 1$. Then the claim follows by induction.

4) Now,

$$\varphi(\lambda) = \log(1 + \lambda) = \sum_{n=1}^{+\infty} (-1)^{n-1} \frac{1}{n} \lambda^n, \quad \text{for } |\lambda| < 1,$$

and $T \in B(L^2([0, 1]))$ with $\|T\| \leq \frac{1}{\sqrt{2}} < 1$, so the operator $\log(I + T)$ is indeed defined by

$$\varphi(T) = \log(I + T) = \sum_{n=1}^{+\infty} (-1)^{n-1} \frac{1}{n} T^n.$$

Each of the T^n is of Volterra type, and $\varphi(T)$ contains only T^n for $n \geq 1$, hence $\varphi(T)$ is also of Volterra type.

5) When we insert the expression for $T^n f$ from (3), we get by purely formal computations that

$$\log(I + T)f(x) = \sum_{n=1}^{+\infty} (-1)^{n-1} \frac{1}{n} \int_0^x \frac{(x-t)^{n-1}}{(n-1)!} f(t) dt = \sum_{n=1}^{+\infty} \int_0^x \frac{(t-x)^{n-1}}{n!} f(t) dt.$$

However, the series $\sum_{n=1}^{+\infty} \frac{(t-x)^{n-1}}{n!}$ is *uniformly* convergent for $0 \leq t \leq x \leq 1$. (Notice that we get the sum 1 for $t = x$). Therefore it is indeed legal to interchange summation and integration. The we get for $0 \leq t < x$ the sum

$$\sum_{n=1}^{+\infty} \frac{(t-x)^{n-1}}{n!} = \frac{1}{t-x} \left\{ \sum_{n=0}^{+\infty} \frac{(t-x)^n}{n!} - 1 \right\} = \frac{e^{t-x} - 1}{t-x} = e^{-x} \cdot \frac{e^x - e^t}{x-t}.$$

Note that we for $t \rightarrow x$ get the limit $e^{-x} \cdot e^x = 1$.

We get by interchanging summation and integration,

$$\log(I + T)f(x) = \int_0^x e^{-x} \cdot \frac{e^x - e^t}{x-t} f(t) dt,$$

so the kernel of the Volterra operator $\log(I + T)$ is given by

$$k(x, t) = \begin{cases} e^{-x} \cdot \frac{e^x - e^t}{x-t} & \text{for } 0 \leq t < x \leq 1, \\ 1 & \text{for } 0 \leq t = x \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Example 2.2 In this example it is allowed to change the order of integrations without justification. Consider the operator

$$Af(x) = \frac{1}{\sqrt{\pi}} \int_0^x \frac{f(t)}{\sqrt{x-t}} dt, \quad x \in [0, 1],$$

whenever this expression gives sense.

- 1) Show that $Af \in L^\infty([0, 1])$ if $f \in L^p([0, 1])$, $p > 2$.
- 2) Find the operator $B = A^2$, that is find the kernel $k(x, t)$ such that

$$Bf(x) = A^2 f(x) = \int_0^x k(x, t) f(t) dt$$

for $f \in L^p([0, 1])$, $p > 2$.

- 3) Show that $B : L^p([0, 1]) \rightarrow L^\infty([0, 1])$, $1 \leq p \leq \infty$ is bounded.
- 4) Solve the equation

$$(I - A)f(x) = 1$$

formally by a Neumann series, and express f as

$$f(x) = g(x) + Ah(x),$$

where g and h are known functions. (Here it is not possible to express $Ah(x)$ as a known function.) Insert and show that this formal solution is a solution.

Remark 2.1 First note that the kernel does *not* belong to $L^2([0, 1]^2)$. In fact, it follows from

$$k(x, t) = \begin{cases} \frac{1}{\sqrt{x-t}} & \text{for } 0 \leq t < x \leq 1, \\ 0 & \text{otherwise,} \end{cases}$$

that

$$\int_0^1 \int_0^1 |k(x, t)|^2 dt dx = \int_0^1 \left\{ \int_0^x \frac{dt}{x-t} \right\} dx = \int_0^1 [-\ln(x-t)]_{t=0}^x dx = +\infty,$$

so we cannot apply the theory of the Hilbert-Schmidt operators. Part of the example is to use other methods. \diamond

- 1) Given $f \in L^p([0, 1])$, where $p > 2$, thus $1 < q < 2$, where q is the conjugated number of p , i.e. $\frac{1}{p} + \frac{1}{q} = 1$. Then by the Hölder inequality

$$\begin{aligned} |Af(x)| &\leq \frac{1}{\sqrt{\pi}} \int_0^x \frac{|f(t)|}{\sqrt{x-t}} dt \leq \frac{1}{\sqrt{\pi}} \left\{ \int_0^x |f(t)|^p dt \right\}^{\frac{1}{p}} \left\{ \int_0^x \frac{dt}{(x-t)^{q/2}} \right\}^{\frac{1}{q}} \\ &\leq \frac{1}{\sqrt{\pi}} \|f\|_p \left\{ \frac{-1}{1-\frac{q}{2}} \left[(x-t)^{1-\frac{q}{2}} \right]_{t=0}^x \right\}^{\frac{1}{q}} = \frac{1}{\sqrt{\pi}} \|f\|_p \left\{ \frac{1}{1-\frac{q}{2}} x^{1-\frac{q}{2}} \right\}^{\frac{1}{q}} \\ &\leq \frac{1}{\sqrt{\pi}} \cdot \left\{ 1 - \frac{q}{2} \right\}^{-\frac{1}{q}} \|f\|_p, \end{aligned}$$

where we have used that $1 - \frac{q}{2} > 0$, because $p > 2$. This holds for all $x \in [0, 1]$, so

$$\|Af\|_{\infty} \leq \frac{1}{\sqrt{\pi}} \cdot \left\{1 - \frac{q}{2}\right\}^{-\frac{1}{q}} \|f\|_p,$$

and $Af \in L^{\infty}([0, 1])$ for $f \in L^p([0, 1])$, when $2 < p < +\infty$.

If instead $p = +\infty$, then we get the following estimate,

$$\begin{aligned} |Af(x)| &\leq \frac{1}{\sqrt{\pi}} \int_0^x \frac{|f(t)|}{\sqrt{x-t}} dt = \frac{1}{\sqrt{\pi}} \|f\|_{\infty} \int_0^x \frac{dt}{\sqrt{x-t}} \\ &= \frac{1}{\sqrt{\pi}} \|f\|_{\infty} \cdot \left[\frac{-1}{1-\frac{1}{2}} \sqrt{x-t} \right]_0^x = \frac{2}{\sqrt{\pi}} \sqrt{x} \cdot \|f\|_{\infty} \leq \frac{2}{\sqrt{\pi}} \|f\|_{\infty}, \end{aligned}$$

and we get in this case that

$$\|Af\|_{\infty} \leq \frac{2}{\sqrt{\pi}} \|f\|_{\infty},$$

hence $Af \in L^{\infty}([0, 1])$ for $f \in L^{\infty}([0, 1])$.

- 2) Assume again that $f \in L^p([0, 1])$, where $p > 2$. Then $Af \in L^{\infty}([0, 1])$ according to (1). From $p_1 = \infty > 2$ follows by another application of (1) that $A^2 f \in L^{\infty}([0, 1])$.

Try this...



The sequence 2, 4, 6, 8, 10, 12, 14, 16, ... is the sequence of even whole numbers. The 100th place in this sequence is the number...?

Challenging? Not challenging? Try more >>

www.alloptions.nl/life

Compute

$$Bf(x) = A^2 f(x) = \frac{1}{\sqrt{\pi}} \int_0^x \frac{1}{\sqrt{x-t}} Af(t) dt = \frac{1}{\sqrt{\pi}} \int_0^x \frac{1}{\sqrt{x-t}} \left\{ \frac{1}{\sqrt{\pi}} \int_0^t \frac{f(u)}{\sqrt{t-u}} du \right\} dt.$$

From $0 \leq u \leq t \leq x \leq 1$ we infer by an interchange of the integrals as follows by the change of variable $xs = t - u$ that

$$\begin{aligned} Bf(x) &= \frac{1}{\pi} \int_0^x \left\{ \int_u^x \frac{dt}{\sqrt{(x-t)(t-u)}} \right\} f(u) du = \frac{1}{\pi} \int_0^x \left\{ \int_0^{x-u} \frac{ds}{\sqrt{\{(x-u)-s\}s}} \right\} f(u) du \\ &= \frac{1}{\pi} \int_0^x \pi f(u) du = \int_0^x f(t) dt, \end{aligned}$$

where we have used that

$$\int_0^a \frac{ds}{\sqrt{(a-s)s}} = \pi \quad \text{for } a = x - u > 0.$$

Remark 2.2 We prove for completeness this formula. We get by the monotonous substitution $s = a \sin^2 \theta$, $\theta \in [0, \frac{\pi}{2}]$,

$$\begin{aligned} \int_0^a \frac{ds}{\sqrt{(a-s)s}} &= \int_0^{\frac{\pi}{2}} \frac{1 \cdot 2 \sin \theta \cos \theta}{\sqrt{(a - a \sin^2 \theta) \cdot a \sin^2 \theta}} d\theta = 2a \int_0^{\frac{\pi}{2}} \frac{\sin \theta \cos \theta}{\sqrt{a^2(1 - \sin^2 \theta) \sin^2 \theta}} d\theta \\ &= \frac{2a}{|a|} \int_0^{\frac{\pi}{2}} \frac{\cos \theta \sin \theta}{|\cos \theta \sin \theta|} d\theta = 2 \int_0^{\frac{\pi}{2}} d\theta = \pi. \quad \diamond \end{aligned}$$

The operator is therefore a well-known integral operator, and A corresponds to “integrating one half time from 0”. The kernel is explicitly given by

$$k(x, t) = \begin{cases} 1 & \text{for } 0 \leq t \leq x \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

3) This follows easily from the Hölder inequality,

$$|Bf(x)| \leq \int_0^x |f(t)| dt \leq \int_0^1 |f(t)| \cdot 1 dt \leq 1 \cdot \|f\|_p,$$

hence $\|Bf\|_\infty \leq \|f\|_p$, and $\|B\| \leq 1$.

4) The Neumann series is given by

$$(I - A)^{-1} = \sum_{n=0}^{+\infty} A^n,$$

so the formal solution is

$$\begin{aligned} f(x) &= \sum_{n=0}^{+\infty} A^n 1(x) = \sum_{n=0}^{+\infty} A^{2n} 1(x) + \sum_{n=0}^{+\infty} A^{2n+1} 1(x) \\ &= \sum_{n=0}^{+\infty} B^n 1(x) + A \sum_{n=0}^{+\infty} B^n 1(x) = g(x) + Ag(x), \end{aligned}$$

hence

$$\begin{aligned} h(x) = g(x) &= \sum_{n=0}^{+\infty} B^n 1(x) = 1 + \sum_{n=1}^{+\infty} B^n 1(x) = 1 + \sum_{n=1}^{+\infty} \int_0^x \frac{t^{n-1}}{(n-1)!} \cdot 1 \, dt \\ &= 1 + \sum_{n=1}^{+\infty} \frac{x^n}{n!} = e^x, \end{aligned}$$

and the formal solution is


$$f(x) = e^x + Ae^x.$$

Then we get by insertion

$$\begin{aligned} (I - A)f(f) &= f(x) - Af(f) = e^x + Ae^x - Ae^x - A^2e^x \\ &= e^x - Be^x = e^x - \int_0^x e^t \, dt = e^x - [e^t]_0^x = e^x - (e^x - 1) = 1, \end{aligned}$$

and we have proved that we have found a solution.

Please click the advert



ALTERNATIVELY (and more elegantly),

$$(I - A)(I + A) = (I + A)(I - A) = I - A^2 = I - B.$$

Since B is a Volterra operator, we have that $(I - B)^{-1} = \sum_{n=0}^{+\infty} B^n$ is bounded. Clearly, A and $B = A^2$ commutes, so

$$(I - A) \{ (I + A)(I - B)^{-1} \} = \{ (I + A)(I - B)^{-1} \} (I - A) = I,$$

proving that

$$(I - A)^{-1} = (I + A)(I - B)^{-1}.$$

Hence the equation $(I - A)f = 1$ is equivalent to

$$f(x) = (I - A)^{-1}A(x) = (I + A) \sum_{n=0}^{+\infty} B^n 1(x) = (I + A)e^x = e^x + Ae^x,$$

where we have applied the computation above.

Example 2.3 Let $H = L^2([0, 1])$ and consider the integral operator

$$Bf(x) = \int_0^x f(t) dt, \quad \text{for } f \in H.$$

1) Show that

$$k(x, t) = \min\{x, t\}, \quad 0 \leq x, t \leq 1,$$

is the kernel for the self adjoint Hilbert-Schmidt operator $K = BB^*$.

2) Let φ be an eigenfunction for K associated with a non-zero eigenvalue λ . Justify that φ can be taken as a C^∞ -function.

Next, show that φ must satisfy the equation

$$\lambda \varphi''(x) = -\varphi(x),$$

and use this to find all non-zero eigenvalues for K and all the associated eigenfunctions.

3) Assuming the $\|BB^*\| = \|B^*\|^2$, show that $\|K\| = \|B\|^2$, and find both $\|K\|$ and $\|B\|$.

1) The operator B has the kernel

$$b(x, t) = \begin{cases} 1 & \text{for } 0 \leq t \leq x \leq 1, \\ 0 & \text{otherwise,} \end{cases}$$

so

$$b^*(x, t) = \overline{b(t, x)} = b(t, x) = \begin{cases} 1 & \text{for } 0 \leq x \leq t \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Then the kernel $k(x, t)$ for $K = BB^*$ is given by

$$\begin{aligned} k(x, t) &= \int_0^1 b(x, s)b^*(s, t) ds = \int_0^1 b(x, s)b(t, s) ds \\ &= \int_0^1 b(\min\{x, t\}, s) ds = \min\{x, t\}, \quad x, t \in [0, 1]. \end{aligned}$$

- 2) Since $k(x, t)$ is continuous, we can choose the eigenfunctions continuous. Hence, if $\varphi(x)$ is an eigenfunction corresponding to an eigenvalue $\lambda \neq 0$, then

$$(10) \quad \lambda \varphi(x) = \int_0^1 k(x, t) \varphi(t) dt = \int_0^x t \varphi(t) dt + x \int_x^1 \varphi(t) dt.$$

If φ is continuous, then the right hand side of (10) is differentiable. If φ is of class C^n , then the right hand side of (10) is of class C^{n+1} , hence φ is also of class C^{n+1} . Then the claim follows by induction, hence $\varphi \in C^\infty$.

When we differentiate (10), we get

$$\lambda \varphi'(x) = x \varphi(x) + \int_x^1 \varphi(t) dt - x \varphi(x) = \int_x^1 \varphi(t) dt,$$

hence by another differentiation,

$$(11) \quad \lambda \varphi''(x) = -\varphi(x),$$

and the claim is proved.

- 3) Let $\alpha \in \mathbb{C} \setminus \{0\}$ satisfy the condition $\alpha^2 = \frac{1}{\lambda}$. Then the equation (11) has the complete solution

$$(12) \quad \varphi(x) = C_1 e^{i\alpha x} + C_2 e^{-i\alpha x}.$$

When (12) is put into (10), and we apply that $\frac{1}{\alpha^2} = \lambda$, then

$$\begin{aligned} \lambda \varphi(x) &= \lambda \{C_1 e^{i\alpha x} + C_2 e^{-i\alpha x}\} \\ &= \int_0^x t \{C_1 e^{i\alpha t} + C_2 e^{-i\alpha t}\} dt + x \int_x^1 \{C_1 e^{i\alpha t} + C_2 e^{-i\alpha t}\} dt \\ &= \left[t \left\{ \frac{C_1}{i\alpha} e^{i\alpha t} - \frac{C_2}{i\alpha} e^{-i\alpha t} \right\} \right]_0^x - \int_0^x \left\{ \frac{C_1}{i\alpha} e^{i\alpha t} - \frac{C_2}{i\alpha} e^{-i\alpha t} \right\} dt \\ &\quad + x \left[\frac{C_1}{i\alpha} e^{i\alpha t} - \frac{C_2}{i\alpha} e^{-i\alpha t} \right]_x^1 \\ &= x \left\{ \frac{C_1}{i\alpha} e^{i\alpha x} - \frac{C_2}{i\alpha} e^{-i\alpha x} \right\} - \left[\frac{C_1}{i^2 \alpha^2} e^{i\alpha t} + \frac{C_2}{i^2 \alpha^2} e^{-i\alpha t} \right]_0^x \\ &\quad + x \left\{ \frac{C_1}{i\alpha} e^{i\alpha} - \frac{C_2}{i\alpha} e^{-i\alpha} \right\} - x \left\{ \frac{C_1}{i\alpha} e^{i\alpha x} - \frac{C_2}{i\alpha} e^{-i\alpha x} \right\} \\ &= \frac{1}{\alpha^2} \{C_1 e^{i\alpha x} + C_2 e^{-i\alpha x}\} - \frac{1}{\alpha^2} \{C_1 + C_2\} + \frac{x}{i\alpha} \{C_1 e^{i\alpha} - C_2 e^{-i\alpha}\} \\ &= \lambda \varphi(x) - \lambda \{C_1 + C_2\} + \frac{x}{i\alpha} \{C_1 e^{i\alpha} - C_2 e^{-i\alpha}\}. \end{aligned}$$

This equation holds for every x , and $\lambda \neq 0$ and $\alpha \neq 0$, so we conclude that

$$C_1 + C_2 = 0 \quad \text{and} \quad C_1 e^{i\alpha} - C_2 e^{-i\alpha} = 0,$$

hence $C_2 = -C_1$, and $C_1 \{e^{i\alpha} + e^{-i\alpha}\} = 2C_1 \cos \alpha = 0$, thus

$$\alpha = \frac{\pi}{2} + n\pi, \quad n \in \mathbb{Z}.$$

It follows from

$$\varphi(x) = C_1 e^{i\alpha x} + C_2 e^{-i\alpha x} = C_1 \{e^{i\alpha x} - e^{-i\alpha x}\} = 2i C_1 \sin \alpha x,$$

that the eigenfunctions for K corresponding to a $\lambda \in \sigma_p(K) \setminus \{0\}$ are some constant times

$$\varphi_n(x) = \sin \left(\left(n - \frac{1}{2} \right) \pi x \right), \quad n \in \mathbb{N},$$

corresponding to the eigenvalue

$$\lambda_n = \frac{1}{\alpha_n^2} = \frac{4}{\pi^2} \cdot \frac{1}{(2n+1)^2}, \quad n \in \mathbb{N}.$$


4) Now, $\|K\|$ is the absolute value of the numerically largest eigenvalue $|\lambda_1|$, so

$$\|K\| = \|BB^*\| = \lambda_1 = \frac{4}{\pi^2} \cdot \frac{1}{(2-1)^2} = \left(\frac{2}{\pi} \right)^2.$$

On the other hand, BB^* is self adjoint, hence

$$\begin{aligned} \|BB^*\| &= \sup\{|(BB^*f, f)| \mid f \in L^2([0, 1]), \|f\|_2 = 1\} \\ &= \sup\{(B^*f, B^*f) \mid f \in L^2([0, 1]), \|f\|_2 = 1\} \\ &= \sup\{\|B^*f\|^2 \mid f \in L^2([0, 1]), \|f\|_2 = 1\} = \|B^*\|^2. \end{aligned}$$

Please click the advert



Masters in Management



Designed for high-achieving graduates across all disciplines, London Business School's Masters in Management provides specific and tangible foundations for a successful career in business.

This 12-month, full-time programme is a business qualification with impact. In 2010, our MiM employment rate was 95% within 3 months of graduation*; the majority of graduates choosing to work in consulting or financial services.

As well as a renowned qualification from a world-class business school, you also gain access to the School's network of more than 34,000 global alumni – a community that offers support and opportunities throughout your career.

For more information visit www.london.edu/mm, email mim@london.edu or give us a call on **+44 (0)20 7000 7573**.

* Figures taken from London Business School's Masters in Management 2010 employment report

Finally, $B \in B(H)$, hence also $B^* \in B(H)$ with $\|B^*\| = \|B\|$, and whence

$$\|K\| = \|BB^*\| = \|B^*\|^2 = \|B\|^2 = \left(\frac{2}{\pi}\right)^2.$$

Then

$$\|B\| = \frac{2}{\pi},$$

where

$$Bf(x) = \int_0^x f(t) dt, \quad f \in L^2([0, 1]).$$

Example 2.4 Let $H = L^2([0, 1])$ and consider the operator K with domain $D(K) = C([0, 1])$ given by

$$Kf(x) = x \int_0^x f(t) dt + \int_x^1 t f(t) dt, \quad f \in D(K).$$

1) Show that $K : D(K) \rightarrow C^2([0, 1])$, and that

$$(Kf)'(0) = 0 \quad \text{and} \quad (Kf)'(1) = (Kf)(1).$$

2) Show that K is injective and that K^{-1} has the domain

$$D(K^{-1}) = \{u \in C^2([0, 1]) \mid u'(0) = 0, u(1) = u'(1)\},$$

and the action $K^{-1}u = u''$.

3) Show that K is an integral operator with continuous and symmetric kernel and find this kernel.

4) Let φ and ψ denote eigenfunctions for K associated to the same eigenvalue λ . Define the function f by

$$f(x) = \psi(0)\varphi(x) - \varphi(0)\psi(x),$$

and use the existence and uniqueness theorem for ordinary differential equations to argue that $f = 0$.

Next show that all eigenspaces for K are of dimension one.

5) Let $\sigma_p(K) = (\lambda_n)$ denote the sequence of eigenvalues for K . Find

$$\sum_{n=1}^{\infty} \lambda_n^2.$$

6) Let λ be a positive eigenvalue and let $\mu = \frac{1}{\sqrt{\lambda}}$. Express the associated eigenfunction with μ as a transcendent equation for μ .

Use a graph argument to show that K has at most one positive eigenvalue.

- 1) If $f \in C([0, 1])$, then we get immediately that Kf is of class $C^1([0, 1])$ and

$$(Kf)'(x) = \int_0^x f(t) dt + x f(x) - x f(x) = \int_0^x f(t) dt.$$

This shows that we even have $(Kf)' \in C^1([0, 1])$, hence $Kf \in C^2([0, 1])$, and

$$(13) \quad (Kf)''(x) = f(x).$$

Furthermore,

$$(Kf)'(0) = \int_0^0 f(t) dt = 0,$$

and

$$(Kf)(1) = 1 \cdot \int_0^1 f(t) dt + \int_1^1 t f(t) dt = \int_0^1 f(t) dt = (Kf)'(1).$$

- 2) Now, K is linear, hence K is injective, If $Kf(x) \equiv 0$ implies that $f = 0$. This follows from (13) in (1), because

$$f(x) = (Kf)''(x) = 0.$$

Assume that $u \in C^2([0, 1])$ satisfies $u'(0) = 0$ and $u(1) = u'(1)$. We shall prove that there is an $f \in C([0, 1])$, for which $u = Kf$. According to (13) the only possibility is $f = u''$, which we now check. Using that $u'' \in C([0, 1])$, we get

$$\begin{aligned} Ku''(x) &= x \int_0^x u''(t) dt + \int_x^1 t u''(t) dt = x \{u'(x) - u'(0)\} + [t u'(t)]_x^1 - \int_x^1 1 \cdot u'(t) dt \\ &= x u'(x) + u'(1) - x u'(x) - [u(t)]_x^1 = u'(1) - u(1) + u(x) = u(x), \end{aligned}$$

and the claim is proved.

- 3) We get from the expression for Kf that

$$Kf(x) = \int_0^1 k(x, t) f(t) dt = \int_0^x f(t) dt + \int_x^1 t f(t) dt = \int_0^1 \max\{x, t\} f(t) dt,$$

thus

$$k(x, t) = \max\{x, t\} \quad \text{for } x, t \in [0, 1],$$

and $k(x, t)$ is clearly continuous in $[0, 1]^2$, hence of class $L^2([0, 1]^2)$.

We note that $k(x, t) = \overline{k(t, x)}$, hence the kernel is Hermitian and K is a self adjoint Hilbert-Schmidt operator.

- 4) This is trivial. We know that K is injective, so $0 \notin \sigma_p(K)$, and if $\lambda \in \sigma_p(K)$, $\lambda \neq 0$, and $K\varphi = \lambda\varphi$, it follows by an application of K^{-1} that

$$\varphi = \lambda K^{-1}\varphi, \quad \text{i.e.} \quad K^{-1}\varphi = \frac{1}{\lambda}\varphi.$$

5) Assume that φ and ψ are eigenvectors for K with the same eigenvalue λ . Then

$$f(x) = \psi(0) \varphi(x) - \varphi(0) \psi(x)$$

is also an eigenfunction corresponding to λ , hence f is according to (4) an eigenvector corresponding to the operator $K^{-1} = \frac{d^2}{dx^2}$ with the eigenvalue $\frac{1}{\lambda}$, so

$$f''(x) = \frac{1}{\lambda} f(x).$$

Now, $(K\varphi)'(0) = 0 = \lambda\varphi'(0)$, and analogously for ψ , so we conclude from (1) that

$$f(0) = \psi(0) \varphi(0) - \varphi(0) \psi(0) = 0$$

and

$$f'(0) = \psi(0) \varphi'(0) - \varphi(0) \psi'(0) = 0.$$

It follows from the existence and uniqueness theorem for linear second order differential equations that

$$(14) \quad \frac{d^2 f}{dx^2} - \frac{1}{\lambda} f(x) = 0, \quad f(0) = 0, \quad f'(0) = 0,$$

does only have the solution $f(x) \equiv 0$, hence

$$(15) \quad \psi(0) \varphi(x) = \varphi(0) \psi(x).$$

Then assume that $\varphi(0) = 0$ for every eigenfunction. Then also $\varphi'(0) = 0$, cf. the above, so φ is a solution of (14), and $\varphi \equiv 0$. This means that φ is not an eigenfunction, contradicting the assumption. Therefore, we conclude that $\varphi(0) \neq 0$ for every eigenfunction. Then it follows from (15) that all eigenfunctions of the same eigenvalue are mutually proportional, hence every eigenspace for K has dimension 1.

6) When we use that K is self adjoint and of Hilbert-Schmidt type, cf. (3), we get that all eigenvalues are real, and

$$\sum_{n=1}^{+\infty} \lambda_n^2 = \|k\|_2^2,$$

where we have used (5) that every eigenspace has dimension 1. Then

$$\begin{aligned} \sum_{n=1}^{+\infty} \lambda_n^2 &= \|k\|_2^2 = \int_0^1 \int_0^1 \max\{x, t\}^2 dt dx = \int_0^1 \left\{ \int_0^x x^2 dt + \int_x^1 t^2 dt \right\} dx \\ &= \int_0^1 \left\{ x^3 + \left[\frac{t^3}{3} \right]_x^1 \right\} dx = \int_0^1 \left\{ x^3 + \frac{1}{3} - \frac{x^3}{3} \right\} dx = \frac{1}{3} \int_0^1 (2x^3 + 1) dx \\ &= \frac{1}{3} \left[\frac{x^4}{2} + x \right]_0^1 = \frac{1}{3} \left\{ \frac{1}{2} + 1 \right\} = \frac{1}{2}. \end{aligned}$$

7) It follows from (4) that if $\lambda > 0$ and $\mu = \frac{1}{\sqrt{\lambda}}$, then

$$\varphi''(x) = \frac{1}{\lambda} \varphi(x) = \mu^2 \varphi(x),$$

the complete solution of which is

$$\varphi(x) = C_1 e^{\mu x} + C_2 e^{-\mu x}.$$




Figure 2: The graphs of $x = \mu$ and $x = \coth \mu$ intersect at $\mu \approx 1.199678640$.

Please click the advert



You're full of *energy*
and *ideas*. And that's
just what we are looking for.

Looking for a career where your ideas could really make a difference? UBS's Graduate Programme and internships are a chance for you to experience for yourself what it's like to be part of a global team that rewards your input and believes in succeeding together.

Wherever you are in your academic career, make your future a part of ours by visiting www.ubs.com/graduates.

www.ubs.com/graduates



© UBS 2010. All rights reserved.

We shall find the values of C_1 , C_2 and μ , for which $\varphi \in D(K^{-1})$. We compute

$$\varphi'(x) = \mu \{C_1 e^{\mu x} - C_2 e^{-\mu x}\},$$

and get the conditions (because $\mu > 0$)

$$\varphi'(0) = \mu \{C_1 - C_2\} = 0, \quad \text{i.e. } C_1 = C_2 = C,$$

and

$$\varphi(1) = C \{e^\mu + e^{-\mu}\} = C\mu \{e^\mu - e^{-\mu}\} = \varphi'(1),$$

so μ is a solution of the equation

$$\cosh \mu = \mu \sinh \mu,$$

which we write as

$$\coth \mu = \mu.$$

Considering the graphs we see that this equation has only one solution $\mu > 0$.

Remark 2.3 Using the iteration

$$\mu_{n+1} = \frac{1}{\tan \mu_n}$$

we get on a pocket calculator that

$$\mu \approx 1.199\,678\,640.$$

Note that

$$\lambda_1^2 = \frac{1}{\mu^4} \approx 0.482\,770\,022 < 0.5,$$

so

$$\sum_{n=2}^{+\infty} \lambda_n^2 = 0.017\,229\,978 \ll \lambda_1^2.$$

The norm of K is approximately

$$\|K\| = \lambda_1 \approx 0.694\,82.$$

We have for any other eigenvalue $\lambda \in \mathbb{R}$ that $\lambda < 0$, so $\mu = \frac{1}{\sqrt{\lambda}}$ is purely imaginary. \diamond

Example 2.5 Let $K \in B(H)$, where $H = L^2([0, 1])$, be given by

$$Kf(x) = \int_{1-x}^1 f(t) dt.$$

- 1) Show that K is actually bounded.
- 2) Show that the kernel $k(x, t)$ for K is Hermitian, and calculate

$$\|k\|^2 = \int_0^1 \int_0^1 |k(x, t)|^2 dt dx.$$

- 3) Show that the kernel $k_2(x, t)$ for K^2 is $\min\{x, t\}$.
- 4) Show that an eigenfunction for K is an eigenfunction for K^2 .
Now, let f denote an eigenfunction for K associated with the eigenvalue λ . Calculate $(K^2 f)''$, justify that it belongs to H and show that f is a solution to the equation

$$\lambda^2 f'' + f = 0.$$

- 5) Find all eigenvalues and associated eigenfunctions for K .
- 6) Determine $\|K\|$.

- 1) Apply the Cauchy-Schwarz inequality in $L^2([1-x, 1])$ for $f \in H$. This gives

$$\|Kf\|_2^2 = \int_0^1 \left| \int_{1-x}^1 1 \cdot f(t) dt \right|^2 dx \leq \int_0^1 \{\sqrt{x} \cdot \|f\|_2\}^2 dx = \|f\|_2^2 \int_0^1 x dx = \frac{1}{2} \|f\|_2^2,$$

and we conclude that $\|K\| \leq \frac{1}{\sqrt{2}}$, thus K is bounded.

- 2) It follows from

$$Kf(x) = \int_0^1 k(x, t) f(t) dt = \int_{1-x}^2 f(t) dt = \int_0^1 1_{[1-x, 1]}(t) f(t) dt,$$

that

$$k(x, t) = 1_{[1-x, 1]}(t) = \begin{cases} 1 & \text{for } 1-x \leq t \leq 1, \\ 0 & \text{otherwise.} \end{cases} \quad x \in [0, 1],$$

Hence, $k(x, t) = 1$, if and only if $x + t \geq 1$, $x, t \in [0, 1]$, and 0 otherwise, i.e. if and only if

$$(x, t) \in B = \{(x, t) \in [0, 1]^2 \mid x + t \geq 1\},$$

so we get (cf. the figure)

$$k(x, t) = 1_B(x, t) = \overline{1_B(t, x)} = \overline{k(t, x)},$$




Figure 3: The domain B , where $k(x, t) = 1$, is the upper triangle.

which shows that the kernel is Hermitian.

Then we get

$$\|k\|_2^2 = \int_0^1 \int_0^1 |k(x, t)|^2 dt dx = \int_0^1 \int_0^1 k(x, t) dt dx = \text{area}(B) = \frac{1}{2},$$

possibly in the variant

$$\|k\|_2^2 = \int_0^1 \int_0^1 k(x, t) dt dx = \int_0^1 (K1)(x) dx = \int_0^1 \left\{ \int_{1-x}^1 dt \right\} dx = \int_0^1 x dx = \frac{1}{2}.$$

3) The kernel for K^2 is given by

$$k_2(x, t) = \int_0^1 k(x, s)k(s, t) ds,$$

where the integrand is $\neq 0$, if and only if

$$1 - x \leq s \leq 1 \quad \text{and} \quad 1 - s \leq t \leq 1.$$

This provides us with the bounds

$$1 - x \leq s \leq 1 \quad \text{and} \quad 1 - t \leq s \leq 1,$$

hence $s \leq 1$ and

$$s \geq \max\{1 - x, 1 - t\} = 1 - \min\{x, t\}.$$

Then by insertion

$$\begin{aligned} k_2(x, t) &= \int_0^1 k(x, s)k(s, t) ds = \int_{1-\min\{x, t\}}^1 k(x, s)k(s, t) ds \\ &= \int_{1-\min\{x, t\}}^1 ds = \min\{x, t\}, \end{aligned}$$

i.e.

$$k_2(x, t) = \min\{x, t\}, \quad (x, t) \in [0, 1]^2.$$

4) If $Kf = \lambda f$, then of course

$$K^2 f = \lambda Kf = \lambda^2 f,$$

so if f is an eigenfunction for K corresponding to the eigenvalue λ , then f is an eigenfunction for K^2 corresponding to the eigenvalue λ^2 .

We get, the kernel for K^2 being k_2 ,

$$K^2 f(x) = \int_0^1 \min\{x, t\} f(t) dt = \int_0^x t f(t) dt + x \int_x^1 f(t) dt.$$

Obviously, $K^2 f$ is differentiable in the weak sense, and we get

$$(K^2 f)'(x) = x f(x) + \int_x^1 f(t) dt - x f(x) = \int_x^1 f(t) dt.$$

This shows that $(K^2 f)'$ also is weakly differentiable, so

$$(K^2 f)''(x) = -f(x).$$

Please click the advert



Discover the truth at www.deloitte.ca/careers

Deloitte.

© Deloitte & Touche LLP and affiliated entities.

If f is an eigenvalue for K corresponding to the eigenvalue λ , i.e. $Kf = \lambda f$, then it follows from the above that

$$(K^2 f)(x) = \lambda^2 f(x)$$

and f is differentiable. It follows by induction that f is infinitely often differentiable, so we get from the above that

$$\lambda^2 f''(x) = (K^2 f)''(x) = -f(x),$$

hence by a rearrangement,

$$(16) \quad \lambda^2 f''(x) + f(x) = 0.$$

Therefore, if f is an eigenfunction for K with eigenvalue λ , then f must also fulfil (16). In particular, $\lambda \neq 0$, if f is an eigenfunction. It is well-known that the solutions of (16) are

$$f(x) = c_1 \exp\left(\frac{i}{\lambda} x\right) + c_2 \exp\left(-\frac{i}{\lambda} x\right).$$

From $K^2 f(0) = 0 = \lambda^2 f(0)$ follows that $f(0) = 0$, so we conclude that $c_1 + c_2 = 0$. Putting $c_1 = \frac{c}{2i}$, we get $c_2 = -\frac{c}{2i}$, and the only possibility of an eigenfunction is

$$f(x) = \frac{c}{2i} \left\{ \exp\left(\frac{i}{\lambda} x\right) - \exp\left(-\frac{i}{\lambda} x\right) \right\} = c \cdot \sin\left(\frac{x}{\lambda}\right).$$

5) It remains to find the possible eigenvalues λ .

Put $c = 1$ and $\alpha = \frac{1}{\lambda}$. It follows from $Kf(x) = \lambda f(x)$ that

$$f(x) = \sin\left(\frac{x}{\lambda}\right) = \sin(\alpha x) = \frac{1}{\lambda} Kf(x) = \alpha \cdot K \sin(\alpha \cdot)(x),$$

hence by insertion into the definition of K ,

$$\begin{aligned} \sin(\alpha x) &= \alpha \int_{1-x}^1 \sin(\alpha t) dt = [-\cos(\alpha t)]_{1-x}^1 = \cos(\alpha(1-x)) - \cos \alpha \\ &= \cos \alpha \cdot \cos \alpha x + \sin \alpha \cdot \sin \alpha x - \cos \alpha, \end{aligned}$$

so

$$(1 - \sin \alpha) \sin \alpha x = \cos \alpha \cdot (\cos \alpha x - 1).$$

This equation is fulfilled for all x , if either $\alpha = 0$, which is not possible because $\alpha = \frac{1}{\lambda}$, or if $\cos \alpha = 0$ and $\sin \alpha = 1$, hence

$$\alpha_p = \frac{\pi}{2} + 2p\pi, \quad p \in \mathbb{Z},$$

and we get

$$\lambda_p = \frac{1}{\alpha_p} = \frac{1}{\frac{\pi}{2} + 2p\pi} = \frac{1}{\pi} \cdot \frac{1}{4p+1}, \quad p \in \mathbb{Z}.$$

Then we derive the point spectrum and the continuous spectrum,

$$\sigma_p(K) = \left\{ \frac{2}{\pi} \cdot \frac{1}{4p+1} \mid p \in \mathbb{Z} \right\} \quad \text{and} \quad \sigma_c(K) = \{0\}.$$

The eigenfunction corresponding to

$$\lambda_p = \frac{2}{\pi} \cdot \frac{1}{4p+1}, \quad p \in \mathbb{Z},$$

is

$$f_p(x) = \sin \left(\left(\frac{\pi}{2} + 2p\pi \right) x \right), \quad x \in [0, 1]; \quad p \in \mathbb{Z}.$$

6) The numerically largest eigenvalue is $\lambda_0 = \frac{2}{\pi} > 0$, hence

$$\|K\| = \max\{|\lambda_p| \mid p \in \mathbb{Z}\} = \frac{2}{\pi}.$$

CHECK. As a *check* we use that we should have

$$\frac{1}{2} = \|k\|_2^2 = \sum_{p \in \mathbb{Z}} |\lambda_p|^2.$$

We get

$$\sum_{p \in \mathbb{Z}} |\lambda_p|^2 = \frac{4}{\pi^2} \sum_{p=-\infty}^{+\infty} \frac{1}{(4p+1)^2} = \frac{4}{\pi^2} \sum_{p=0}^{+\infty} \frac{1}{(2p+1)^2} = \frac{4}{\pi^2} \cdot \frac{\pi^2}{8} = \frac{1}{2} = \|k\|_2^2,$$

because it follows from

$$\begin{aligned} \frac{\pi^2}{6} &= \sum_{n=1}^{+\infty} \frac{1}{n^2} = \left\{ 1 + \frac{1}{2^2} + \frac{1}{2^4} + \cdots \right\} \sum_{p=0}^{+\infty} \frac{1}{(2p+1)^2} = \sum_{n=0}^{+\infty} \frac{1}{4^n} \sum_{p=0}^{+\infty} \frac{1}{(2p+1)^2} \\ &= \frac{4}{3} \sum_{p=0}^{+\infty} \frac{1}{(2p+1)^2}, \end{aligned}$$

that

$$\sum_{p=0}^{+\infty} \frac{1}{(2p+1)^2} = \frac{\pi^2}{8}.$$

Index

canonical kernel of finite rank, 27
 Cauchy-Schwarz inequality, 10, 16, 42, 47, 61
 differentiability in weak sense, 63
 fractional integration, 51
 general Hilbert-Schmidt operator, 5, 7, 27
 Hölder inequality, 49
 Hermitian kernel, 33, 37, 42
 Hilbert Schmidt norm, 7
 Hilbert-Schmidt operator, 5, 33
 iterated kernels, 13
 Legendre polynomial, 33
 Neumann series, 49
 Parseval's equation, 7
 trace, 15, 17, 27
 Volterra integral operator, 47

Please click the advert

It's only an opportunity if you act on it

IKEA.SE/STUDENT

© Inter IKEA Systems B.V. 2009